


SFB  
823

Nonparametric identification of  
endogenous and heterogeneous  
aggregate demand models:  
Complements, bundles and the  
market level

---

Fabian Dunker, Stefan Hoderlein,  
Hiroaki Kaido

Nr. 28/2015



Discussion Paper



# Nonparametric Identification of Endogenous and Heterogeneous Aggregate Demand Models: Complements, Bundles and the Market Level

Fabian Dunker\*

Ruhr-Universität Bochum

Stefan Hoderlein<sup>†</sup>

Boston College

Hiroaki Kaido<sup>‡</sup>

Boston University

June 20, 2015

## Abstract

This paper studies nonparametric identification in market level demand models for differentiated products. We generalize common models by allowing for the distribution of heterogeneity parameters (random coefficients) to have a nonparametric distribution across the population and give conditions under which the density of the random coefficients is identified. We show that key identifying restrictions are provided by (i) a set of moment conditions generated by instrumental variables together with an inversion of aggregate demand in unobserved product characteristics; and (ii) an integral transform (Radon transform) that maps the random coefficient density to the aggregate demand. This feature is shown to be common across a wide class of models, and we illustrate this by studying leading demand models. Our examples include demand models based on the multinomial choice (Berry, Levinsohn, Pakes, 1995), the choice of bundles of goods that can be substitutes or complements, and the choice of goods consumed in multiple units.

---

\*Ruhr-Universität Bochum, Department of Mathematics, Institute of Statistics, 44780 Bochum, Germany

<sup>†</sup>Department of Economics, Boston College, 140 Commonwealth Avenue, Chestnut Hill, MA 02467, USA, Email: stefan\_hoderlein@yahoo.com.

<sup>‡</sup>Department of Economics, Boston University, 270 Bay State Road, Boston, MA 02215, USA, Email: hkaido@bu.edu. We thank Steve Berry, Jeremy Fox, Amit Gandhi, Phil Haile, Kei Hirano, Arthur Lewbel, Marc Rysman, and Elie Tamer for their helpful comments. We also thank seminar and conference participants at Harvard, BU-BC joint mini-conference, the Demand Estimation and Modeling Conference 2013 and the Cowles Summer Conference 2014. Dunker gratefully acknowledges financial support from DFG FOR 916 and SFB 823. Kaido gratefully acknowledges financial support from NSF Grant SES-1357643.

# 1 Introduction

Modeling consumer demand for products that are bought in single or discrete units has a long and colorful history in applied Economics, dating back to at least the foundational work of McFadden (1974, 1981). While allowing for heterogeneity, much of the earlier work on this topic, however, was not able to deal with the fact that in particular the own price is endogenous. In a seminal paper that provides the foundation for much of contemporaneous work on discrete choice consumer demand, Berry, Levinsohn and Pakes (1994, BLP) have proposed a solution to the endogeneity problem. Indeed, this work is so appealing that it is not just applied in discrete choice demand and empirical IO, but also increasingly in many adjacent fields, such as health and urban economics, education and many others. From a methodological perspective, this line of work is quite different from traditional multivariate choice, as it uses data on the aggregate level and integrates out individual characteristics<sup>1</sup> to obtain a system of nonseparable equations. This system is then inverted for unobservables for which in turn a moment condition is then supposed to hold.

Descending in parts from the parametric work of McFadden (1974, 1981), BLP share many of its features, in particular (parametric) distributional assumptions, but also a linear random coefficients (RCs) structure for the latent utility. Not surprisingly, there is increasing interest in the properties of the model, in particular which features of the model are nonparametrically point identified, and how the structural assumptions affect identification of the parameters of interest. Why is the answer to these questions important? Because an empiricist working with this model wants to understand whether the results she obtained are a consequence of the specific parametric assumptions she invoked, or whether they are at least qualitatively robust. In addition, nonparametric identification provides some guidance on essential model structure and on data requirements, in particular about instruments. Finally, understanding the basic structure of the model makes it easier to understand how the model can be extended. Extensions of the BLP framework that are desirable are in particular to allow for consumption of bundles and multiple units of a product without modeling every choice as a new separate alternative.

We are not the first to ask the nonparametric identification question for market demand models. In a series of elegant papers, Berry and Haile (2011, 2013, BH henceforth) provide important answers to many of the identification questions. In particular, they establish conditions under which the “Berry inversion”, a core building block of the BLP model named after Berry (1994), which allows to solve for unobserved product characteristics, as well as the distribution

---

<sup>1</sup>There are extensions of the BLP framework that allow for the use of Microdata, see Berry, Levinsohn and Pakes (2004, MicroBLP). In this paper, we focus on the aggregate demand version of BLP, and leave an analogous work to MicroBLP for future research.

of a heterogeneous utility index are nonparametrically identified.

Our work complements this line of work in that we follow more closely the original BLP specification and assume in addition that the utility index has a linear random coefficients (RCs) structure. More specifically, we show how to nonparametrically identify the distribution of random coefficients in this framework. This result does not just close the remaining gap in the proof of nonparametric identification of the original BLP model, but is also important for applications because the distribution of random coefficients allows to characterize the distribution of the changes in welfare due to a change in regressors, in particular the own price (to borrow an analogy from the treatment effect literature, if we think of a price as a treatment, BH recover the treatment effect on the distribution, while we recover the distribution of treatment effects). For example, consider a change in the characteristics of a good. The change may be due to a new regulation, an improvement of the quality of a product, or an introduction of a new product. Knowledge of the random coefficient density would allow the researcher to calculate the distribution of the welfare change up to location and scale normalization. This allows one to answer various questions. For example, one may investigate whether the change gives rise to a Pareto improvement. This is possible because, with the distribution of the random coefficients being identified, one can track each individual's welfare before and after the change. If a change in one of the product characteristics is not Pareto improving, one can also calculate the proportion of individuals who would benefit from the change and therefore prefers the product with new characteristics.<sup>2</sup> Identification of the random coefficient distribution allows one to conduct various types of welfare analysis that are not possible by only identifying the demand function. Our focus therefore will be on the set of conditions under which one can uniquely identify the random coefficient distribution from the observed demand.

The arguments in establishing nonparametric identification of these changes are constructive and permit the construction of sample counterparts estimators, using theory in Hoderlein, Klemelä and Mammen (2010). From this theory it is well known that these estimators reveal that the random coefficients density is only weakly identified, suggesting that numerical instabilities and problems frequently reported and discussed in the BLP literature, e.g., Dube, Fox and Su (2013), are caused or aggravated by this feature of the model.

The second contribution in this paper is that we use the insights obtained from the identification results to extend the market demand framework to cover bundle choice (i.e., consume complementary goods together), as well as consumption of multiple units. Note that bundles and multiple purchases can in principle be accommodated within the BLP framework by

---

<sup>2</sup>Note that a simultaneous change in the product characteristic and price is allowed. Hence, one can also investigate how much price change can be made to compensate for a change (e.g. downgrading of a feature) in one of the product characteristics to let a certain fraction of individuals receive a non-negative utility change, i.e.  $P(\Delta U_{ijt} \geq 0) \geq \tau$  for some prespecified  $\tau \in [0, 1]$ , where  $\Delta U_{ijt}$  denotes the utility change.

treating them as separate alternatives. However, this is not parsimonious as the number of alternatives increases rapidly and with it the number of unobserved product characteristics, making the system quickly intractable. To fix ideas, suppose there were two goods, say good A and B. First, we allow for the joint consumption of goods A and B, and second, we allow for the consumption of several units of either A and/or B, without labeling it a separate alternative. We model the utility of each bundle as a combination of the utilities for each good and an extra utility from consuming the bundle. This structure in turn implies that the dimension of the unobservable product characteristic equals the number of goods  $J$  instead of the number of bundles. There are three conclusions we draw from this contribution: first, depending on the type of model, the data requirements vary. In particular, to identify all structural parts of the model, in, say, the model on bundle choice, market shares are not the correct dependent variable any more. Second, depending on the object of interest, the data requirements and assumptions may vary depending on whether we want to just recover demand elasticities, or the entire distribution of random coefficients. Third, the parsimonious features of the structural model result in significant overidentification of the model, which opens up the way for specification testing, and efficient estimation. As in the classical BLP setup, in all setups we may use the identification argument to propose a nonparametric sample counterpart estimators, but we also use the insights obtained to propose a parametric estimator for models where there had not been an estimator before.

**Related literature:** as discussed above, this paper is closely related to both the original BLP line of work (Berry, Levinsohn and Pakes (1994, 2004)), as well as to the recent identification analysis of Berry and Haile (2011, 2013). Because of its generality, our approach also provides identification analysis for the “pure characteristics” model of Berry and Pakes (2007), see also Ackerberg, Benkard, Berry and Pakes (2007) for an overview. Other important work in this literature that is completely or partially covered by the identification results in this paper include Petrin (2002) and Nevo (2001). Moreover, from a methodological perspective, we note that BLP continues a line of work that emanates from a broader literature which in turn was pioneered by McFadden (1974, 1981); some of our identification results extend therefore beyond the specific market demand model at hand. Other important recent contributions in discrete choice demand include Armstrong (2013) and Moon, Shum, and Weidner (2013). Less closely related is the literature on hedonic models, see Heckman, Matzkin and Nesheim (2010), and references therein.

In addition to this line of work, we also share some commonalities with the work on bundle choice in IO, most notably Gentzkow (2007), and Fox and Lazzati (2013). For some of the examples discussed in this paper, we use Gale-Nikaido inversion results, which are related to arguments in Berry, Gandhi and Haile (2013). Because of the GMM type endogeneity, our

approach also relates to nonparametric IV, in particular to Newey and Powell (2003), Andrews (2011), and Dunker, Florens, Hohage, Johannes, and Mammen (2014). Finally, our arguments are related to the literature on random coefficients in discrete choice model, see Ichimura and Thompson (1995), Gautier and Kitamura (2013), Dunker, Hoderlein and Kaido (2013), Fox and Gandhi (2012), and Matzkin (2012). Since we use the Radon transform introduced by Hoderlein, Klemelä and Mammen (2010, HKM) into Econometrics, possibly in conjunction with tensor products as in Dunker, Hoderlein and Kaido (2013), this work is particularly close to the literature that uses the Radon transform, in particular HKM and Gautier and Hoderlein (2013).

**Structure of the paper:** The second section lays out preliminaries we require for our main result: We first introduce the class of models and detail the structure of our two main setups. Still in the same section, for completeness we quickly recapitulate the results of Berry and Haile (2013) concerning the identification of structural demands, adapted to our setup. The third section contains the key novel result in this paper, the nonparametric (point-)identification of the distribution of random coefficients in the BLP setup. The fourth section contains various extensions: We discuss the identification in the bundles case, including how the structural demand identification results of Berry and Haile (2013) have to be adapted, but again focusing on the random coefficients density. As another set of extensions, we discuss the multiple units case, and the pure characteristics model that does not contain a market-product-individual specific (“logit”) error. Finally, we discuss how full independence assumptions may be utilized to increase the strength of identification, in particular in the identification of structural demands. The fifth section discusses estimation. The objective here is twofold, first we sketch how a nonparametric sample counterparts estimator that utilizes the insights of the identification sections could be constructed, and we propose a simple parametric estimator for the bundles model which we believe to be relevant for applications. We end with an outlook.

## 2 Preliminaries

### 2.1 Model

We begin with a setting where a consumer faces  $J \in \mathbb{N}$  products and an outside good which is labeled good 0. Throughout, we index individuals by  $i$ , products by  $j$  and markets by  $t$ . We use upper-case letters, e.g.  $X_{jt}$ , for random variables (or vectors) that vary across markets and lower-case letters, e.g.  $x_j$ , for particular values the random variables (vectors) can take. In addition, we use letters without a subscript for products e.g.  $X_t$  to represent vectors e.g.  $(X_{1t}, \dots, X_{Jt})$ . For individual  $i$  in market  $t$ , the (indirect) utility from consuming

good  $j$  depends on its (log) price  $P_{jt}$ , a vector of observable characteristics  $X_{jt} \in \mathbb{R}^{d_x}$ , and an unobservable scalar characteristic  $\Xi_{jt} \in \mathbb{R}$ . Following Berry, Levinsohn and Pakes (1995), we model the utility from consuming good  $j$  using the linear random coefficient specification:

$$U_{ijt}^* \equiv X_{jt}'\beta_{it} + \alpha_{it}P_{jt} + \Xi_{jt} + \epsilon_{ijt}, \quad j = 1, \dots, J, \quad (2.1)$$

where  $(\alpha_{it}, \beta_{it})' \in \mathbb{R}^{d_x+1}$  is a vector of random coefficients on the product characteristics, which varies across individuals.  $\epsilon_{ijt}$  is an additive stochastic taste shifter. In what follows, we call  $\epsilon_{it} \equiv (\epsilon_{i1t}, \dots, \epsilon_{iJt})'$  “tastes for products” following Berry and Pakes (2007). Throughout, we assume that  $X_{jt}$  is exogenous, while  $P_{jt}$  can be correlated with the unobserved product characteristic  $\Xi_{jt}$  in an arbitrary way. Without loss of generality, we normalize the utility from the outside good to 0. This mirrors the setup considered in BH (2013).

Throughout we think of a large sample of individuals as *iid* copies of this population model. The random coefficients  $\theta_{it} \equiv (\alpha_{it}, \beta_{it}, \epsilon_{i1t}, \dots, \epsilon_{iJt})'$  vary across individuals in any given market (or, alternatively, have a distribution in any given market in the population), while the product characteristics vary solely across markets. These coefficients are assumed to follow a distribution with a density function  $f_\theta$  with respect to Lebesgue measure, i.e., be continuously distributed. This density is assumed to be common across markets, and is therefore not indexed by  $t$ . As we will show, an important aspect of our identification argument is that, once the demand function is identified, one may recover  $\Xi_t$  from the market shares and other product characteristics  $(X_t, P_t)$ . Then, by creating exogenous variations in the product characteristics and exploiting the linear random coefficients structure, one may trace out the distribution  $f_\theta$  of the preference that is common across markets. We note that we can allow for the coefficients  $(\alpha_{it}, \beta_{it})$  to be alternative  $j$  specific, and will indeed do so below. However, parts of the analysis will subsequently change, and we start out with the more common case where the coefficients are the same across  $j$ .

Having specified the model on individual level, the outcomes of individual decisions are then aggregated in every market. The econometrician observes exactly these market level outcomes  $S_{l,t}$ , where  $l$  belongs to some index set denoted by  $\mathbb{L}$ . Below, we give two examples. The first example is the setting of BLP, which is our main focus where individuals choose a single good out of multiple products.

**Example 1** (BLP). Each individual chooses the product that maximizes her utility out of  $J \in \mathbb{N}$  products. Hence, product  $j$  is chosen if

$$U_{jt}^* > U_{kt}^*, \quad \forall k \neq j. \quad (2.2)$$

The demand for good  $j$  in market  $t$  is obtained by aggregating the individual demand with



respect to the distribution of individual preferences.

$$\begin{aligned} \varphi_j(X_t, P_t, \Xi_t) = & \int 1\{X'_{jt}b + aP_{jt} + e_j > -\Xi_{jt}\}1\{(X_{jt} - X_{1t})'b + a(P_{jt} - P_{1t}) + (e_j - e_1) > -(\Xi_{jt} - \Xi_{1t})\} \\ & \cdots 1\{(X_{jt} - X_{Jt})'b + a(P_{jt} - P_{Jt}) + (e_j - e_J) > -(\Xi_{jt} - \Xi_{Jt})\}f_\theta(b, a, e)d\theta, \end{aligned} \quad (2.3)$$

for  $j = 1, \dots, J$ , while the aggregate demand for good 0 is given by

$$\varphi_0(X_t, P_t, \Xi_t) = \int 1\{X'_{1t}b + aP_{1t} + e_1 < -\Xi_{1t}\} \cdots 1\{X'_{Jt}b + aP_{Jt} + e_J < -\Xi_{Jt}\}f_\theta(b, a, e)d\theta, \quad (2.4)$$

where  $(b, a, e_1, \dots, e_J)$  are placeholders for the random coefficients  $\theta_{it} = (\beta_{it}, \alpha_{it}, \epsilon_{i1t}, \dots, \epsilon_{iJt})$ . The researcher then observes the market shares of products  $S_{lt} = \varphi_l(X_t, P_t, \Xi_t)$ ,  $l \in \mathbb{L}$ , where  $\mathbb{L} = \{0, 1, \dots, J\}$ .

The second example considers choice of bundles.

**Example 2 (Bundles).** Each individual faces  $J = 2$  products and decides whether or not to consume a single unit of each of the products. There are therefore four possible combinations  $(Y_1, Y_2)$  of consumption units, which we call *bundles*. In addition to the utility from consuming each good as in (2.1), the individuals gain additional utility (or disutility)  $\Delta_{it}$  if the two goods are consumed simultaneously. Here,  $\Delta_{it}$  is also allowed to vary across individuals. The utility  $U_{i,(Y_1, Y_2),t}^*$  from each bundle is therefore specified as follows:

$$\begin{aligned} U_{i,(0,0),t}^* &= 0, \\ U_{i,(1,0),t}^* &= X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t}, \quad U_{i,(0,1),t}^* = X'_{2t}\beta_{it} + \alpha_{it}P_{2t} + \Xi_{2t} + \epsilon_{i2t}, \\ U_{i,(1,1),t}^* &= X'_{1t}\beta_{it} + X'_{2t}\beta_{it} + \alpha_{it}P_{1t} + \alpha_{it}P_{2t} + \Xi_{1t} + \Xi_{2t} + \epsilon_{i1t} + \epsilon_{i2t} + \Delta_{it}, \end{aligned} \quad (2.5)$$

Each individual chooses a bundle that maximizes her utility. Hence, bundle  $(y_1, y_2)$  is chosen when  $U_{i,(y_1, y_2),t}^* > U_{i,(y'_1, y'_2),t}^*$  for all  $(y'_1, y'_2) \neq (y_1, y_2)$ . For example, bundle  $(1, 0)$  is chosen if

$$\begin{aligned} X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t} > 0, \quad \text{and} \quad X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t} > X'_{2t}\beta_{it} + \alpha_{it}P_{2t} + \Xi_{2t} + \epsilon_{i2t}, \quad \text{and} \\ X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t} > X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t} + X'_{2t}\beta_{it} + \alpha_{it}P_{2t} + \Xi_{2t} + \epsilon_{i2t} + \Delta_{it}. \end{aligned} \quad (2.6)$$

Suppose the random coefficients  $\theta_{it} = (\beta'_{it}, \alpha_{it}, \Delta_{it}, \epsilon_{i1t}, \epsilon_{i2t})$  have a joint density  $f_\theta$ . The aggregate structural demand for  $(1, 0)$  can then be obtained by integrating over the set of individuals

satisfying (2.6) with respect to the distribution of the random coefficients:

$$\begin{aligned} \varphi_{(1,0)}(X_t, P_t, \Xi_t) = & \int 1\{X'_{1t}b+aP_{1t}+e_1 > -\Xi_{1t}\}1\{(X_{1t}-X_{2t})'b+a(P_{1t}-P_{2t})+(e_1-e_2) > \Xi_{2t}-\Xi_{1t}\} \\ & \times 1\{X'_{2t}b+aP_{2t}+e_2+\Delta < -\Xi_{2t}\}f_\theta(b, a, \Delta, e)d\theta. \end{aligned} \quad (2.7)$$

The aggregate demand on other bundles can be obtained similarly. The econometrician then observes a vector of aggregate demand on the bundles:  $S_{l,t} = \varphi_l(X_t, P_t, \Xi_t), l \in \mathbb{L}$  where  $\mathbb{L} \equiv \{(0, 0), (1, 0), (0, 1), (1, 1)\}$ .

In Examples 2, we assume that the econometrician observes the aggregate demand for all the respective bundles. We emphasize this point as it changes the data requirement, and an interesting open question arises about what happens if these requirements are not met. Example of data sets that would satisfy these requirements are when 1. individual observations are collected through direct survey or scanner data on individual consumption (in every market), 2. aggregate variables (market shares) are collected, but augmented with a survey that asks individuals whether they consume each good separately or as a bundle. 3. Finally, another possible data source are producer's direct record of sales of bundles, provided each bundles are recorded separately (e.g., when they are sold through promotional activities). When discussing Example 2 (and Example 3 in Section 4), we henceforth tacitly assume to have access to such data in principle.

## 2.2 Structural Demand

The first step toward identification of  $f_\theta$  is to use a set of moment conditions generated by instrumental variables to identify the aggregate demand function  $\varphi$ . Following BH (2013), we partition the covariates as  $X_{jt} = (X_{jt}^{(1)}, X_{jt}^{(2)}) \in \mathbb{R} \times \mathbb{R}^{d_x-1}$ , and make the following assumption.

**Assumption 2.1.** *The coefficient  $\beta_{ij}^{(1)}$  on  $X_{jt}^{(1)}$  is non-random for all  $j$  and is normalized to 1.*

Assumption 2.1 requires that at least one coefficient on the covariates is non-random. Since we may freely choose the scale of utility, we normalize the utility by setting  $\beta_{ij}^{(1)} = 1$  for all  $j$ . Under Assumption 2.1, the utility for product  $j$  can be written as  $U_{jt}^* = X_{jt}^{(2)'}\beta_{ij}^{(2)} + \alpha_{ij}P_{jt} + \epsilon_{ijt} + D_{jt}$ , where  $D_{jt} \equiv X_{jt}^{(1)} + \Xi_{jt}$  is the part of the utility that is common across individuals. Assumption 2.1 (i) is arguably strong but will provide a way to obtain valid instruments required to identify the structural demand (see BH, 2013, Section 7 for details). Under this assumption,  $U_{ijt}^*$  is strictly increasing in  $D_{jt}$  but unaffected by  $D_{it}$  for all  $i \neq j$ . In Example 1, together with a mild regularity condition, this is sufficient for inverting the demand system to obtain  $\Xi_t$  as a function of the market shares  $S_t$ , price  $P_t$ , and exogenous

covariates  $X_t$  (Berry, Gandhi, and Haile, 2013). In what follows, we redefine the aggregate demand as a function of  $(X_t^{(2)}, P_t, D_t)$  instead of  $(X_t, P_t, \Xi_t)$  by  $\phi(X_t^{(2)}, P_t, D_t) \equiv \varphi(X_t, P_t, \Xi_t)$ , where  $X_t = (X_t^{(1)}, X_t^{(2)})$  and  $D_t = \Xi_t + X_t^{(1)}$  and make the following assumption

**Assumption 2.2.** *For some subset  $\tilde{\mathbb{L}}$  of  $\mathbb{L}$  whose cardinality is  $J$ , there exists a unique function  $\psi : \mathbb{R}^{J \times (d_X - 1)} \times \mathbb{R}^J \times \mathbb{R}^J \rightarrow \mathbb{R}^J$  such that  $D_{jt} = \psi_j(X_t^{(2)}, P_t, \tilde{S}_t)$  for  $j = 1, \dots, J$ , where  $\tilde{S}_t$  is a subvector of  $S_t$ , which stacks the components of  $S_t$  whose indices belong to  $\tilde{\mathbb{L}}$ .*

Under Assumption 2.2, we may write

$$\Xi_{jt} = \psi_j(X_t^{(2)}, P_t, \tilde{S}_t) - X_{jt}^{(1)}. \quad (2.8)$$

This can be used to generate moment conditions in order to identify the aggregate demand function.

**Example 1** (BLP, continued). Let  $\tilde{\mathbb{L}} = \{1, \dots, J\}$ . In this setting, the inversion discussed above is the standard Berry inversion. A key condition for the inversion is that the products are *connected substitutes* (Berry, Gandhi, and Haile (2013)). The linear random coefficient specification as in (2.1) is known to satisfy this condition. Then, Assumption 2.2 follows.

In Example 2, one may employ an alternative inversion strategy to obtain  $\psi$  in (2.8) with  $\tilde{\mathbb{L}} = \{(1, 0), (1, 1)\}$  or  $\tilde{\mathbb{L}} = \{(0, 0), (0, 1)\}$ . To keep a tight focus on the BLP example, we defer details on this case to Section 4.

The inverted system in (2.8), together with the following assumption, yields a set of moment conditions the researcher can use to identify the structural demand.

**Assumption 2.3.** *There is a vector of instrumental variables  $Z_t \in \mathbb{R}^{d_Z}$  such that (i)  $E[\Xi_{jt}|Z_t, X_t] = 0$ , a.s.; (ii) for any  $B : \mathbb{R}^{Jk_2} \times \mathbb{R}^J \times \mathbb{R}^J \rightarrow \mathbb{R}$  with  $E[|B(X_t^{(2)}, P_t, \tilde{S}_t)|] < \infty$ , it holds that*

$$E[B(X_t^{(2)}, P_t, \tilde{S}_t)|Z_t, X_t] = 0 \implies B(X_t^{(2)}, P_t, \tilde{S}_t) = 0, \text{ a.s.}$$

Assumption 2.3 (i) is a mean independence assumption on  $\Xi_{jt}$  given a set of instruments  $Z_t$ , which also normalizes the location of  $\Xi_{jt}$ . Assumption 2.3 (ii) is a completeness condition, which is common in the nonparametric IV literature, see BH (2013) for a detailed discussion. However, the role it plays here is slightly different, as the moment condition leads to an integral equation which is different from nonparametric IV (Newey & Powell, 2003), and more resembles GMM. As such, the construction of a sample counterpart estimator is less clear. In Section 4.5, we discuss an approach based on a strengthening of the mean independence condition to full independence. In case such a strengthening is economically palatable, we still retain the sum  $X_{jt}^{(1)} + \Xi_{jt}$ , which has a closer analogy to a dependent variable in nonparametric IV.

Given Assumption 2.3 and (2.8), the unknown function  $\psi$  can be identified through the following conditional moment restrictions:

$$E[\psi_j(X_t^{(2)}, P_t, S_t) - X_{jt}^{(1)} | Z_t, X_t] = 0, \quad j = 1, \dots, J. \quad (2.9)$$

We here state this result as a theorem.

**Theorem 2.1.** *Suppose Assumptions 2.1-2.3 hold. Then,  $\psi$  is identified.*

Once  $\psi$  is identified, the structural demand  $\phi$  can be identified nonparametrically in Examples 1 and 2.

**Example 1** (BLP, continued). Recall that  $\psi$  is a unique function such that

$$S_{jt} = \phi_j(X_t^{(2)}, P_t, D_t), \quad j = 1, \dots, J \quad \Leftrightarrow \quad \Xi_{jt} = \psi_j(X_t^{(2)}, P_t, \tilde{S}_t) - X_{jt}^{(1)}, \quad j = 1, \dots, J, \quad (2.10)$$

where  $\tilde{S}_t = (S_{1t}, \dots, S_{Jt})$ . Hence, the structural demand  $(\phi_1, \dots, \phi_J)$  is identified by Theorem 2.1 and the equivalence relation above. In addition,  $\phi_0$  is identified through the identity:  $\phi_0 = 1 - \sum_{j=1}^J \phi_j$ .

**Example 2** (Bundles, continued). Let  $\tilde{\mathbb{L}} = \{(1, 0), (1, 1)\}$ .  $\psi$  is then a unique function such that

$$S_{lt} = \phi_l(X_t^{(2)}, P_t, D_t), \quad l \in \tilde{\mathbb{L}} \quad \Leftrightarrow \quad \Xi_{jt} = \psi_j(X_t^{(2)}, P_t, \tilde{S}_t) - X_{jt}^{(1)}, \quad j = 1, 2, \quad (2.11)$$

where  $\tilde{S}_t = (S_{(1,0),t}, S_{(1,1),t})$ . Theorem 2.1 and the equivalence relation above then identify the demand for bundles (1, 0) and (1, 1). This, therefore, only identifies subcomponents of  $\phi$ . Although these subcomponents are sufficient for recovering the random coefficient density, one may also identify the rest of the subcomponents by taking  $\tilde{\mathbb{L}} = \{(0, 0), (0, 1)\}$  and applying Theorem 2.1 again.

### 3 Identification of the Random Coefficient Density in the BLP model

This section contains the main innovation in this paper: We establish that the density of random coefficients in a BLP setup is nonparametrically identified. Our strategy for identification of the random coefficient density is to construct a function from the structural demand, which

is related to the density through an integral transform known as the *Radon transform*. More precisely, we construct a function  $\Phi(w, u)$  such that

$$\frac{\partial \Phi(w, u)}{\partial u} = \mathcal{R}[f](w, u), \quad (3.1)$$

where  $f$  is the density of interest,  $w$  is a unit vector in  $\mathbb{R}^q$  where  $q$  is the dimension of the random coefficients, and  $u \in \mathbb{R}$  is a scalar. In what follows, we let  $\mathbb{S}^q \equiv \{v \in \mathbb{R}^q : \|v\| = 1\}$  denote the unit sphere in  $\mathbb{R}^q$ .  $\mathcal{R}$  is the Radon transform defined pointwise by

$$\mathcal{R}[f](w, u) = \int_{P_{w,u}} f(v) d\mu_{w,u}(v). \quad (3.2)$$

where  $P_{w,u}$  denotes the hyperplane  $\{v \in \mathbb{R}^q : v'w = u\}$ , and  $\mu_{w,u}$  is the Lebesgue measure on  $P_{w,u}$ . (See for example Helgason (1999) for details on the properties of the Radon transform including its injectivity.) Our identification strategy is constructive and will therefore suggest a natural nonparametric estimator. Applications of the Radon transform to random coefficients models have been studied in Hoderlein, Klemelä, and Mammen (2010), and Gautier and Hoderlein (2013).

Throughout, we maintain the following assumption.

**Assumption 3.1.** (i) For all  $k \in \{1, \dots, J\}$ ,  $(X_{kt}^{(2)}, P_{kt}, D_{kt})$  are absolutely continuous with respect to Lebesgue measure on  $\mathbb{R}^{dx-1} \times \mathbb{R} \times \mathbb{R}$ ; (ii) the random coefficients  $\theta$  are independent of  $(X_t, P_t, D_t)$ .

Assumption 3.1 (i) requires that  $(X_{it}^{(2)}, P_{it}, D_{it})$  are continuously distributed for all  $i$ . By Assumption 3.1 (ii), we assume that the covariates  $(X_t, P_t, D_t)$  are exogenous to the individual heterogeneity and  $f_\theta$  has a bounded support. These conditions are used to invert the Radon transform in (3.2).

We now discuss the construction of  $\Phi$  for the BLP model. Recall that the demand for good  $j$  with the product characteristics  $(X_t, P_t, \Xi_t)$  is as given in (2.3). Since  $D_t = X_t^{(1)} + \Xi_t$ , the demand in market  $t$  with  $(X_t^{(2)}, P_t, D_t) = (x^{(2)}, p, \delta)$  is given by:

$$\begin{aligned} \phi_j(x^{(2)}, p, \delta) &= \int 1\{x_j^{(2)'}b^{(2)} + ap_j + e_j > -\delta_j\} 1\{(x_j^{(2)} - x_1^{(2)})'b^{(2)} + a(p_j - p_1) + (e_j - e_1) > -(\delta_j - \delta_1)\} \\ &\quad \cdots 1\{(x_j^{(2)} - x_J^{(2)})'b^{(2)} + a(p_j - p_J) + (e_j - e_J) > -(\delta_j - \delta_J)\} f_\theta(b^{(2)}, a, e) d\theta. \end{aligned} \quad (3.3)$$

If  $D_{kt}, k \neq j$  have a large support conditional on  $(X_t^{(2)}, P_t, D_{jt})$ , one may consider letting  $D_{kt} \rightarrow -\infty$  for all  $k \neq j$ . The demand for good  $j$  in such a setting is  $\phi_j(x^{(2)}, p, \delta) = \int 1\{x_j^{(2)'}b^{(2)} + ap_j + e_j > -\delta_j\} f_{\vartheta_j}(b^{(2)}, a, e_j) d\vartheta_j$ , where  $f_{\vartheta_j}$  denotes the joint density of the subvector  $\vartheta_{ijt} \equiv (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{ijt})$  of the random coefficients.

Under the condition we provide below, we may construct a function  $\tilde{\Phi}_j$ , which can be written in general as  $\tilde{\Phi}_j(X_j^{(2)}, P_j, D_j) = \int 1\{X_j^{(2)'}b^{(2)} + aP_j + e_j < -D_j\} f_{\vartheta_j}(b^{(2)}, a, e_j) d\vartheta_j$ . Let  $w \equiv (x_j^{(2)}, p_j, 1) / \|(x_j^{(2)}, p_j, 1)\|$  and  $u \equiv \delta_j / \|(x_j^{(2)}, p_j, 1)\|$ . Define

$$\begin{aligned} \Phi(w, u) &\equiv \tilde{\Phi}_j\left(\frac{x_j^{(2)}}{\|(x_j^{(2)}, p_j, 1)\|}, \frac{p_j}{\|(x_j^{(2)}, p_j, 1)\|}, \frac{\delta_j}{\|(x_j^{(2)}, p_j, 1)\|}\right) \\ &= \tilde{\Phi}_j(x_j^{(2)}, p_j, \delta_j), \quad (x_j^{(2)}, p_j, \delta_j) \in \text{supp}(X_{jt}^{(2)}, P_{jt}, D_{jt}), \end{aligned} \quad (3.4)$$

where the second equality holds because normalizing the scale of  $(x_j^{(2)}, p_j, \delta_j)$  does not change the value of  $\tilde{\Phi}_j$ .  $\Phi$  then satisfies

$$\begin{aligned} \Phi(w, u) &= - \int 1\{w'\vartheta_j < -u\} f_{\vartheta_j}(b^{(2)}, a, e_j) d\vartheta_j \\ &= - \int_{-\infty}^{-u} \int_{P_{w,r}} f_{\vartheta_j}(b^{(2)}, a, e_j) d\mu_{w,r}(b^{(2)}, a, e_j) dr = - \int_{-\infty}^{-u} \mathcal{R}[f_{\vartheta_j}](w, r) dr, \end{aligned} \quad (3.5)$$

Hence, by taking a derivative with respect to  $u$ , we may relate  $\Phi$  to the random coefficient density through the Radon transform:

$$\frac{\partial \Phi(w, u)}{\partial u} = \mathcal{R}[f_{\vartheta_j}](w, u). \quad (3.6)$$

Note that since the structural demand  $\phi$  is identified by Theorem 2.1,  $\Phi$  is nonparametrically identified as well. Hence, Eq. (3.6) gives an operator that maps the random coefficient density to an object identified by the moment condition studied in the previous section. To construct  $\Phi$  described above and to invert the Radon transform, we formally make the following assumptions.

**Assumption 3.2.** (i) For each  $j$ , the joint distribution of  $\{D_{kt}, k \neq j\}$  conditional on  $(X_t, P_t, D_{jt})$  has a full support a.s.; (ii)  $\bigcup_{j=1}^J \text{supp}(X_{jt}^{(2)}, P_{jt}, D_{jt}) = \mathbb{R}^{d_x-1} \times \mathbb{R} \times \mathbb{R}$ .

Assumption 3.2 (i) requires that one may vary  $D_{kt}$  for  $k \neq j$  on a large support so that the demand for product  $j$  is determined through its choice between product  $j$  and the outside good. This identification argument therefore uses a “thin” (lower-dimensional) subset of the support of the covariates, which is due to the presence of the tastes for products  $\epsilon_{ijt}, j = 1, \dots, J$  in the model. On the other hand, if the researcher uses a model without the tastes for products (called the pure characteristics model), one can achieve identification without relying on a lower-dimensional subset of the support of the covariates. We will revisit this point in Section 4.3. Assumption 3.2 (ii) requires that the union of the supports of the product characteristics  $(X_{jt}^{(2)}, P_{jt}, D_{jt}), j = 1, \dots, J$  jointly span the full support. This includes as a special case the

setting where there exists a “special product” whose product characteristics  $(X_{jt}^{(2)}, P_{jt}, D_{jt})$  has a full support. Even if such a product does not exist, identification of the random coefficient density is possible as long as the full support condition can be met by combining the supports of all products. This means that our identification strategy can use the variations in the product characteristics across multiple products.<sup>3</sup>

Under the conditions given in the theorem below, inverting the Radon transform in (3.2) identifies  $f_{\vartheta_j}$ . If one is interested in the joint density of the coefficients on the product characteristics  $(\beta_{it}^{(2)}, \alpha_{it})$ , one may stop here as marginalizing  $f_{\vartheta_j}$  gives the desired density. The joint distribution of all coefficients including the tastes for products can be identified under an additional assumption. We state this result in the following theorem.

**Theorem 3.1.** *Suppose Assumptions 2.1-3.2 hold. Suppose that the conditional distribution of  $\epsilon_{ijt}$  given  $(\beta_{it}^{(2)}, \alpha_{it})$  is identical for  $j = 1, \dots, J$ . Then, (i)  $f_{\vartheta_j}$  is identified in Example 1, where  $\vartheta_{ijt} = (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{ijt})$ ; (ii) If  $(\epsilon_{i1t}, \dots, \epsilon_{iJt})$  are independently distributed (across  $j$ ) conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$ ,  $f_{\theta}$  is also identified.*

Several remarks are in order.

**Remark 3.1.** The identical distribution assumption on  $\{\epsilon_{ijt}\}_{j=1}^J$  in Theorem 3.1 is compatible with commonly used utility specifications and can also be relaxed at the cost of a stronger support condition on the product characteristics. In applications, it is often assumed that the utility of product  $j$  is

$$U_{ijt}^* = \beta_{it}^0 + \tilde{X}_{jt}'\beta_{it} + \alpha_{it}P_{jt} + \Xi_{jt} + \tilde{\epsilon}_{ijt}, \quad (3.7)$$

where  $\tilde{X}_{jt}$  is a vector of non-constant product characteristics,  $\beta_{it}^0$  is an individual specific intercept, which measures the utility difference between inside goods and the outside good, and  $\tilde{\epsilon}_{ijt}$  is a mean zero error that follows the Type-I extreme value distribution. The requirement that  $\epsilon_{ijt} = \beta_{it}^0 + \tilde{\epsilon}_{ijt}$  are i.i.d. across  $j$  (conditional on  $(\beta_{it}, \alpha_{it})$ ) can be met if  $\tilde{\epsilon}_{ijt}$  are i.i.d. across  $j$ .

If for each  $j$ ,  $(X_{jt}^{(2)}, P_{jt}, D_{jt})$  has a full support, one can drop the identical distribution assumption on  $\{\epsilon_{ijt}\}_{j=1}^J$ . This is because one can identify  $f_{\vartheta_j}$  for all  $j = 1, \dots, J$  by inverting the Radon transform in (3.6) repeatedly. This in turn implies that the distribution of  $\epsilon_{ijt}$  conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$  is identified for each  $j$ . If one assumes  $(\epsilon_{i1t}, \dots, \epsilon_{iJt})$  are independent of each other conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$ , the joint distribution of  $(\epsilon_{i1t}, \dots, \epsilon_{iJt})$  conditional on

---

<sup>3</sup>More precisely, the Radon transform  $\mathcal{R}[f_{\vartheta_j}](w, u)$  gives  $f_{\vartheta_j}$ 's integral along each hyperplane  $P_{w,u} = \{v \in \mathbb{R}^{d_{\vartheta}} : v'w = u\}$  defined by the *angle*  $w = (x_j^{(2)}, p_j, 1)/\|(x_j^{(2)}, p_j, 1)\|$  and *offset*  $u = \delta_j/\|(x_j^{(2)}, p_j, 1)\|$ . For recovering  $f_{\vartheta_j}$  from its Radon transform, one needs exogenous variations in both. Our proof uses the fact that varying  $w$  over the hemisphere  $\mathbb{H}_+ \equiv \{w = (w_1, w_2, \dots, w_{d_{\vartheta_j}}) \in \mathbb{S}^{d_{\vartheta_j}-1} : w_{d_{\vartheta_j}} \geq 0\}$  and  $u$  over  $\mathbb{R}$  suffices to recover  $f_{\vartheta_j}$ .

$(\beta_{it}^{(2)}, \alpha_{it})$  is also identified, which in turn implies that the joint distribution  $f_\theta$  of all coefficients is identified.

**Remark 3.2.** Theorems 2.1 and 3.1 shed light on the roles played by the key features of the BLP-type demand model: the invertibility of the demand system, instrumental variables, and the linear random coefficients specification. In Theorem 2.1, the invertibility and instrumental variables play key roles in identifying the demand. Once the demand is identified, one may “observe” the vector  $(X_t^{(2)}, P_t, D_t)$  of covariates. This is possible because the invertibility of demand allows one to recover the unobserved product characteristics  $\Xi_t$  from the market shares  $S_t$  (together with other covariates). One may then vary  $(X_t^{(2)}, P_t, D_t)$  across markets in a manner that is exogenous to the individual heterogeneity  $\theta_{it}$ . Theorem 3.1 shows this exogenous variation combined with the linear random coefficients specification then allows one to trace out the distribution of  $\theta_{it}$ .

**Remark 3.3.** Our identification result reveals the nature of the BLP-type demand model. A positive aspect of our result is that the preference is nonparametrically identified if one observes rich variations in the consumers’ choice sets (represented by  $(X_{jt}^{(2)}, P_{jt}, D_{jt})$ ) across markets. On the other hand, if the product characteristics have limited variations, the identifying power of the model on the distribution of preferences may be limited. In particular, identification is not achieved only with discrete covariates. Hence, for such settings, one needs to augment the model structure with a parametric specification or independence assumptions.

There are several potential ways to extend our result. One direction is to relax the support condition. In practice, one may not be able to vary  $w$  in (3.6) flexibly when  $(X_{jt}^{(2)}, P_{jt}, D_{jt})$  has a limited support. Even in such a setting, identification of  $f_\theta$  is still possible under an additional assumption on the distribution of the random coefficients. One such condition is as follows:

**Assumption 3.3.** *All the absolute moments of each component of  $\theta_{it}$  are finite, and for any fixed  $z \in \mathbb{R}_+$ ,  $\lim_{l \rightarrow \infty} \frac{z^l}{l!} (E[|\theta_{it}^{(1)}|^l] + \dots + E[|\theta_{it}^{(d_\theta)}|^l]) = 0$ .*

Under Assumption 3.3, the characteristic function  $w \mapsto \varphi_\theta(tw)$  of  $\theta$  (a key element of the Radon inversion) is uniquely determined by its restriction to a non-empty full dimensional subset of  $\mathbb{S}^{d_\theta}$ .<sup>4</sup> Hence,  $f_{\vartheta_j}$  can still be identified if one may vary  $w$  on a non-empty full dimensional subset. For example, if the union of the supports of  $(X_{jt}^{(2)}, P_{jt}, D_{jt}), j = 1, \dots, J$  contains an open ball in  $\mathbb{R}^{d_x-1} \times \mathbb{R} \times \mathbb{R}$ , this is sufficient for the identification of  $f_{\vartheta_j}$ . Alternatively, if the support of the random coefficients is compact, one may employ another integral transform, the limited-angle Radon transform, which is also known to be injective. Finally, another interesting

---

<sup>4</sup>This follows from analytic continuation. See Hoderlein, Holzman, and Meister (2014) and Masten (2014) for details.



direction would be to conduct partial identification analysis on functionals of  $f_\theta$ , while imposing weak support restrictions.

## 4 Extensions

Below, we show that our strategy set forth in the previous section can also be applied to extended models that share the key features of the BLP-type demand model. We further discuss identification of the random coefficient density in pure characteristics demand models, the analysis of the case in which random coefficients are alternative specific, and an alternative approach to nonparametrically identifying the demand under a full independence assumption.

### 4.1 Bundle choice (Example 2)

We consider an alternative procedure for inverting the demand in Example 2. This is because this example (and also the example in the next section) has a specific structure. We note that the inversion of Berry, Gandhi, and Haile (2013) can still be applied to bundles if one treats each bundle as a separate good and recast the bundle choice problem into a standard multinomial choice problem. However, as can be seen from (2.5), Example 2 has the additional structure that the utility of a bundle is the combination of the utilities for each good and extra utilities, and hence the model does not involve any bundle specific unobserved characteristic. This structure in turn implies that the dimension of the unobservable product characteristic  $\Xi_t$  equals the number of goods  $J$ , while the econometrician observes  $\dim(S) = \prod_{j=1}^J (d_j + 1)$  aggregate choice probabilities over bundles, where  $d_j$  is the maximum number of consumption units allowed for each good (e.g. in Example 2,  $J = 2$ , and  $\dim(S) = 4$ ). This suggests that (i) using only a part of the demand system is sufficient for obtaining an inversion, which can be used to identify  $f_\theta$  and (ii) using additional subcomponents of  $S$ , one may potentially overidentify the parameter of interest. We therefore consider an inversion that exploits a monotonicity property of the demand system that follows from this structure.<sup>5</sup> For this, we assume that the following condition is met.

**Condition 4.1.** *The random coefficient density  $f_\theta$  is continuously differentiable.  $(\epsilon_{i1t}, \epsilon_{i2t})$  and  $(D_{1t}, D_{2t})$  have full supports in  $\mathbb{R}^2$  respectively.*

---

<sup>5</sup>The additional structure can potentially be tested. In Example 2, one may identify the demand for bundles (1,0) and (1,1) using the inversion described below under the hypothesis that eq. (2.5) holds. Further, treating (1,0), (0,1), and (1,1) as three separate goods (and (0,0) as an outside good) and applying the inversion of Berry, Gandhi, and Haile (2013), one may identify the demand for bundles (1,0) and (1,1) without imposing (2.5). The specification can then be tested by comparing the demand functions obtained from these distinct inversions. We are indebted to Phil Haile for this point.

Let  $\tilde{\mathbb{L}} = \{(1, 0), (1, 1)\}$ . From (2.7), it is straightforward to show that  $\varphi_{(1,0)}$  is strictly increasing in  $D_{1t}$  but is strictly decreasing in  $D_{2t}$ , while  $\varphi_{(1,1)}$  is strictly increasing both in  $D_{1t}$  and  $D_{2t}$ . Hence, the Jacobian matrix is non-degenerate. Together with a mild support condition on  $(D_{1t}, D_{2t})$ , this allows to invert the demand (sub)system and write  $\Xi_{jt} = \psi_j(X_t^{(2)}, P_t, \tilde{S}_t) - X_{jt}^{(1)}$ , where  $\tilde{S}_t = (S_{(1,0),t}, S_{(1,1),t})$ . This ensures Assumption 2.2 in this example (see Lemma B.1 given in the appendix). By Theorem 2.1, one can then nonparametrically identify subcomponents  $(\varphi_{(1,0)}, \varphi_{(1,1)})$  of the demand function  $\varphi$ .

One may alternatively choose  $\tilde{\mathbb{L}} = \{(0, 0), (0, 1)\}$ , and the argument is similar, which then identifies  $(\varphi_{(0,0)}, \varphi_{(0,1)})$ , and hence all components of the demand function  $\varphi$  are identified. This inversion is valid even if the two goods are complements. This is because the inversion uses the monotonicity property of the aggregate choice probabilities on bundles (e.g.  $\phi_{(1,0)}$  and  $\phi_{(1,1)}$ ) with respect to  $(D_{1t}, D_{2t})$ . Hence, even if the aggregate share of each good (e.g. aggregate share on good 1:  $\sigma_1 = \phi_{(1,0)} + \phi_{(1,1)}$ ) is not invertible in the price  $P_t$  due to the presence of complementary goods, one can still obtain a useful inversion provided that aggregate choice probabilities on bundles are observed.

Given the demand for bundles, we now analyze identification of the random coefficient density. By (2.5), the demand for bundle (0,0) is given by

$$\begin{aligned} & \phi_{(0,0)}(x^{(2)}, p, \delta) \\ &= \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 < -\delta_1\}1\{x_2^{(2)'}b^{(2)} + ap_2 + e_2 < -\delta_2\} \\ & \quad \times 1\{(x_1^{(2)} + x_2^{(2)})'b^{(2)} + a(p_1 + p_2) + (e_1 + e_2) + \Delta < -\delta_1 - \delta_2\}f_\theta(b^{(2)}, a, e, \Delta)d\theta. \end{aligned} \quad (4.1)$$

Given product  $j \in \{1, 2\}$ , let  $-j$  denote the other product. We then define  $\tilde{\Phi}_l$  with  $l = (0, 0)$  as in the BLP example by letting  $D_{-jt}$  take a large negative value. For each  $(x^{(2)}, p, \delta)$ , let

$$\tilde{\Phi}_{(0,0)}(x_j^{(2)}, p_j, \delta_j) \equiv - \lim_{\delta_{-j} \rightarrow -\infty} \phi_{(0,0)}(x^{(2)}, p, \delta), \quad j = 1, 2. \quad (4.2)$$

We then define  $\Phi_{(0,0)}$  as in (3.4).<sup>6</sup> Consider for the moment  $j = 1$  in (4.2). Then,  $\Phi_{(0,0)}$  is related to the joint density  $f_{\vartheta_1}$  of  $\vartheta_{i1t} \equiv (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t})$  through a Radon transform.<sup>7</sup> Arguing as in (3.5), it is straightforward to show that  $\partial\Phi_{(0,0)}(w, u)/\partial u = \mathcal{R}[f_{\vartheta_1}](w, u)$  with  $w \equiv (x_1^{(2)}, p_1, 1)/\|(x_1^{(2)}, p_1, 1)\|$  and  $u \equiv \delta_1/\|(x_1^{(2)}, p_1, 1)\|$ . Hence, one may identify  $f_{\vartheta_1}$  by inverting

<sup>6</sup>In the BLP example, we invert a Radon transform only once. Hence  $\Phi$  in (3.4) does not have any subscript. In Examples 2 and 3, we invert Radon transform multiple times, and to make this point clear we add subscripts to  $\Phi$  (e.g.  $\Phi_{(0,0)}$  and  $\Phi_{(1,1)}$ ).

<sup>7</sup>Since the bundle effect  $\Delta_{it}$  does not appear in (4.1), one may only identify the joint density of the subvector  $(\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t})$  from the demand for bundle (0,0).

the Radon transform under Assumptions 3.1 and 3.2 with  $J = 2$ .

If the researcher is only interested in the distribution of  $(\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{ijt})$  but not in the bundle effect, the demand for (0,0) is enough for recovering their density. However,  $\Delta_{it}$  is often of primary interest. The demand on (1,1) can be used to recover its distribution by the following argument.

The demand for bundle (1,1) is given by

$$\begin{aligned} \phi_{(1,1)}(x^{(2)}, p, \delta) &= \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta > -\delta_1\}1\{x_2^{(2)'}b^{(2)} + ap_2 + e_2 + \Delta > -\delta_2\} \\ &\times 1\{(x_1^{(2)} + x_2^{(2)})'b^{(2)} + a(p_1 + p_2) + (e_1 + e_2) + \Delta > -\delta_1 - \delta_2\}f_\theta(b^{(2)}, a, e, \Delta)d\theta. \end{aligned} \quad (4.3)$$

Note that  $\Delta_{it}$  can be viewed as an additional random coefficient on the constant whose sign is fixed. Hence, the set of covariates includes a constant. Again, conditioning on an event where  $D_{-jt}$  takes a large negative value and normalizing the arguments by the norm of  $(x_j^{(2)}, p_j, 1)$  yield a function  $\Phi_{(1,1)}$  that is related to the density of  $\eta_{ijt} \equiv (\beta_{it}^{(2)}, \alpha_{it}, \Delta_{it} + \epsilon_{ijt})$  through the Radon transform in (3.2). Note that the last component of  $\eta_j$  and  $\vartheta_j$  differ only in the bundle effect  $\Delta_{it}$ . Hence, if  $\epsilon_{ijt}$  is independent of  $\Delta_{it}$  conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$ , the distribution of  $\Delta_{it}$  can be identified via deconvolution. For this, let  $\Psi_{\epsilon_j | (\beta^{(2)}, \alpha)}$  denote the characteristic function of  $\epsilon_{ijt}$  conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$ . We summarize these results below.

**Theorem 4.2.** *Suppose Assumptions 2.1-3.2 and Condition 4.1 hold with  $J = 2$  and  $\theta_{it} = (\beta_{it}^{(2)}, \alpha_{it}, \Delta_{it}, \epsilon_{i1t}, \epsilon_{i2t})$ . Suppose the conditional distribution of  $\epsilon_{ijt}$  given  $(\beta_{it}^{(2)}, \alpha_{it})$  is identical for  $j = 1, 2$ .*

*Then, (a)  $f_{\vartheta_j}, f_{\eta_j}$  are nonparametrically identified in Example 2; (b) If, in addition,  $\Delta_{it} \perp \epsilon_{ijt} | (\beta_{it}^{(2)}, \alpha_{it})$  and  $\Psi_{\epsilon_j | (\beta^{(2)}, \alpha)}(t) \neq 0$  for almost all  $t \in \mathbb{R}$  and for some  $j$ , and  $\epsilon_{ijt}, j = 1, 2$  are independently distributed (across  $j$ ) conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$ , then  $f_\theta$  is nonparametrically identified in Example 2.*

The identification of the distribution of the bundle effect requires the characteristic function of  $\epsilon_{ijt}$  to have isolated zeros (see e.g. Devroye, 1989, Carrasco and Florens, 2010). This condition can be satisfied by various distributions including the Type-I extreme value distribution and normal distribution.

**Remark 4.1.** *Note that the conditions of Theorem 4.2 do not impose any sign restriction on  $\Delta_{it}$ . Hence, the two goods can be substitutes ( $\Delta_{it} < 0$ ) for some individuals and complements ( $\Delta_{it} > 0$ ) for others. This feature, therefore, can be useful for analyzing bundles of goods whose substitution pattern can significantly differ across individuals (e.g. E-books and print books).*

## 4.2 Multiple units of consumption (Example 3)

One may also consider settings where multiple units of consumption are allowed. For simplicity, we consider the simplest setup where  $J = 2$  and  $Y_1 \in \{0, 1, 2\}$  and  $Y_2 \in \{0, 1\}$ . The utility from consuming  $y_1$  units of product 1 and  $y_2$  units of product 2 is specified as follows:

$$U_{i,(y_1,y_2),t}^* = y_1 U_{i1t}^* + y_2 U_{i2t}^* + \Delta_{i,(y_1,y_2),t} , \quad (4.4)$$

where  $\Delta_{i,(y_1,y_2),t}$  is the additional utility (or disutility) from consuming the particular bundle  $(y_1, y_2)$ . This specification allows, e.g., for decreasing marginal utility (with the number of units), as well as interaction effects. We assume that  $\Delta_{(1,0)} = \Delta_{(0,1)} = 0$  as  $U_{i1t}^*$  and  $U_{i2t}^*$  give the utility from consuming a single unit of each of the two goods. Throughout this example, we assume that  $U_{i,(y_1,y_2),t}^*$  is concave in  $(y_1, y_2)$ . Then, a bundle is chosen if its utility exceeds those of the neighboring alternatives. For example, bundle  $(2, 0)$  is chosen if it is preferred to bundles  $(1,0)$ ,  $(1,1)$  and  $(2,1)$ . That is,

$$\begin{aligned} & 2(X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t}) + \Delta_{i,(2,0),t} > X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t} , \\ & 2(X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t}) + \Delta_{i,(2,0),t} \\ & \quad > X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t} + X'_{2t}\beta_{it} + \alpha_{it}P_{2t} + \Xi_{2t} + \epsilon_{i2t} + \Delta_{i,(1,1),t} \\ & 2(X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t}) + \Delta_{i,(2,0),t} , \\ & \quad > 2(X'_{1t}\beta_{it} + \alpha_{it}P_{1t} + \Xi_{1t} + \epsilon_{i1t}) + X'_{2t}\beta_{it} + \alpha_{it}P_{2t} + \Xi_{2t} + \epsilon_{i2t} + \Delta_{i,(2,1),t}. \end{aligned} \quad (4.5)$$

The aggregate structural demand can be obtained as

$$\begin{aligned} \varphi_{(2,0)}(X_t, P_t, \Xi_t) &= \int 1\{X'_{1t}b + aP_{1t} + e_1 + \Delta_{(2,0)} > -\Xi_{1t}\} \\ & \quad \times 1\{(X_{1t} - X_{2t})'b + a(P_{1t} - P_{2t}) + (e_1 - e_2) + \Delta_{(2,0)} - \Delta_{(1,1)} > -\Xi_{1t} + \Xi_{2t}\} \\ & \quad \times 1\{X'_{2t}b + aP_{2t} + e_2 + \Delta_{(2,1)} - \Delta_{(2,0)} < -\Xi_{2t}\} f_{\theta}(b, a, e, \Delta) d\theta . \end{aligned} \quad (4.6)$$

The observed aggregate demand on the bundles are similarly defined for  $S_{l,t} = \varphi_l(X_t, P_t, \Xi_t)$ ,  $l \in \mathbb{L}$  where  $\mathbb{L} \equiv \{(0, 0), (1, 0), (0, 1), (1, 1), (2, 0), (2, 1)\}$ .

Let  $\tilde{\mathbb{L}} = \{(2, 0), (2, 1)\}$ . From (4.5),  $\varphi_{(2,0)}$  is increasing in  $D_1$  but is decreasing in  $D_2$ . Similarly,  $\varphi_{(2,1)}$  is increasing in both  $D_1$  and  $D_2$ . The rest of the argument is similar to Example 2. This ensures Assumption 2.2 in this example, and by Theorem 2.1, one can then nonparametrically identify subcomponents  $\{\varphi_l, l \in \tilde{\mathbb{L}}\}$  of the demand function  $\varphi$ . One may alternatively take  $\tilde{\mathbb{L}} = \{(0, 0), (0, 1)\}$  and use the same line of argument. Note, however, that  $(1,0)$  or  $(1,1)$  cannot be included in  $\tilde{\mathbb{L}}$  as  $\phi_{(1,0)}$  and  $\phi_{(1,1)}$  are not monotonic in one of  $(D_1, D_2)$ .

This is because increasing  $D_1$  while fixing  $D_2$ , for example, makes good 1 more attractive and creates both an inflow of individuals who move from (0,0) to (1,0) and an outflow of individuals who move from (1,0) to (2,0). Hence, the demand for (1,0) does not necessarily change monotonically.

The nonparametric IV step identifies  $\phi_l$  for  $l \in \{(0,0), (0,1), (2,0), (2,1)\}$ . Using them, we may first recover the joint density of some of the random coefficients:  $\theta_{it} = (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t}, \epsilon_{i2t}, \Delta_{i,(1,1),t}, \Delta_{i,(2,0),t}, \Delta_{i,(2,1),t})'$ . We begin with the demand for (0,0), (0,1), (2,0), and (2,1) given by

$$\begin{aligned}
\phi_{(0,0)}(x^{(2)}, p, \delta) &= \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 < -\delta_1\} \\
&\quad \times 1\{x_2^{(2)'}b^{(2)} + ap_2 + e_2 < -\delta_2\} \\
&\quad \times 1\{(x_1^{(2)} + x_2^{(2)})'b^{(2)} + a(p_1 + p_2) + (e_1 + e_2) < -\delta_1 - \delta_2\} f_\theta(b^{(2)}, a, e, \Delta) d\theta, \\
\phi_{(0,1)}(x^{(2)}, p, \delta) &= \int 1\{x_2^{(2)'}b^{(2)} + ap_2 + e_2 > -\delta_2\} \\
&\quad \times 1\{(x_1^{(2)} - x_2^{(2)})'b^{(2)} + a(p_1 - p_2) + (e_1 - e_2) < -\delta_1 + \delta_2\} \\
&\quad \times 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(1,1)} > -\delta_1\} f_\theta(b^{(2)}, a, e, \Delta) d\theta, \\
\phi_{(2,0)}(x^{(2)}, p, \delta) &= \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(2,0)} > -\delta_1\} \\
&\quad \times 1\{(x_1^{(2)} - x_2^{(2)})'b^{(2)} + a(p_1 - p_2) + (e_1 - e_2) + \Delta_{(2,0)} - \Delta_{(1,1)} > -\delta_1 + \delta_2\} \\
&\quad \times 1\{x_2^{(2)'}b^{(2)} + ap_2 + e_2 + \Delta_{(2,1)} - \Delta_{(2,0)} < -\delta_2\} f_\theta(b^{(2)}, a, e, \Delta) d\theta, \\
\phi_{(2,1)}(x^{(2)}, p, \delta) &= \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(2,1)} - \Delta_{(1,1)} > -\delta_1\} \\
&\quad \times 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(2,1)} - \Delta_{(2,0)} > -\delta_2\} \\
&\quad \times 1\{(x_1^{(2)} + x_2^{(2)})'b^{(2)} + a(p_1 + p_2) + (e_1 + e_2) + \Delta_{(2,1)} > -\delta_1 - \delta_2\} f_\theta(b^{(2)}, a, e, \Delta) d\theta.
\end{aligned}$$

Hence, if  $D_{2t}$  has a large support, by taking  $\delta_2$  sufficiently small or sufficiently large, we may

define

$$\begin{aligned}\tilde{\Phi}_{(0,0)}(x_1^{(2)}, p_1, \delta_1) &\equiv - \lim_{\delta_2 \rightarrow -\infty} \phi_{(0,0)}(x^{(2)}, p, \delta) \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 < -\delta_1\} f_\theta(b^{(2)}, a, e, \Delta) d\theta ,\end{aligned}\quad (4.7)$$

$$\begin{aligned}\tilde{\Phi}_{(0,1)}(x_1^{(2)}, p_1, \delta_1) &\equiv - \lim_{\delta_2 \rightarrow \infty} \phi_{(0,1)}(x^{(2)}, p, \delta) \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(1,1)} > -\delta_1\} f_\theta(b^{(2)}, a, e, \Delta) d\theta ,\end{aligned}\quad (4.8)$$

$$\begin{aligned}\tilde{\Phi}_{(2,0)}(x_1^{(2)}, p_1, \delta_1) &\equiv - \lim_{\delta_2 \rightarrow -\infty} \phi_{(2,0)}(x^{(2)}, p, \delta) \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(2,0)} > -\delta_1\} f_\theta(b^{(2)}, a, e, \Delta) d\theta ,\end{aligned}\quad (4.9)$$

$$\begin{aligned}\tilde{\Phi}_{(2,1)}(x_1^{(2)}, p_1, \delta_1) &\equiv - \lim_{\delta_2 \rightarrow \infty} \phi_{(2,1)}(x^{(2)}, p, \delta) \\ &= \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(2,1)} - \Delta_{(1,1)} > -\delta_1\} f_\theta(b^{(2)}, a, e, \Delta) d\theta .\end{aligned}\quad (4.10)$$

For each  $l \in \{(0, 0), (0, 1), (2, 0), (2, 1)\}$ , define  $\Phi_l$  as in (3.4). Arguing as in Example 2,  $\Phi_l$  is then related to the random coefficient densities by

$$\frac{\partial \Phi_l(w, u)}{\partial u} = \mathcal{R}[f_{\vartheta_l}](w, u), \quad l \in \{(0, 0), (0, 1), (2, 0), (2, 1)\},$$

where  $w \equiv -(x_1^{(2)}, p_1, 1) / \|(x_1^{(2)}, p_1, 1)\|$  and  $u \equiv \delta_1 / \|(x_1^{(2)}, p_1, 1)\|$ . Here, for each  $l$ ,  $f_{\vartheta_l}$  is the joint density of a subvector  $\vartheta_{i,l,t}$  of  $\theta_{it}$ , which is given by<sup>8</sup>

$$\begin{aligned}\vartheta_{i,(0,0),t} &= (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t}), \quad \vartheta_{i,(0,1),t} = (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t} + \Delta_{i,(1,1),t}), \\ \vartheta_{i,(2,0),t} &= (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t} + \Delta_{i,(2,0),t}), \quad \vartheta_{i,(2,1),t} = (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t} + \Delta_{i,(2,1),t} - \Delta_{i,(1,1),t}).\end{aligned}\quad (4.11)$$

The joint density of  $\theta_{it}$  is identified by making the following assumption.

**Assumption 4.1.** (i)  $(\Delta_{i,(1,1),t}, \Delta_{i,(2,0),t}, \Delta_{i,(2,1),t}) \perp \epsilon_{ijt} | (\beta_{it}^{(2)}, \alpha_{it})$  and  $\Psi_{\epsilon_j | (\beta^{(2)}, \alpha)}(t) \neq 0$  for almost all  $t \in \mathbb{R}$  and for some  $j \in \{1, 2\}$ ; (ii)  $\epsilon_{ijt}, j = 1, 2$  are independently and identically distributed (across  $j$ ) conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$ ; (iii)  $(\Delta_{i,(1,1),t}, \Delta_{i,(2,0),t}, \Delta_{i,(2,1),t})$  are independent of each other conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$  and  $\Psi_{\Delta_{(1,1)} | (\beta^{(2)}, \alpha)}(t) \neq 0$  for almost all  $t \in \mathbb{R}$ .

Assumption 4.1 (iii) means that, relative to the benchmark utility given as an index function of  $(X_t^{(2)}, P_t, D_t)$ , the additional utilities from the bundles are independent of each other.

---

<sup>8</sup>Alternative assumptions can be made to identify the joint density of different components of the random coefficient vector. For example, a large support assumption on  $D_{1t}$  would allow one to recover the joint density of  $(\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i2t} + \Delta_{i,(2,1),t} - \Delta_{i,(2,0),t})$  from the demand for bundle (2,0).

Assumption 4.1 (iii) also adds a regularity condition for recovering the distribution of  $\Delta_{i,(2,1),t}$  from those of  $\Delta_{i,(2,1),t} - \Delta_{i,(1,1),t}$  and  $\Delta_{i,(1,1),t}$  through deconvolution.

Identification of the joint density  $f_\theta$  allows one to recover the demand for the middle alternative: (1,0), which remained unidentified in our analysis in the nonparametric IV step. To see this, we note that the demand for this bundle is given by

$$\begin{aligned} \phi_{(1,0)}(x^{(2)}, p, \delta) &= \int \mathbf{1}\{0 < x_1^{(2)'} + ap_1 + e_1 + \delta_1 < -\Delta_{(2,0)}\} \\ &\times \mathbf{1}\{x_2^{(2)'} + ap_2 + e_2 + \delta_2 < -\Delta_{(1,1)}\} \mathbf{1}\{(x_1^{(2)} - x_2^{(2)})' + a(p_1 - p_2) + (e_1 - e_2) < -(\delta_1 - \delta_2)\} \\ &\times \mathbf{1}\{(x_1^{(2)} + x_2^{(2)})'b^{(2)} + a(p_1 + p_2) + (e_1 + e_2) + \Delta_{(2,1)} < -(\delta_1 + \delta_2)\} f_\theta(b^{(2)}, a, e, \Delta) d\theta. \end{aligned} \quad (4.12)$$

Since the previously unknown density  $f_\theta$  is identified, this demand function is identified. This and  $\phi_{(1,1)} = 1 - \sum_{l \in \mathbb{L} \setminus \{(1,1)\}} \phi_l$  further imply that all components of  $\phi$  are now identified. We summarize these results below as a theorem.<sup>9</sup>

**Theorem 4.3.** *Suppose  $U_{(y_1, y_2), t}$  is concave in  $(y_1, y_2)$ . Suppose Condition 4.1 and Assumptions 2.1, 2.3-3.1 hold with  $J = 2$  and  $\theta_{it} = (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t}, \epsilon_{i2t}, \Delta_{i,(1,1),t}, \Delta_{i,(2,0),t}, \Delta_{i,(2,1),t})$ . Suppose that  $(X_{1t}, P_{1t}, D_{1t})$  has a full support. Then, (a)  $f_{\vartheta_l}, l \in \{(0, 0), (0, 1), (2, 0), (2, 1)\}$  are nonparametrically identified in Example 3; (b) Suppose further that Assumption 4.1 holds. Then,  $f_\theta$  is identified in Example 3. Further, all components of the structural demand  $\phi$  are identified.*

### 4.3 Pure Characteristics Demand Models

Berry and Pakes (2007) study a model called the *pure characteristics demand model*, in which the utility for individual  $i$  in market  $t$  is given by

$$U_{ijt}^* \equiv X'_{jt}\beta_{it} + \alpha_{it}P_{jt} + \Xi_{jt}, \quad j = 1, \dots, J. \quad (4.13)$$

In other words, the model does not contain the unobserved tastes for products  $\{\epsilon_{ijt}, j = 1, \dots, J\}$ .<sup>10</sup> For this model, one may employ an alternative strategy to construct a function  $\Phi$  in (3.1), which does not rely on the identification at infinity argument but still achieves identification of the density of the random coefficients  $\theta_{it} = (\beta_{it}, \alpha_{it})$ . We present this alternative identification strategy by taking the multinomial choice setting as an example. Below, we maintain Assumptions 2.1-2.3, which ensure the identification of demand by Theorem 2.1.

<sup>9</sup>For simplicity, we only consider the case where  $\delta_2 \rightarrow -\infty$  or  $\infty$  in (4.7)-(4.8). This requires a full support condition on  $(X_{1t}, P_{1t}, D_{1t})$ . It is possible to replace this assumption with an analog of Assumption 3.2 (ii) by also considering the case where  $\delta_1 \rightarrow -\infty$  or  $\infty$  and imposing an additional restriction on the distribution of  $(\epsilon_{i1t}, \epsilon_{i2t}, \Delta_{i,(1,1),t}, \Delta_{i,(2,0),t}, \Delta_{i,(2,1),t})$ .

<sup>10</sup>Berry and Pakes (2007) provide detailed discussions on how a model with unobserved tastes for products differs from the pure characteristics model in terms of allowed substitution patterns and welfare implications.

The demand for good  $j$  with the product characteristics  $(X_t, P_t, \Xi_t)$  is as given in (2.3) but without the tastes for products. Since  $D_t = X_t^{(1)} + \Xi_t$ , the demand in market  $t$  with  $(X_t^{(2)}, P_t, D_t) = (x^{(2)}, p, \delta)$  is given by:

$$\begin{aligned} \phi_j(x^{(2)}, p, \delta) &= \int 1\{x_j^{(2)'}b^{(2)} + ap_j > -\delta_j\}1\{(x_j^{(2)} - x_1^{(2)})'b^{(2)} + a(p_j - p_1) > -(\delta_j - \delta_1)\} \\ &\quad \dots 1\{(x_j^{(2)} - x_J^{(2)})'b^{(2)} + a(p_j - p_J) > -(\delta_j - \delta_J)\}f_\theta(b^{(2)}, a)d\theta. \end{aligned} \quad (4.14)$$

Recall that, in the BLP model, we used Assumption 3.2 (i) (the large support condition on  $D_{kt}, k \neq j$ ) to obtain a function related to the random coefficient density through a Radon transform. In the current setting, one can take the following alternative approach.

For any subset  $\mathcal{J}$  of  $\{1, \dots, J\} \setminus \{j\}$ , let  $\mathcal{M}_{\mathcal{J}}$  denote the map  $(x^{(2)}, p, \delta) \mapsto (\acute{x}^{(2)}, \acute{p}, \acute{\delta})$  that is uniquely defined by the following properties:

$$(\acute{x}_j^{(2)} - \acute{x}_i^{(2)}, \acute{p}_j - \acute{p}_i, \acute{\delta}_j - \acute{\delta}_i) = -(x_j^{(2)} - x_i^{(2)}, p_j - p_i, \delta_j - \delta_i), \quad \forall i \in \mathcal{J}, \quad (4.15)$$

$$(\acute{x}_i^{(2)}, \acute{p}_i, \acute{\delta}_i) = (x_i^{(2)}, p_i, \delta_i), \quad \forall i \notin \mathcal{J}. \quad (4.16)$$

We then define

$$\tilde{\Phi}_j(x_j^{(2)}, p_j, \delta_j) \equiv - \sum_{\mathcal{J} \subseteq \{1, \dots, J\} \setminus \{j\}} \phi_j \circ \mathcal{M}_{\mathcal{J}}(x^{(2)}, p, \delta). \quad (4.17)$$

Eq (4.17) combines the structural demand function for good  $j$  in different markets to define a function which can be related to the random coefficient density in a simple way. This operation can be easily understood when  $J = 2$ , where for example  $\phi_1$  is given by

$$\begin{aligned} \phi_1(x^{(2)}, p, \delta) &= \int 1\{x_1^{(2)'}b^{(2)} + ap_1 < -\delta_1\} \\ &\quad \times 1\{(x_1^{(2)} - x_2^{(2)})'b^{(2)} + a(p_1 - p_2) < -(\delta_1 - \delta_2)\}f_\theta(b^{(2)}, a)d\theta. \end{aligned} \quad (4.18)$$

Then,  $\tilde{\Phi}_1$  is given by

$$\begin{aligned} \tilde{\Phi}_1(x_1^{(2)}, p_1, \delta_1) &= -\phi_1 \circ \mathcal{M}_\emptyset(x_1^{(2)} - x_2^{(2)}, p_1 - p_2, \delta_1 - \delta_2) - \phi_1 \circ \mathcal{M}_{\{2\}}(x_1^{(2)} - x_2^{(2)}, p_1 - p_2, \delta_1 - \delta_2) \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 < -\delta_1\} \left( 1\{(x_1^{(2)} - x_2^{(2)})'b^{(2)} + a(p_1 - p_2) < -(\delta_1 - \delta_2)\} \right. \\ &\quad \left. + 1\{(x_1^{(2)} - x_2^{(2)})'b^{(2)} + a(p_1 - p_2) > -(\delta_1 - \delta_2)\} \right) f_\theta(b^{(2)}, a)d\theta \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 < -\delta_1\} f_\theta(b^{(2)}, a)d\theta \end{aligned} \quad (4.19)$$

This shows that aggregating the demand in the two markets with  $(X_t^{(2)}, P_t, D_t) = (x^{(2)}, p, \delta)$



and  $(X_{t'}^{(2)}, P_{t'}, D_{t'}) = (\hat{x}^{(2)}, \hat{p}, \hat{\delta})$  yields  $\tilde{\Phi}_1$  which depends only on the utility from products 1 and 0, where the latter is normalized to 0. Eq. (4.17) generalizes this argument to settings with  $J \geq 2$ . For the general setting, we make the following assumption.

**Assumption 4.2.** (i) For any  $\mathcal{J} \subseteq \{1, \dots, J\} \setminus \{j\}$  and any  $(x^{(2)}, p, \delta) \in \text{supp}(X_t^{(2)}, P_t, D_t)$ , we have  $\mathcal{M}_{\mathcal{J}}(x^{(2)}, p, \delta) \in \text{supp}(X_t^{(2)}, P_t, D_t)$ .

Assumption 4.2 ensures that  $\Phi$  is well-defined. It requires that for any  $(x^{(2)}, p, \delta) \in \text{supp}(X_t^{(2)}, P_t, D_t)$  and  $\mathcal{J} \subseteq \{1, \dots, J\} \setminus \{j\}$ , the operation  $\mathcal{M}_{\mathcal{J}}$  gives another point in the support. A full support assumption on  $(X_t^{(2)}, P_t, D_t)$  is sufficient for this condition. One thing to note is that our identification argument based on Assumption 4.2 constructs  $\tilde{\Phi}_j$  without relying on a “thin” (lower-dimensional) subset of the support of  $(X_{jt}^{(2)}, P_{jt}, D_{jt})$  as done in the BLP model. This comes from the fact that we no longer need to let  $D_{kt}, k \neq j$  tend to infinity. Instead, one can construct  $\tilde{\Phi}_j$  in (4.17) by combining the demand in different markets.<sup>11</sup> Under Assumption 4.2, one can construct a function  $\tilde{\Phi}_l$ , which can be written in general as  $\tilde{\Phi}_l(X_j^{(2)}, P_j, D_j) = \int 1\{X_j^{(2)'}\beta^{(2)} + aP_j < -D_j\}f_{\theta}(b^{(2)}, a)d\theta$ . Let  $w \equiv (x_j^{(2)}, p_j)/\|(x_j^{(2)}, p_j)\|$  and  $u \equiv \delta_j/\|(x_j^{(2)}, p_j)\|$ . Define

$$\Phi(w, u) \equiv \tilde{\Phi}_l\left(\frac{x_j^{(2)}}{\|(x_j^{(2)}, p_j)\|}, \frac{p_j}{\|(x_j^{(2)}, p_j)\|}, \frac{\delta_j}{\|(x_j^{(2)}, p_j)\|}\right) = \tilde{\Phi}_j(x_j^{(2)}, p_j, \delta_j). \quad (4.20)$$

The rest of the analysis is then similar to the BLP model. One should note, however, that one does not need to invoke additional independence (or i.i.d.) assumptions on  $(\epsilon_{i1t}, \dots, \epsilon_{iJt})$  to identify the joint distribution of  $\theta$ .

**Theorem 4.4.** Suppose Assumptions 2.1-3.1 and 4.2 hold. Then,  $f_{\theta}$  is identified in the pure characteristics demand model, where  $\theta_{it} = (\beta_{it}^{(2)}, \alpha_{it})$ .

One thing to note is that the utility specification adopted in the pure characteristics model can also be combined with the bundle choice and multiple units of consumption. The identification of the random coefficients can be achieved using arguments similar to the ones in this section<sup>12</sup>.

## 4.4 Alternative specific coefficients

So far, we have maintained the assumption that  $(\beta_{ijt}, \alpha_{ijt}) = (\beta_{it}, \alpha_{it}), \forall j$  almost surely. This excludes alternative specific random coefficients. However, this is not essential in our analysis.

<sup>11</sup>If Assumption 4.2 does not hold, however, one may alternatively rely on an identification argument that uses a lower dimensional subset and retain the same identification result as done in the BLP model.

<sup>12</sup>The analysis of these settings are contained in an earlier version of this paper, which is available from the authors upon request.

One may allow some or all components of  $(\beta_{ijt}, \alpha_{ijt})$  to be different random variables across  $j$  and identify their joint distribution under an extended support condition on the product characteristics.

We first note that the aggregate demand is identified as long as Assumptions 2.1-2.3 hold. In the BLP model, the marginal density  $f_{\vartheta_j}$  of  $\vartheta_{ijt} = (\beta_{ijt}^{(2)}, \alpha_{ijt}, \epsilon_{ijt})$  can be identified for any  $j$  as long as the corresponding product characteristics  $(X_{jt}^{(2)}, P_{jt}, D_{jt})$  has a full support using the same identification strategy in Section 3 (see Remark 3.1). For the pure characteristics demand model, we note that the maps  $\mathcal{M}_{\mathcal{J}}$  cannot be used as the use of this map is justified when  $(\beta_{ijt}, \alpha_{ijt}) = (\beta_{it}, \alpha_{it}), \forall j$ . However, the large support assumption on  $D_{kt}, k \neq j$  (Assumption 3.2 (i)) can still be used to construct  $\Phi$ . Hence, the analysis of this case becomes similar to the BLP model. In both models, the joint density  $f_{\theta}$  of  $\theta_{it} = (\vartheta_{i1t}, \dots, \vartheta_{iJt})$  can be recovered under the assumption that  $\vartheta_{ijt}$  are independent across  $j$ .

When the covariates  $(X_t^{(2)}, P_t, D_t)$  have rich variations jointly, it is also possible to identify the joint density  $f_{\theta}$  without the independence assumption invoked above. This requires us to extend our identification strategy. To see this, we take Example 2 as an illustration below. Consider identifying the joint density of  $\theta_{it} = (\beta_{i1t}^{(2)}, \beta_{i2t}^{(2)}, \alpha_{i1t}, \alpha_{i2t}, \epsilon_{i1t}, \epsilon_{i2t} + \Delta_{it})$  under the assumption that the two goods are complements, i.e.  $\Delta_{it} > 0, a.s.$  In this setting, we may use the demand for bundle  $(1, 0)$ , which can be written as

$$\begin{aligned} \phi_{(1,0)}(x^{(2)}, p, \delta) &= \int 1\{x_1^{(2)'}b_1^{(2)} + a_1p_1 + e_1 > -\delta_1\} \\ &\quad \times 1\{x_2^{(2)'}b_2^{(2)} + a_2p_2 + e_2 + \Delta < -\delta_2\} f_{\theta}(b_1^{(2)}, b_2^{(2)}, a_1, a_2, \Delta) d\theta. \end{aligned} \quad (4.21)$$

To recover the joint density, one has to directly work with this demand function without simplifying it further. A key feature of (4.21) is that it involves multiple indicator functions and that distinct subsets of  $\theta$  show up in each of these indicator functions. For example, the first indicator function in (4.21) involves  $(\beta_{i1t}^{(2)}, \alpha_{i1t}, \epsilon_{i1t})$ , while the second indicator function involves  $(\beta_{i2t}^{(2)}, \alpha_{i2t}, \epsilon_{i2t} + \Delta_{it})$ . Integral transforms of this form are studied in Dunker, Hoderlein, and Kaido (2013) in their analysis of random coefficients discrete game models. They use tensor products of integral transforms to study nonparametric identification of random coefficient densities. Using their framework, one may show that

$$\frac{\partial^2 \phi_{(1,0)}(w_1, w_2, u_1, u_2)}{\partial u_1 \partial u_2} = (\mathcal{R} \otimes \mathcal{R})[f_{\theta}](w_1, w_2, u_1, -u_2), \quad (4.22)$$

where  $w_1 = -(x_1^{(2)}, p_1, 1) / \|(x_1^{(2)}, p_1, 1)\|$ ,  $w_2 = (x_2^{(2)}, p_2, 1) / \|(x_2^{(2)}, p_2, 1)\|$ ,  $u_1 = -\delta_1 / \|(x_1^{(2)}, p_1, 1)\|$ ,  $u_2 = \delta_2 / \|(x_2^{(2)}, p_2, 1)\|$ , and  $\mathcal{R} \otimes \mathcal{R}$  is the tensor product of Radon transforms, which can be inverted to identify  $f_{\theta}$ . The main principle of our identification strategy is therefore the same

as before. Inverting the transform in (4.22) to identify  $f_\theta$  requires Assumption 3.1 (ii) to be strengthened as follows.

**Assumption 4.3.**  $(X_{1t}^{(2)}, P_{1t}, D_{1t}, X_{2t}^{(2)}, P_{2t}, D_{2t})$  has a full support.

This is a stronger support condition than Assumption 3.2 (ii) as it requires a joint full support condition for the characteristics of both goods. This condition is violated, for example, when there is a common covariate that enters the characteristics of both goods. This is in line with the previous findings in the literature that identifying the joint distribution of potentially correlated unobservable tastes for products (e.g.  $\epsilon_1$  and  $\epsilon_2$ ) requires variables that are excluded from one or more goods (see e.g. Keane, 1992 and Gentzkow, 2007). Identification of  $f_\theta$  is then established by the following theorem.<sup>13</sup>

**Theorem 4.5.** *In Example 2, let  $\theta_{it} = (\beta_{i1t}^{(2)}, \beta_{i2t}^{(2)}, \alpha_{i1t}, \alpha_{i2t}, \epsilon_{i1t}, \epsilon_{i2t} + \Delta_{it})$ . Suppose that Assumptions 2.1-2.3, 3.1 (i), (iii), and 4.3 hold. Suppose further that  $\Delta_{it} > 0$ , a.s. Then,  $f_\theta$  is identified.*

## 4.5 Nonparametric identification of $\psi$ with full independence

In Section 2.2, we discussed the the nonparametric identification of the functions  $\psi_j$  in the equation  $\Xi_{jt} = \psi_j(X_t^{(2)}, P_t, \tilde{S}_t) - X_{jt}^{(1)}$ . Following BH (2013), we proposed to identify the structural functions by the conditional moment equations

$$E\left[\psi_j\left(X_t^{(2)}, P_t, S_t\right) \middle| Z_t = z_t, X_t = \left(x_t^{(1)}, x_t^{(2)}\right)\right] = x_{jt}^{(1)}, \quad j = 1, \dots, J.$$

with instrumental variables  $Z_t$ . The identification relies on the assumption that the unobservable  $\Xi_{jt}$  is mean independent of the instruments. However, in many applications researchers choose instruments by arguing that they are independent of the unobservable. Using only mean independence means using only parts of the available information. Thereby, the identifying power is weakened. Adding the stronger independence assumption when it is justified will improve identification as well as estimation. Therefore, we propose an approach similar to Dunker et. al. (2014) by formally assuming

$$\Xi_{jt} \perp\!\!\!\perp (Z_t, X_t) \quad \text{and} \quad E[\Xi_{jt}] = 0 \quad \text{for all } j, t.$$

---

<sup>13</sup>We omit the proof of this result for brevity. It is also possible to disentangle the distribution of  $\Delta_{it}$  from that of  $\epsilon_{i2t} + \Delta_{it}$  using a deconvolution argument as done in Theorem 4.2.

This leads to the nonlinear equation

$$0 = \begin{pmatrix} P[\psi_j(X_t^{(2)}, S_t, P_t) - X_{jt}^{(1)} \leq \xi] - P[\psi_j(X_t^{(2)}, S_t, P_t) - X_{jt}^{(1)} \leq \xi | Z_t = z_t, X_t = x_t] \\ E[\psi_j(X_t^{(2)}, S_t, P_t) - X_{jt}^{(1)}] \end{pmatrix}$$

for all  $\xi, z_t, x_t$ . Nonparametric estimation of problems involving this type of nonlinear restrictions are studied in Chen and Pouzo (2012) and Dunker et. al. (2014). To give sufficient conditions for identification, we define the operator

$$F(\varphi)(\xi, z_t, x_t) := \begin{pmatrix} P[\varphi(X_t^{(2)}, S_t, P_t) - X_{jt}^{(1)} \leq \xi] - P[\varphi(X_t^{(2)}, S_t, P_t) - X_{jt}^{(1)} \leq \xi | Z_t = z_t, X_t = x_t] \\ E[\varphi(X_t^{(2)}, S_t, P_t) - X_{jt}^{(1)}] \end{pmatrix}.$$

The function  $\psi_j$  is a root of the operator  $F$ . It is, therefore, globally identified under the following assumption.

**Assumption 4.4.** *The operator  $F$  has a unique root.*

On first sight this may appear as a strong assumption due to the complexity of the operator. It is, however, weaker than the usual completeness assumption for the mean independence assumption. This is because, if  $\Xi_{jt} \perp (Z_t, X_t)$  and the usual completeness assumption hold, then  $F$  has only one root. On the other hand, completeness is not necessary for  $F$  to have a unique root. Hence, when  $\Xi_{jt} \perp (Z_t, X_t)$ , Assumption 4.4 is weaker than Assumption 2.3. Another important advantage of this method is that because the  $D_j$  do not vanish, we have a close analog to nonparametric IV with full independence, see, e.g., Dunker et al (2014), where  $D_j$  now plays the role of the dependent variable.

## 5 Suggested estimation methods

### 5.1 Nonparametric estimator

The structure of the nonparametric identification suggests a nonparametric estimation strategy in a natural way. It consists of three steps. The first step is the estimation of the structural function  $\psi_j$ . The second step is to derive the function  $\Phi$  from the estimated  $\hat{\psi}_j$ . This requires only straightforward algebraic manipulation which were presented in Section 3. We will not further comment on these computations. The last step of the estimation is the inversion of a Radon transform.

The mathematical structure of the first step is similar to nonparametric IV. The conditional

expectation operator on the left hand side of the equation

$$E[\psi_j(x_t^{(2)}, P_t, S_t) | Z_t = z_t, X_t = x_t] = x_{jt}^{(1)} \quad \text{for all } x_t, z_t$$

has to be inverted. Let us denote this linear operator by  $T$  and rewrite the problem as  $(T\psi_j)(z_t, x_t) = x_{jt}^{(1)}$ . Here  $x_{jt}^{(1)}$  should be interpreted as a function in  $x_t$  and  $z_t$  which is constant in  $x_t^{(2)}$ ,  $z_t$ , and  $x_{it}^{(1)}$  for  $i \neq j$ . The operator depends on the joint density of  $(X_t, P_t, S_t, Z_t)$  which has to be estimated nonparametrically, e.g. by kernel density estimation. This gives an estimator  $\widehat{T}$ . As in nonparametric IV the operator equation is usually ill-posed. Regularized inversion schemes must be applied. We propose Tikhonov regularization for this purpose:

$$\widehat{\psi}_j := \min_{\psi} \|\widehat{T}\psi - x_{jt}^{(1)}\|_{L^2(X_t, Z_t)}^2 + \alpha \mathfrak{R}(\psi). \quad (5.1)$$

As usual  $\alpha \geq 0$  is a regularization parameter and  $\mathfrak{R}$  a regularization functional. The usual choice would be  $\mathfrak{R}(\psi) = \|\psi\|_{L^2}^2$ . If more smoothness is expected, this could be a squared Sobolev norm or some other norm as well. In the case of bundles and multiple goods we know that  $\psi$  must be monotonically increasing or decreasing in  $S_t$ . One may incorporate this a priori knowledge by setting  $\mathfrak{R}(\psi) = \infty$  for all functions  $\psi$  not having this property. Monotonicity is a convex constraint. Hence, even with this choice of  $\mathfrak{R}$ , equation (5.1) is a convex minimization problem. Solving the problem is computationally feasible. We refer to Eggermont (1993), Burger and Osher (2004), and Resmerita (2005) for regularization with general convex regularization functional. Furthermore, we refer to Newey and Powell (2003) for the related nonparametric IV problem.

The third step of our nonparametric estimation strategy is the inversion of a Radon transform. A popular and efficient method for the problem is the filtered back projection

$$\widehat{f}_\theta(\vartheta) = \mathcal{R}^* \left( \Omega_r *_{\delta} \frac{\partial \Phi_j(x_j^{(2)}, p_j, \delta_j)}{\partial \delta_j} \right) (\vartheta).$$

Here  $\vartheta = (b, a)$  or  $(b, a, \Delta)$  depending on the application,  $(R^*g)(x) := \int_{\|w\|=1} g(w, w'x)dw$  is the adjoint of the Radon transform, and  $*_{\delta}$  denotes the convolution with respect to the last variable  $\delta_j$ , and  $\Omega_r$  is the function

$$\Omega_r(s) := \frac{1}{4\pi^2} \begin{cases} (\cos(rs) - 1)/s^2 & \text{for } s \neq 0, \\ r^2/2 & \text{for } s = 0. \end{cases}$$

For more details on this algorithm in a deterministic setting we refer to Natterer (2001). A similar estimator for random coefficients is proposed and analyzed in HKM.

## 5.2 Parametric estimators for bundle choice models

Our nonparametric identification analysis shows that the choice of bundles and multiple units of consumption can be studied very much in the same way as the standard BLP model (or the pure characteristic model). This suggests that one may construct parametric estimators for these models by extending standard estimation methods, given appropriate data. Below, we take Example 2 and illustrate this idea.

Let  $\theta_{it} = (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{1it}, \epsilon_{2it}, \Delta_{it})$  be random coefficients and let  $f_{\theta}(\cdot; \gamma)$  be a parametric density function, where  $\gamma$  belongs to a finite dimensional parameter space  $\Gamma \subset \mathbb{R}^{d_{\gamma}}$ . The estimation procedure consists of the following steps:

**Step 1** : Compute the aggregate share of bundles as a function of parameter  $\gamma$  conditional on the set of covariates.

**Step 2** : Use numerical methods to solve demand systems for  $(D_{1t}, D_{2t})$ , where  $D_{jt} = \Xi_{jt} + X_{jt}^{(1)}$ ,  $j = 1, 2$  and obtain the inversion in eq. (2.8).

**Step 3** : Form a GMM criterion function using instruments and minimize it with respect to  $\gamma$  over the parameter space.

The first step is to compute the aggregate share. In the pure characteristic model, one may approximate the aggregate share of each bundle such as the one in (2.7) by simulating  $\theta$  from  $f_{\theta}(\cdot; \gamma)$  for each  $\gamma$ . Specifically, if the conditional CDF of  $\epsilon_{ijt}$  given  $(\beta_{it}^{(2)}, \alpha_{it}, \Delta_{it})$  has an analytic form, the two-step method in BLP and Berry and Pakes (2007) can be employed.<sup>14</sup> We take the demand for bundle (0,0) in eq. (4.1) as an example. Conditional on the product characteristics  $y \equiv (x^{(2)}, p, \delta)$  and the rest of the random coefficients  $(\beta_{it}^{(2)}, \alpha_{it}, \Delta_{it})$ , bundle (0,0) is chosen when

$$\epsilon_{i1t} < h_1(y, b^{(2)}, a, \Delta) \quad \text{and} \quad \epsilon_{i2t} < h_2(y, b^{(2)}, a, \Delta), \quad \text{if } \Delta < 0 \quad (5.2)$$

$$\epsilon_{i1t} < h_2(y, b^{(2)}, a, \Delta) \quad \text{and} \quad \epsilon_{i1t} + \epsilon_{i2t} < h_3(y, b^{(2)}, a, \Delta), \quad \text{if } \Delta \geq 0 \quad (5.3)$$

where

$$h_1(y, \beta^{(2)}, a, \Delta) \equiv -x_1^{(2)'} b^{(2)} - ap_1 - \delta_1, \quad h_2(y, \beta^{(2)}, a, \Delta) \equiv -x_2^{(2)'} b^{(2)} - ap_2 - \delta_2, \\ h_3(y, \beta^{(2)}, a, \Delta) \equiv -(x_1^{(2)} + x_2^{(2)})' b^{(2)} - a(p_1 + p_2) - (\delta_1 - \delta_2). \quad (5.4)$$

---

<sup>14</sup>For the pure characteristics demand model, a similar strategy can be taken if the conditional CDF of  $\alpha_{it}$  given  $(\beta_{it}^{(2)}, \Delta_{it})$  has an analytic form.

Specify the conditional distribution of  $(\epsilon_{i1t}, \epsilon_{i2t})$  given  $(\beta_{it}^{(2)}, \alpha_{it}, \Delta_{it})$ . For each  $(y, b^{(2)}, a, \Delta)$ , define

$$G(y, b^{(2)}, a, \Delta) \equiv \begin{cases} Pr(\epsilon_{i1t} < h_1(y, b^{(2)}, a, \Delta), \epsilon_{i2t} < h_2(y, b^{(2)}, a, \Delta) | y, b^{(2)}, a, \Delta) & \Delta < 0 \\ Pr(\epsilon_{i1t} < h_2(y, b^{(2)}, a, \Delta), \epsilon_{i1t} + \epsilon_{i2t} < h_3(y, b^{(2)}, a, \Delta) | y, b^{(2)}, a, \Delta) & \Delta > 0. \end{cases} \quad (5.5)$$

The value of  $G(y, b^{(2)}, a, \Delta)$  can be calculated analytically, for example, if one specifies the joint distribution of  $(\epsilon_{i1t}, \epsilon_{i2t})$  as normal. Eq. (5.2)-(5.3) then imply that the aggregate share of bundle (1,0) is given by

$$\phi_{(1,0)}(x^{(2)}, p, \delta; \gamma) = \int G(y, b^{(2)}, a, \Delta) f_{\beta^{(2)}, a, \Delta}(b, a, \Delta; \gamma) d\theta. \quad (5.6)$$

This can be approximated by the simulated moment:

$$\hat{\phi}_{(1,0)}(x^{(2)}, p, \delta; \gamma) = \frac{1}{n_S} \sum_{i=1}^{n_S} G(y, b_i^{(2)}, a_i, \Delta_i), \quad (5.7)$$

where the simulated sample  $\{(b_i^{(2)}, a_i, \Delta_i), i = 1, \dots, n_S\}$  is generated from  $f_{\beta^{(2)}, a, \Delta}(\cdot; \gamma)$ .<sup>15</sup> Computation of the aggregate demand for other bundles is similar. This step therefore gives the model predicted aggregate demand  $\hat{\phi}_l$  for all bundles under a chosen parameter value  $\gamma$ .

The next step is then to invert subsystems of demand and obtain  $\psi$  numerically. Given  $\hat{\phi}_l, l \in \mathbb{L}$  from Step 1, this step can be carried out by numerically calculating inverse mappings. For example, take  $\tilde{\mathbb{L}} = \{(1, 0), (1, 1)\}$ . Then,  $(\delta_1, \delta_2) \mapsto (\hat{\phi}_{(1,0)}(x^{(2)}, p, \delta; \gamma), \hat{\phi}_{(1,1)}(x^{(2)}, p, \delta; \gamma))$  defines a mapping from  $\mathbb{R}^2$  to  $[0, 1]^2$ . Standard numerical methods such as the Newton-Raphson method or the homotopy method (see Berry and Pakes, 2007) can then be employed to calculate the inverse of this mapping<sup>16</sup>, which then yields  $\hat{\psi}(\cdot; \gamma) \equiv (\hat{\psi}_1(\cdot; \gamma), \hat{\psi}_2(\cdot; \gamma))$  such that

$$\Xi_{1,t} = \hat{\psi}_1(X_t^{(2)}, P_t, S_{(1,0),t}, S_{(1,1),t}; \gamma) - X_{1t}^{(1)}, \quad \Xi_{2,t} = \hat{\psi}_2(X_t^{(2)}, P_t, S_{(1,0),t}, S_{(1,1),t}; \gamma) - X_{2t}^{(1)} \quad (5.8)$$

where  $(S_{(1,0),t}, S_{(1,1),t})$  are observed shares of bundles. One may further repeat this step with  $\tilde{\mathbb{L}} = \{(0, 0), (0, 1)\}$ , which yields

$$\Xi_{1,t} = \hat{\psi}_3(X_t^{(2)}, P_t, S_{(0,0),t}, S_{(0,1),t}; \gamma) - X_{1t}^{(1)}, \quad \Xi_{2,t} = \hat{\psi}_4(X_t^{(2)}, P_t, S_{(0,0),t}, S_{(0,1),t}; \gamma) - X_{2t}^{(1)} \quad (5.9)$$

This helps generate additional moment restrictions in the next step.

<sup>15</sup>One may also use an importance sampling method.

<sup>16</sup>Whether the demand subsystems admit an analog of BLP's contraction mapping method is an interesting open question, which we leave for future research.

The third step is to use (5.8)-(5.9) to generate moment conditions and estimate  $\gamma$  by GMM. There are four equations in total, while because the shares sum up to 1 one equation is redundant. Hence, by multiplying instruments to the residuals from the first three equations, we define the sample moment:

$$g_n(X_t, P_t, S_t, Z_t; \gamma) \equiv \frac{1}{n} \sum_{t=1}^n \begin{pmatrix} \hat{\psi}_1(X_t^{(2)}, P_t, S_{(1,0),t}, S_{(1,1),t}; \gamma) - X_{1t}^{(1)} \\ \hat{\psi}_2(X_t^{(2)}, P_t, S_{(1,0),t}, S_{(1,1),t}; \gamma) - X_{2t}^{(1)} \\ \hat{\psi}_3(X_t^{(2)}, P_t, S_{(0,0),t}, S_{(0,1),t}; \gamma) - X_{1t}^{(1)} \end{pmatrix} \otimes \begin{pmatrix} Z_t \\ X_t \end{pmatrix}.$$

Letting  $W_n(\gamma)$  be a (possibly data dependent) positive definite matrix, define the GMM criterion function by

$$Q_n(\gamma) \equiv g_n(X_t, P_t, S_t, Z_t; \gamma)' W_n(\gamma) g_n(X_t, P_t, S_t, Z_t; \gamma).$$

The GMM estimator  $\hat{\gamma}$  of  $\gamma$  can then be computed by minimizing  $Q_n$  over the parameter space. A key feature of this method is that it uses the familiar BLP methodology (simulation, inversion & GMM) but yet allows one to estimate models that do not fall in the class of multinomial choice models. Employing our procedure may, for example, allow one to estimate bundle choices (e.g. print newspaper, online newspaper, or both) or platform choices using market level data.

## 6 Outlook

This paper is concerned with the nonparametric identification of models of market demand. It provides a general framework that nests several important models, including the workhorse BLP model, and provides conditions under which these models are point identified. Important conclusions include that the assumption necessary to recover various objects differ; in particular, it is easier to identify demand elasticities and more difficult to identify the individual specific random coefficient densities. Moreover, the data requirements are also shown to vary with the model considered. The identification analysis is constructive, extends the classical nonparametric BLP identification as analyzed in BH to other models, and opens up the way for future research on sample counterpart estimation. A particularly intriguing part hereby is the estimation of the demand elasticities, as the moment condition is different from the one used in nonparametric IV. Understanding the properties of these estimators, and evaluating their usefulness in an application, is an open research question that we hope this paper stimulates.



# References

- [1] ACKERBERG, D., C. L. BENKARD, S. BERRY, AND A. PAKES (2007): “Chapter 63 Econometric Tools for Analyzing Market Outcomes,” vol. 6, Part A of *Handbook of Econometrics*, pp. 4171 – 4276. Elsevier.
- [2] ANDREWS, D. (2011): “Examples of L2-Complete and Boundedly-complete Distributions,” Cowles Foundation Discussion Paper.
- [3] ARMSTRONG, T. (2013): “Large Market Asymptotics for Differentiated Product Demand Estimators with Economic Models of Supply,” Discussion Paper, Yale University.
- [4] BENNING, M., AND M. BURGER (2011): “Error estimates for general fidelities,” *Electron. Trans. Numer. Anal.*, 38, 44–68.
- [5] BERAN, R., AND P. HALL (1992): “Estimating coefficient distributions in random coefficient regressions,” *The annals of Statistics*, 20(4), 1970–1984.
- [6] BERRY, S. T., A. GANDHI, AND P. A. HAILE (2013): “Connected Substitutes and Invertibility of Demand,” *Econometrica*, 81(5), 2087–2111.
- [7] BERRY, S. T., AND P. A. HAILE (2011): “Nonparametric Identification of Multinomial Choice Demand Models With Heterogeneous Consumers,” Cowles Foundation Discussion Paper.
- [8] ——— (2013): “Identification in Differentiated Products Markets Using Market Level Data,” Cowles Foundation Discussion Paper.
- [9] BERRY, S. T., J. A. LEVINSOHN, AND A. PAKES (1995): “Automobile Prices in Market Equilibrium,” *Econometrica: Journal of the Econometric Society*, pp. 841–890.
- [10] ——— (2004): “Differentiated Products Demand Systems from a Combination of Micro and Macro Data: The New Car Market,” *Journal of Political Economy*, 112(1), 68–105.
- [11] BERRY, S. T., AND A. PAKES (2007): “The Pure Characteristics Demand Model,” *International Economic Review*, 48(4), 1193–1225.
- [12] BURGER, M., AND S. OSHER (2004): “Convergence rates of convex variational regularization,” *Inverse problems*, 20(5), 1411.
- [13] CARRASCO, M., AND J. FLORENS (2010): “A spectral method for deconvolving a density,” *Econometric Theory*, 27(3), 546–581.

- [14] CHEN, X., AND D. POUZO (2012): “Estimation of nonparametric conditional moment models with possibly nonsmooth generalized residuals,” *Econometrica*, 80(1), 277–321.
- [15] CRAMÉR, H., AND H. WOLD (1936): “Some Theorems on Distribution Functions” *Journal of the London Mathematical Society*, s1-11(4), 290–294.
- [16] DUBÉ, J.-P., J. T. FOX, AND C.-L. SU (2012): “Improving the Numerical Performance of Static and Dynamic Aggregate Discrete Choice Random Coefficients Demand Estimation,” *Econometrica*, 80(5), 2231–2267.
- [17] DUNKER, F., J.-P. FLORENS, T. HOHAGE, J. JOHANNES, AND E. MAMMEN (2014): “Iterative Estimation of Solutions to Noisy Nonlinear Operator Equations in Nonparametric Instrumental Regression,” *Journal of Econometrics*, 178(3), 444 – 455.
- [18] DUNKER, F., S. HODERLEIN, AND H. KAIDO (2013): “Random Coefficients in Static Games of Complete Information,” CEMMAP Working Paper.
- [19] EGGERMONT, P. P. B. (1993): “Maximum entropy regularization for Fredholm integral equations of the first kind,” *SIAM J. Math. Anal.*, 24, 1557–1576.
- [20] FOX, J., AND N. LAZZATI (2013): “Identification of Discrete Choice Models for Bundles and Binary Games,” CEMMAP Working Paper.
- [21] FOX, J. T., AND A. GANDHI (2012): “Using selection decisions to identify the joint distribution of outcomes,” .
- [22] GALE, D., AND J. NIKAIDO (1965): “The Jacobian matrix and global univalence of mappings,” *Math. Ann.*, 159(2), 81–93.
- [23] GAUTIER, E., AND S. HODERLEIN (2013): “Estimating the Distribution of Treatment Effects,” *Arxiv preprint arXiv:1109.0362*.
- [24] GAUTIER, E., AND Y. KITAMURA (2013): “Nonparametric Estimation in Random Coefficients Binary Choice Models,” *Econometrica*, 81(2), 581–607.
- [25] GENTZKOW, M. A. (2007): “Valuing New Goods in a Model with Complementarity: Online Newspapers,” *American Economic Review*, 97(3), 713–744.
- [26] HECKMAN, J. J., R. L. MATZKIN AND L. NESHEIM (2010): “Nonparametric Identification and Estimation of Nonadditive Hedonic Models,” *Econometrica*, 78(5), 1569–1591.
- [27] HELGASON, S. (1999): *The Radon Transform*. Birkhauser, Boston-Basel-Berlin, 2nd edition edn.

- [28] HODERLEIN, S., H. HOLZMANN, AND A. MEISTER (2014): “The Triangular Model with Random Coefficients,” Working Paper., Boston College.
- [29] HODERLEIN, S., J. KLEMELÄ, AND E. MAMMEN (2010): “Analyzing the Random Coefficient Model Nonparametrically,” *Econometric Theory*, 26(03), 804–837.
- [30] ICHIMURA, H., AND T. THOMPSON (1998): “Maximum Likelihood Estimation of a Binary Choice Model with Random Coefficients of Unknown Distribution,” *Journal of Econometrics*, 86(2), 269–295.
- [31] KEANE, M. P. (1992): “A note on identification in the multinomial probit model,” *Journal of Business & Economic Statistics*, 10(2), 193–200.
- [32] KRANTZ, S. G., AND H. R. PARKS (2002): *The implicit function theorem: history, theory, and applications*. Birkhäuser, Boston, USA.
- [33] MASTEN, M. A. (2014): “Random Coefficients on Endogenous Variables in Simultaneous Equations Models,” .
- [34] MATZKIN, R. L. (2012): “Identification in Nonparametric Limited Dependent Variable Models with Simultaneity and Unobserved Heterogeneity,” *Journal of Econometrics*, 166(1), 106–115.
- [35] MCFADDEN, D. L. (1974): “Conditional Logit Analysis of Qualitative Choice Behavior,” *Frontiers in Econometrics*, pp. 105–142.
- [36] MCFADDEN, D. L. (1981): “Structural Discrete Probability Models Derived from Theories of Choice,” In Charles Manski and Daniel McFadden, editors, *Structural Analysis of Discrete Data and Econometric Applications* , The MIT Press, Cambridge.
- [37] MOON, H. R., M. SHUM, AND M. WEIDNER (2014): “Estimation of Random Coefficients Logit Demand Models with Interactive Fixed Effects,” CEMMAP Working Paper.
- [38] NATTERER, F. (2001): *The mathematics of computerized tomography*, vol. 32 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, Reprint of the 1986 original.
- [39] NEVO, A. (2001): “Measuring Market Power in the Ready-to-Eat Cereal Industry,” *Econometrica*, 69, 307–342.
- [40] NEWEY, W. K., AND J. L. POWELL (2003): “Instrumental Variable Estimation of Nonparametric Models,” *Econometrica*, 71(5), 1565–1578.

- [41] PETRIN, A. (2002): “Quantifying the Benefits of New Products: The Case of the Minivan,” *Journal of Political Economy*, 110(4), 705–729.
- [42] RESMERITA, E. (2005): “Regularization of ill-posed problems in Banach spaces: convergence rates,” *Inverse Problems*, 21(4), 1303–1314.

## A Notation and Definitions

The following is a list of notations and definitions used throughout the appendix.

- $\mathbb{S}^{q-1}$ : The unit sphere  $\mathbb{S}^{q-1} \equiv \{v \in \mathbb{R}^q : \|v\| = 1\}$ .
- $\mathbb{H}_+$ : The hemisphere  $\mathbb{H}_+ \equiv \{v = (v_1, v_2, \dots, v_q) \in \mathbb{S}^{q-1} : v_q \geq 0\}$ .
- $P_{w,r}$ : The hyperplane:  $P_{w,r} \equiv \{v \in \mathbb{R}^q : v'w = r\}$ .
- $\mu_{w,r}$ : Lebesgue measure on  $P_{w,r}$ .
- $\mathcal{R}$ : Radon transform:  $\mathcal{R}[f](w, u) = \int_{P_{w,u}} f(v) d\mu_{w,u}(v)$ .

## B Proofs

**Proof of Theorem 2.1.** The proof of the theorem is immediate from Theorem 1 in BH (2013). We therefore give a brief sketch. By Assumptions 2.1 and 2.2, we note that there exists a function  $\psi : \mathbb{R}^{Jk_2} \times \mathbb{R}^J \times \mathbb{R}^J \rightarrow \mathbb{R}^J$  such that for some subvector  $\tilde{S}_t$  of  $S_t$ ,

$$\Xi_{jt} = \psi_j(X_t^{(2)}, P_t, \tilde{S}_t) - X_{jt}^{(1)}, \quad j = 1, \dots, J,$$

and by Assumption 2.3, the following moment condition holds:

$$E[\psi_j(X_t^{(2)}, P_t, \tilde{S}_t) - X_{jt}^{(1)} | Z_t, X_t] = 0.$$

Identification of  $\psi$  then follows from applying the completeness argument in the proof of Theorem 1 in BH (2013). □

**Proof of Theorem 3.1.** (i) First, under the linear random coefficient specification, the connected substitutes assumption in Berry, Gandhi, and Haile (2013) is satisfied. By Theorem 1 in Berry, Gandhi, and Haile (2013), Assumption 2.2 is satisfied. Then, by Assumptions 2.1-2.3 and Theorem 2.1,  $\psi$  is identified. Further, the aggregate demand  $\phi$  is identified by (2.10) and the identity  $\phi_0 = 1 - \sum_{j=1}^J \phi_j$ .

For any product  $j$  and product characteristics  $(x_j^{(2)}, p_j, \delta_j)$  define the new function

$$\tilde{\Phi}_j(x_j^{(2)}, p_j, \delta_j) = - \lim_{\delta_1, \dots, \delta_{j-1}, \delta_{j+1}, \dots, \delta_J \rightarrow -\infty} \phi_j(x^{(2)}, p, \delta)$$

point wise. Here  $\phi_j(x^{(2)}, p, \delta)$  can be any fixed vector of product characteristics where  $(x_j^{(2)}, p_j, \delta_j)$  coincide with the values on the l.h.s. of the equation. The limit on the r.h.s. exists and is unique. This can be seen by using the definition of  $\phi_j$ , Lebesgue's theorem, and Assumption 3.2 (i). Consequently,

$$\tilde{\Phi}(x_j^{(2)}, p_j, \delta_j) = - \int 1\{x_j^{(2)'}b^{(2)} + ap_j + \epsilon_j < -\delta_j\} f_{\vartheta_j}(b^{(2)}, a, e_j) d\vartheta_j.$$

Now define  $\Phi$  as in (3.4) and conclude

$$\begin{aligned} \Phi(w, u) &= - \int 1\{w'\theta < -u\} f_{\vartheta_j}(b^{(2)}, a, e_j) d\theta \\ &= - \int_{-\infty}^{-u} \int_{P_{w,r}} f_{\vartheta_j}(b^{(2)}, a, e_j) d\mu_{w,r}(b^{(2)}, a, e_j) dr = - \int_{-\infty}^{-u} \mathcal{R}[f_{\vartheta_j}](w, r) dr. \end{aligned} \quad (\text{B.1})$$

Taking a derivative with respect to  $u$  yields (3.6). By the assumption that the conditional distribution of  $\epsilon_{ijt}$  given  $(\beta_{it}^{(2)}, \alpha_{it})$  is identical for  $j = 1, \dots, J$ , it follows that  $f_{\vartheta_j} = f_{\vartheta}, \forall j$  for some common density  $f_{\vartheta}$ . Hence, we may rewrite (3.6) as

$$\frac{\partial \Phi(w, u)}{\partial u} = \mathcal{R}[f_{\vartheta}](w, u). \quad (\text{B.2})$$

Note that by Assumptions 3.1 (i) and 3.2 (ii),  $\partial \Phi(w, u)/\partial u$  is well-defined for all  $(w, u) \in \mathbb{H}_+ \times \mathbb{R}$ . This is because, under Assumption 3.2 (ii), for any  $(w, u) \in \mathbb{H}_+ \times \mathbb{R}$ , one can find  $(x_j, p_j, d_j) \in \bigcup_{k=1}^J \text{supp}(X_{kt}^{(2)}, P_{kt}, D_{kt})$  such that  $w = (x_j^{(2)}, p_j, 1)/\|(x_j^{(2)}, p_j, 1)\|$  and  $u = \delta_j/\|(x_j^{(2)}, p_j, 1)\|$ . The identification of  $f_{\vartheta}$  then follows from Assumption 3.1 (iii) and the injectivity of the Radon transform (Theorem I in Cramér and Wold, 1936).

(ii) In the first part of the proof  $f_{\vartheta_j}, j = 1, 2, \dots, J$  were identified (as  $f_{\vartheta}$ ). Hence, the conditional distribution  $f_{\epsilon_j|\beta^{(2)}, \alpha}$  of  $\epsilon_{ijt}$  given  $(\beta_{it}^{(2)}, \alpha_{it})$  and the marginal distribution  $f_{\beta^{(2)}, \alpha}$  of  $(\beta_{it}^{(2)}, \alpha_{it})$  are identified for any  $j$ . Under the additional assumption that  $\epsilon_{i1t}, \epsilon_{i2t}, \dots, \epsilon_{iJt}$  are independent conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$  we get the joint distribution of  $\theta_{it}$  by

$$f_{\theta}(b^{(2)}, \alpha, e_1, \dots, e_J) = \prod_{j=1}^J f_{\epsilon_j|\beta^{(2)}, \alpha}(e_j|b^{(2)}, \alpha) \times f_{\beta^{(2)}, \alpha}(b^{(2)}, \alpha). \quad (\text{B.3})$$

Hence,  $f_{\theta}$  is identified. □

The following lemma is used in the proof of Theorem 4.2.

**Lemma B.1.** *Suppose the Assumptions 2.1 and Condition 4.1 hold and that  $\phi_l$  is given as in Example 2 or Example 3 with  $l \in \tilde{\mathbb{L}} = \{(0, 1), (0, 0)\}$ . Then for all  $(x^{(2)}, p) = (x_1^{(2)}, x_2^{(2)}, p_1, p_2) \in \mathbb{R}^{2k}$  with  $(x_1^{(2)}, p_1) \neq (x_2^{(2)}, p_2)$  the function  $\phi : \mathbb{R}^{2k} \times \mathbb{R}^2 \rightarrow [0, 1]^2$  defined as*

$$\phi(x_1^{(2)}, x_2^{(2)}, p_1, p_2, d_1, d_2) \equiv \left[ \phi_{(0,0)}(x_1^{(2)}, x_2^{(2)}, p_1, p_2, d_1, d_2), \phi_{(0,1)}(x_1^{(2)}, x_2^{(2)}, p_1, p_2, d_1, d_2) \right]$$

*is invertible in  $(d_1, d_2)$  on any bounded subset of  $\mathbb{R}^2$ . This holds for other appropriate choices of  $\tilde{\mathbb{L}}$  as well (e.g.  $\tilde{\mathbb{L}} = \{(1, 0), (1, 1)\}$ ).*

**Proof of Lemma B.1.** We start with the observation that  $\phi_{(0,0)}(x^{(2)}, p, d)$  is monotonically decreasing in  $d_1$  and also in  $d_2$  while  $\phi_{(0,1)}(x^{(2)}, p, d)$  is monotonically decreasing in  $d_1$  and monotonically increasing in  $d_2$  by definition. Furthermore, the full support of  $\epsilon_1$  and  $\epsilon_2$  implies that  $\phi_{(0,0)}$  and  $\phi_{(0,1)}$  are strictly increasing or decreasing in  $d_1$  and  $d_2$

$$\frac{\partial \phi_{(0,0)}(x^{(2)}, p, d)}{\partial d_1} < 0, \quad \frac{\partial \phi_{(0,0)}(x^{(2)}, p, d)}{\partial d_2} < 0, \quad \frac{\partial \phi_{(0,1)}(x^{(2)}, p, d)}{\partial d_1} < 0, \quad \frac{\partial \phi_{(0,1)}(x^{(2)}, p, d)}{\partial d_2} > 0.$$

Hence, the determinant of the Jacobian of  $d \mapsto \phi(x^{(2)}, p, d)$  as well as their principle minors are strictly negative for all  $d \in \text{supp}(D)$

$$\det(J_\phi)(x, d) = \frac{\partial \phi_{(0,0)}(x^{(2)}, p, d)}{\partial d_1} \frac{\partial \phi_{(0,1)}(x^{(2)}, p, d)}{\partial d_2} - \frac{\partial \phi_{(0,1)}(x^{(2)}, p, d)}{\partial d_1} \frac{\partial \phi_{(0,0)}(x^{(2)}, p, d)}{\partial d_2} < 0.$$

Thus, on every rectangular domain in  $\mathbb{R}^2$  the assumptions of the Gale-Nikaido theorem are fulfilled. Since any bounded subset in  $\mathbb{R}^2$  is contained in some rectangular domain,  $\phi$  is invertible on any bounded subset of  $\mathbb{R}^2$ .  $\square$

**Proof of Theorem 4.2.** (a) First, let  $\tilde{\mathbb{L}} = \{(1, 0), (1, 1)\}$ . By Condition 4.1 and Lemma B.1, Assumption 2.2 is satisfied. By Assumptions 2.1-2.3 and Theorem 2.1,  $\psi$  is identified. Further, the aggregate demand  $\{\phi_l, l = (1, 0), (1, 1)\}$  is identified by Lemma B.1. Second, take  $\tilde{\mathbb{L}} = \{(0, 0), (0, 1)\}$ . Then by the same argument, the aggregate demand  $\{\phi_l, l = (0, 0), (0, 1)\}$  is identified as well. Hence, the entire aggregate demand vector  $\phi$  is identified.

Recall that the demand for bundle (0,0) satisfies (4.1). Together with Assumption 3.2 and

Lebesgue's theorem the limits

$$\begin{aligned}
\tilde{\Phi}_{(0,0),1}(x_1^{(2)}, p_1, \delta_1) &= - \lim_{\delta_2 \rightarrow -\infty} \phi_{(0,0)}(x^{(2)}, p, \delta) = - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 < -\delta_1\} f_\theta(b^{(2)}, a, e, \Delta) d\theta \\
&= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 < -\delta_1\} f_{\vartheta_1}(b^{(2)}, a, e_1) d\vartheta_1 \\
\tilde{\Phi}_{(0,0),2}(x_2^{(2)}, p_2, \delta_2) &= - \lim_{\delta_1 \rightarrow -\infty} \phi_{(0,0)}(x^{(2)}, p, \delta) = - \int 1\{x_2^{(2)'}b^{(2)} + ap_2 + e_2 < -\delta_2\} f_\theta(b^{(2)}, a, e, \Delta) d\theta \\
&= - \int 1\{x_2^{(2)'}b^{(2)} + ap_2 + e_2 < -\delta_2\} f_{\vartheta_2}(b^{(2)}, a, e_2) d\vartheta_2
\end{aligned}$$

exist and are unique. Note that in both equations  $\Delta$  and  $e_1$  or  $e_2$  are integrated out. Hence, the first equation connects  $f_{\vartheta_1}$  to  $\tilde{\Phi}_{(0,0),1}$  and the second equation connects  $f_{\vartheta_2}$  to  $\tilde{\Phi}_{(0,0),2}$ . Following the argumentation in the proof of Theorem 3.1 yields that  $f_{\vartheta_1}$  and  $f_{\vartheta_2}$  are identified.

As a second step we repeat the argument for  $\phi_{(1,1)}$ . The demand for bundle (1,1) can be written as (4.3). By taking the limits

$$\begin{aligned}
\tilde{\Phi}_{(1,1),1}(x_1^{(2)}, p_1, \delta_1) &= - \lim_{\delta_2 \rightarrow -\infty} \phi_{(1,1)}(x^{(2)}, p, \delta) \\
&= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta < -\delta_1\} f_\theta(b^{(2)}, a, e, \Delta) d\theta \\
&= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta < -\delta_1\} f_{\eta_1}(b^{(2)}, a, e_1 + \Delta) d\eta_1 \\
\tilde{\Phi}_{(1,1),2}(x_2^{(2)}, p_2, \delta_2) &= - \lim_{\delta_1 \rightarrow -\infty} \phi_{(1,1)}(x^{(2)}, p, \delta) \\
&= - \int 1\{x_2^{(2)'}b^{(2)} + ap_2 + e_2 + \Delta < -\delta_2\} f_\theta(b^{(2)}, a, e, \Delta) d\theta \\
&= - \int 1\{x_2^{(2)'}b^{(2)} + ap_2 + e_2 + \Delta < -\delta_2\} f_{\vartheta_2}(b^{(2)}, a, e_2 + \Delta) d\eta_2
\end{aligned}$$

and following the argument in the proof of Theorem 3.1 the identification of  $f_{\eta_1}$  and  $f_{\eta_1}$  is proven.

(b) With  $f_{\eta_j}$  for  $j = 1, 2$  the characteristic function  $\Psi_{\Delta + \epsilon_j | (\beta^{(2)}, \alpha)}$  of  $(\Delta_{it} + \epsilon_{ijt})$  conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$  is identified as well. With the conditional independence assumption  $\Delta_{it} \perp \epsilon_{ijt} | (\beta_{it}^{(2)}, \alpha_{it})$  and  $\Psi_{\epsilon_j | (\beta^{(2)}, \alpha)}(t) \neq 0$  for almost all  $t \in \mathbb{R}$  the densities  $f_{\eta_j}$  and  $f_{\vartheta_j}$  can be disentangled by the deconvolution:

$$f_{\Delta | \beta^{(2)}, \alpha} = \mathcal{F}^{-1} \left( \frac{\Psi_{\Delta + \epsilon_j | (\beta^{(2)}, \alpha)}}{\Psi_{\epsilon_j | (\beta^{(2)}, \alpha)}} \right),$$

where  $\mathcal{F}$  denotes the Fourier transform with respect to  $\Delta$ . This obviously identifies  $f_{\beta^{(2)}, \alpha, \Delta}$  as well. If in addition  $\epsilon_{i1t}$  and  $\epsilon_{i2t}$  are independent conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$ , the density of  $f_\theta$  is

identified by

$$f_{\theta}(b^{(2)}, a, e, \Delta) = f_{\epsilon_1|\beta^{(2)},\alpha}(e_1|b^{(2)}, a) f_{\epsilon_2|\beta^{(2)},\alpha}(e_2|\beta^{(2)}, \alpha) f_{\beta^{(2)},\alpha,\Delta}(b^{(2)}, a, \Delta)$$

This completes the proof of the theorem.  $\square$

**Proof of Theorem 4.3.** First, let  $\tilde{\mathbb{L}} = \{(2, 0), (2, 1)\}$ . By Condition 4.1 and Lemma B.1, Assumption 2.2 is satisfied. By Assumptions 2.1-2.3 and Theorem 2.1,  $\psi$  is identified. This implies that the aggregate demand  $\{\phi_l, l = (2, 0), (2, 1)\}$  is identified. Second, take  $\tilde{\mathbb{L}} = \{(0, 0), (0, 1)\}$ . Then by the same argument, the aggregate demand  $\{\phi_l, l = (0, 0), (0, 1)\}$  is identified as well. Again by Condition 4.1, we can take the limits

$$\begin{aligned} \tilde{\Phi}_{(0,0)}(x_1^{(2)}, p_1, \delta_1) &= - \lim_{\delta_2 \rightarrow -\infty} \phi_{(0,0)}(x^{(2)}, p, \delta) \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 < -\delta_1\} f_{\theta}(b^{(2)}, a, e, \Delta) d\theta \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 < -\delta_1\} f_{(\beta^{(2)},\alpha,\epsilon_1)}(b^{(2)}, a, e_1) d\theta \\ \tilde{\Phi}_{(0,1)}(x_1^{(2)}, p_1, \delta_1) &= - \lim_{\delta_2 \rightarrow \infty} \phi_{(0,1)}(x^{(2)}, p, \delta) \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(1,1)} > -\delta_1\} f_{\theta}(b^{(2)}, a, e, \Delta_{(1,1)}) d\theta \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(1,1)} > -\delta_1\} f_{(\beta^{(2)},\alpha,\epsilon_1+\Delta_{(1,1)})}(b^{(2)}, a, e + \Delta_{(1,1)}) d\theta \\ \tilde{\Phi}_{(2,0)}(x_1^{(2)}, p_1, \delta_1) &= - \lim_{\delta_2 \rightarrow -\infty} \phi_{(2,0)}(x^{(2)}, p, \delta) \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(2,0)} > -\delta_1\} f_{\theta}(b^{(2)}, a, e, \Delta_{(2,0)}) d\theta \\ &= - \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(2,0)} > -\delta_1\} f_{(\beta^{(2)},\alpha,\epsilon_1+\Delta_{(2,0)})}(b^{(2)}, a, e_1 + \Delta_{(2,0)}) d\theta \\ \tilde{\Phi}_{(2,1)}(x_1^{(2)}, p_1, \delta_1) &= - \lim_{\delta_2 \rightarrow \infty} \phi_{(2,1)}(x^{(2)}, p, \delta) \\ &= \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(2,1)} - \Delta_{(1,1)} > -\delta_1\} f_{\theta}(b^{(2)}, a, e, \Delta_{(1,1)}) d\theta \\ &= \int 1\{x_1^{(2)'}b^{(2)} + ap_1 + e_1 + \Delta_{(2,1)} - \Delta_{(1,1)} > -\delta_1\} f_{(\beta^{(2)},\alpha,\epsilon_1+\Delta_{(2,1)}-\Delta_{(1,1)})}(b^{(2)}, a, e_1 + \Delta_{(2,1)} - \Delta_{(1,1)}) d\theta . \end{aligned}$$

By the argument in the proof of Theorem 3.1 and the assumption that  $(X_{1t}^{(2)}, P_{1t}, D_{1t})$  has a full support, this identifies the joint densities of  $(\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t})$ ,  $(\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t} + \Delta_{i,(1,1),t})$ ,  $(\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t} + \Delta_{i,(2,0),t})$ , and  $(\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t} + \Delta_{i,(2,1),t} - \Delta_{i,(1,1),t})$  respectively.

In what follows, the arguments are made conditional on  $(\beta_{it}^{(2)}, \alpha_{it})$  unless otherwise noted. By Assumption 4.1 (i), we may disentangle the distribution of  $\epsilon_{i1t}$  with that of  $\Delta_{i,(1,1),t}$ ,  $\Delta_{i,(2,0),t}$ , and  $\Delta_{i,(2,1),t} - \Delta_{i,(1,1),t}$  respectively by deconvolution as done in the proof of Theorem 4.2.



Thus, the marginal densities of  $\Delta_{i,(2,1),t} - \Delta_{i,(1,1),t}$  and  $\Delta_{i,(1,1),t}$  are identified. Further, we note that  $\Delta_{i,(2,1),t} - \Delta_{i,(1,1),t}$  is a convolution of  $\Delta_{i,(2,1),t}$  and  $-\Delta_{i,(1,1),t}$ . By Assumption 4.1 (ii), Proposition 8 of Carrasco and Florens (2010) applies. Hence, the marginal density of  $\Delta_{i,(2,1),t}$  is identified. By Assumption 4.1 (i),  $\Delta_{i,(1,1),t} \perp \Delta_{i,(2,0),t} \perp \Delta_{i,(2,1),t}$  conditional on  $(\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t})$ , and each of the marginal densities was identified in the previous step. Therefore, the joint density  $f_{(\Delta_{(1,1)}, \Delta_{(2,0)}, \Delta_{(2,1)}) | (\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t})}$  is identified as the product of the marginal densities. Since the density of  $(\beta_{it}^{(2)}, \alpha_{it}, \epsilon_{i1t})$  is identified as well, we may identify the joint density  $f_{\vartheta_1}$  as  $f_{\vartheta_1} = f_{(\Delta_{(1,1)}, \Delta_{(2,0)}, \Delta_{(2,1)}) | (\beta^{(2)}, \alpha, \epsilon_1)} f_{(\beta^{(2)}, \alpha, \epsilon_1)}$ .  $f_{\vartheta_2}$  is identified as  $f_{\vartheta_1}$  by Assumption 4.1 (ii). By Assumption 4.1 (ii) and arguing as in (B.3),  $f_{\theta}$  is identified. Given  $f_{\theta}$ , all components of  $\phi$  is identified. This completes the proof of the theorem.  $\square$

**Proof of Theorem 4.4.** First, under the linear random coefficient specification, the connected substitutes assumption in Berry, Gandhi, and Haile (2013) is satisfied. By Theorem 1 in Berry, Gandhi, and Haile (2013), Assumption 2.2 is satisfied. Then, by Assumptions 2.1-2.3 and Theorem 2.1,  $\psi$  is identified. Further, the aggregate demand  $\phi$  is identified by (2.10) and the identity  $\phi_0 = 1 - \sum_{j=1}^J \phi_j$ . By Assumption 4.2, for each  $(x^{(2)}, p, \delta)$  and  $\mathcal{J} \subseteq \{1, \dots, J\} \setminus \{j\}$ ,  $\mathcal{M}_{\mathcal{J}}(x^{(2)}, p, \delta)$  is in the support of  $(X_t^{(2)}, P_t, D_t)$ . Hence, one may construct

$$\tilde{\Phi}_j(x_j^{(2)}, p_j, \delta_j) = \sum_{\mathcal{J} \subseteq \{1, \dots, J\} \setminus \{j\}} \phi_j \circ \mathcal{M}_{\mathcal{J}}(x^{(2)}, p, \delta) = \int 1\{x_j^{(2)'} b^{(2)} + ap_j < -\delta_j\} f_{\theta}(b^{(2)}, a) d\theta, \quad (\text{B.4})$$

where the second equality follows because of the following. First,  $\mathcal{M}_{\mathcal{J}}$  replaces the indicators in  $\phi_j$  of the form  $1\{(x_j^{(2)} - x_i^{(2)})' b^{(2)} + a(p_j - p_i) < -(\delta_j - \delta_i)\}$  with  $1\{(x_j^{(2)} - x_i^{(2)})' b^{(2)} + a(p_j - p_i) > -(\delta_j - \delta_i)\}$  for  $i \in \mathcal{J}$ . The random coefficients are assumed to be continuously distributed. We therefore have

$$1\{(x_j^{(2)} - x_i^{(2)})' b^{(2)} + a(p_j - p_i) < -(\delta_j - \delta_i)\} + 1\{(x_j^{(2)} - x_i^{(2)})' b^{(2)} + a(p_j - p_i) > -(\delta_j - \delta_i)\} = 1, \quad a.s.$$

Therefore,  $\sum_{\mathcal{J} \subseteq \{1, \dots, J\}} \phi_j \circ \mathcal{M}_{\mathcal{J}}(x^{(2)}, p, \delta) = 1$ . Since  $\tilde{\Phi}_j$  is constructed by summing  $\phi_j \circ \mathcal{M}_{\mathcal{J}}$  over subsets of  $\{1, \dots, J\}$  except  $\{j\}$ , we are left with the integral of the single indicator function  $1\{x_j^{(2)'} b^{(2)} + ap_j < -\delta_j\}$  with respect to  $f_{\theta}$ . This ensures (B.4). Now define  $\Phi$  as in (3.4). Then,

it follows that

$$\begin{aligned}\Phi(w, u) &= - \int 1\{w'\theta < -u\} f_\theta(b^{(2)}, a) d\theta \\ &= - \int_{-\infty}^{-u} \int_{P_{w,r}} f_\theta(b^{(2)}, a) d\mu_{w,r}(b^{(2)}, a) dr = - \int_{-\infty}^{-u} \mathcal{R}[f_\theta](w, r) dr .\end{aligned}\quad (\text{B.5})$$

Taking a derivative with respect to  $u$  then yields (3.6). Note that by Assumption 4.2 (ii),  $\bigcup_{j=1}^J (X_{jt}^{(2)}, P_{jt}, D_{jt}) = \mathbb{R}^{d_X-1} \times \mathbb{R} \times \mathbb{R}$ . This implies that  $\partial\Phi(w, u)/\partial u$  is well-defined for all  $(w, u) \in \mathbb{H}_+ \times \mathbb{R}$ . The conclusion of the theorem then follows from Assumption 3.1 (iii), injectivity of the Radon transform, and  $\Phi$  being identified.  $\square$



