

Original article:

**IDENTIFICATION OF PROTEIN TYPES IN BAMBARA NUT SEEDS:
PERSPECTIVES FOR DIETARY PROTEIN SUPPLY
IN DEVELOPING COUNTRIES**

J. Okpuzor^{1*}, HA. Ogbunugafor², U. Okafor³, MO. Sofidiya⁴

¹. Department of Cell Biology and Genetics, Faculty of Science, University of Lagos, Akoka, Lagos

². Applied Biochemistry Department, Faculty of Natural Sciences, Nnamdi Azikiwe University, Awka, Nigeria

³. Department of Biochemistry, College of Medicine, University of Lagos, Idi-Araba, Lagos, Nigeria

⁴. Department of Pharmacology, College of Medicine, University of Lagos, Idi-Araba, Lagos, Nigeria

* Corresponding author: J.Okpuzor; email: joyokpuzor@yahoo.com; Phone: +234 802 3065999

ABSTRACT

This study aims to identify the types of proteins in malted and dry Bambara groundnut seeds and through a comparative analysis, identify similarities and their known uses. Dry viable bambara seed was stored for five days to malt. The proteins in the dry and malted seed were subsequently extracted in potassium phosphate buffer pH 7.0 and precipitated with saturated ammonium sulphate. MudPit (multidimensional protein identification technology) and LC-MALDI TOF-TOF (liquid chromatography - matrix-assisted laser desorption ionization tandem time-of-flight) mass spectrometry were thereafter used to identify the different types of proteins. A total of ten and twelve different types of proteins present in other legume species were identified in the malted and dry seeds respectively from the 214 peptides isolated after searching 586 proteins of the genus *Vigna*. Seed storage protein B and vicilin were observed to be the major proteins common to both malted and dry seeds and are similar to *Vigna luteola*. Some of the other proteins observed showed amino acid sequence homology with *Vigna radiata* and *Vigna unguiculata* species. The following proteins BV1, Heat shock and Bowman-Birk Inhibitor (a protease), were observed only in the malted state. This information may enhance the appreciation of the nutritional and health benefits of the seed.

Keywords: *Voandzeia subterranea* L. Verdc., Malted Seed, MudPit, LC-MALDI TOF-TOF mass spectrometry, *Vigna species*, Peptides

INTRODUCTION

Bambara groundnut (*Voandzeia subterranea* L. Verdc) is a seed crop of African origin used locally as a vegetable or snack. It is cultivated principally by farmers as a "famine culture" crop because it has several natural

agronomic advantages including high nutritional value, drought tolerance and the ability to produce in soils considered insufficiently fertile for cultivation of other more favoured species such as common beans and groundnuts (*Arachid hypogea*) (Anchirinah et al., 2001; Azam-Ali et al., 2001). Despite these

numerous advantages, it is unfortunately, one of the neglected and under-utilized crops in sub Saharan Africa. The seed is regarded as a balanced food because when compared to most food legumes, it is rich in iron and the protein contains high lysine and methionine (Adu-Dapaah and Sangwan, 2004). In addition, bambara groundnut is known to contain 63 % carbohydrates, 18 % oil and the fatty acid content is predominantly linoleic, palmitic and linolenic acids (Minka and Bruneteau, 2000).

In view of world food shortages caused by drought, wars and inadequate government attention to agriculture especially in developing countries, it becomes expedient to harness proteins from all available food sources to minimise food and nutritional crises. As observed by Padulosi et al. (2002), neglected and underutilized crops could play prominent roles in sustaining the impoverished rural African populations by increasing their available food and protein basket.

Bambara groundnut is eaten in several ways and at different stages of maturation. The young fresh seeds may be boiled and eaten as a snack in a manner similar to boiled peanuts and could be made into pudding locally called *Moi Moi* or *Okpa* (bean porridge) in the some parts of Nigeria. It has been reported that in Zambia, bambara groundnut is used for bread making (Brough et al., 1993) while Poulter and Caygill, (2006) also reported that it could be used for milk making.

Wazael (2004) described the great genetic diversity potential of bambara groundnut. Although, the types of proteins available in other species have been studied and reported, to the best of our knowledge, there has been no documented information on the types of protein in either the dry or the malted bambara groundnut (*Voandzeia subterranea* L. Verdc).

The goal of this investigation is to identify the different types of proteins in the

malted and dry seeds of *Voandzeia subterranea* L. Verdc, their similarities to other known vigna species and the various uses which are expected to enhance the appreciation of the nutritional and health values of this seed.

MATERIALS AND METHODS

Source of bambara groundnut

Bambara groundnuts (*V. subterranea*) were purchased in April, 2008 from a local market in Lagos metropolis and identified in the Department of Botany & Microbiology, University of Lagos, Akoka, Yaba, Lagos, Nigeria.

Malting of bambara groundnut

Fifty grams of bambara groundnuts were washed with distilled water and spread in a tray lined with three layers of Whatman filter paper No 3. The seeds were then covered completely with the filter paper and allowed to stand on the laboratory bench for five days at room temperature (37 ± 2 °C) for the seed to malt.

Ammonium sulphate precipitation

Twenty grams of malted bambara groundnuts were macerated and extracted in 50 ml potassium phosphate buffer, pH 7.0 using an electric blender, original millennium Nakai (blender 248) special. The macerate was centrifuged at 10,000 rpm for 10 mins and the supernatant decanted. The resulting suspension which is the crude protein extract was subjected to saturated ammonium sulphate precipitation. The precipitate was dialyzed overnight in phosphate buffer pH 7.0 at 4 °C.

Sample preparation for LC-MALDI.

Digestion and sample preparation

A 50 μ L aliquot of the sample was centrifuged at 4000 rpm for 5 mins. The supernatant was extracted and the pellet washed with digestion buffer (50 mM ammonium bicarbonate). The digestion buffer wash was combined with the original supernatant and subjected to denaturation and proteolytic digestion using trypsin. Samples collected were lyophilized and dissolved in 100 μ L of 50 mM ammonium bicarbonate in 20 % acetonitrile for trypsin digestion. The samples were then reduced and alkylated with 10 μ L of 250 mM DTT (60 min / 55 °C) and 10 μ L of 625 mM iodoacetamide (60 min/room temperature / in the dark). Proteolytic digestion was performed in 50 mM ammonium bicarbonate buffer using a trypsin to protein ratio of 1:100. The digestion was carried out overnight at 37 °C. The digests were cleaned by repeated lyophilizing and reconstituting in a 0.1 M acetic acid solution. After final lyophilization, the digests were reconstituted in strong cation exchange (SCX) loading buffer (5 mM ammonium formate in 20 % acetonitrile, pH 3).

Strong cation exchange fractionation

After digestion, the protein samples were fractionated using SCX ProteaTip spin Tips. The tips were first washed to wet the packing material by adding 50 μ L of strong cation exchange loading buffer and centrifuging the system at 4000rpm for 3 mins. The sample was then loaded in the Spin Tip and centrifuged at 4000rpm for 3 min after which it was washed to elute salts and other non-retained components by adding 50 μ L of the rinse solution (5 mM ammonium format in 20 % acetonitrile) to the top of the Spin Tip. The Spin Tip was transferred to a new clean centrifuge tube to collect the sample during elution with 200 μ L of elution solution. Eight different elution solutions were used to fractionate the peptides and these were 20, 60,

100, 150, 200, 250, 400, 500 mM ammonium formate in 20 % acetonitrile. The collected fractions were cleaned by repeated lyophilizing and reconstituting in a 0.1 M acetic acid solution. After final lyophilization, the digests were reconstituted in LC run buffer.

LC-MALDI spotting

Lyophilized digested samples were reconstituted in 20 μ L of 0.1 % TFA in de-ionized water and 5 μ L corresponding to 25 % of the sample (5 μ L) was injected into ABI Tempo LC MALDI (Proteabio Inc.) with a Separating column - Merck Chromolith CapRod monolith column – 150 X 0.1 mm with separation gradient of 15 mins.

Mass spectrometry and data analysis

Mass spectrometric analysis was performed using an ABI 4800 MALDI TOF/TOF analyzer in positive ion reflector mode. Data was collected over a mass range of $m/z = 850 - 4000$. For MS acquisition, a total of 400 laser shots per spectrum with an S/N threshold of 10. For MS/MS acquisition, the 15 strongest precursors chosen for MS/MS with S/N threshold were set at 30 for MSMS precursors. Each MALDI spot was interrogated until at least 4 peaks in the MSMS spectra achieved an S/N of 70. The MS/MS data acquired data was processed with Protein ProteinPilot software 2.0 (Applied Biosystems) using the Paragon and Pro Group search algorithms. The search parameter for sample type was set for identification, trypsin was set as the digestion enzyme and the search database was the latest NCBI nr database. *Vigna* was selected as the species and other special factors were cysteine alkylation and I.D focus on biological modifications. The search mode was set to thorough identification, and detected protein confidence threshold was set at 66 %.

RESULTS

A total of ten and twelve different proteins were identified in the malted and dry seeds of bambara groundnut respectively from 214 peptides after searching 586 proteins of the genus *Vigna*. Of the twenty two different proteins identified, B storage protein and vicilin were the proteins most common to the malted and dry seeds.

The Swissprot protein database containing 586 proteins of the genus *vigna* which was used for this investigation did not contain any *V. subterranea* proteins. The latest NCBI nr database also contained only but a few. Most of the proteins identified in both conditions of the seed had amino acid homology to other known species of *Vigna*.

Malted seed

Ten proteins were identified in the malted state of the seed (Table 1). The major protein observed is the Seed storage protein B which exhibited a total protscore of 52.03 and percentage coverage of approx 43 that was found to be similar to that of *Vigna luteola*. Forty percent of the ten proteins identified showed amino acid sequence homology to *Vigna radiata* species, twenty percent is similar to *Vigna unguiculata* and only ten percent of the proteins were identical to *Vigna angularis* and *Vigna luteola*. The B storage protein of the latter incidentally, possesses the highest unused protscore (measure of the protein confidence for a detected protein, calculated from the peptide confidence for peptides from spectra that have not already been completely “used” by higher scoring proteins) in malted seed. However, the other proteins BV1, Heat shock protein, 8S globulin alpha isoform and Chain I, Crystal Structure of The Bowman-Birk Inhibitor protein had protscore (measure of the total amount of evidence for a detected protein) lower than the cut-off threshold of 0.47 used

at 66 % confidence (the confidence for the peptide identification, expressed as a percentage), but presented fairly good percentage coverage (the number of amino acids matching at least one identified peptide divided by the total number of amino acids in the protein sequence, expressed as a percentage (Table 2)). Interestingly, the peptide sequences of Chain I, Crystal Structure of The Bowman-Birk Inhibitor protein had low percentage coverage of only 25 % but exhibited high confidence level of 83 % indicating the accuracy of protein identification process.

Dry seed

In the dry seeds, twelve proteins were identified (Table 3). Three had protscore above 19 at 66 % confidence level with percentage coverage above 32. The other types of protein however, had lower protscores ranging between 0.96-14.0 and percentage coverage of between 8 and 38.

Seven identified proteins of *V. subterranea*, 8S globulin alpha subunit, Mung bean seed albumin, Em protein, ARG10, PDF1, 10 kDa protein precursor (Clone PSAS10) and Actin had sequence homology to *Vigna radiata* specie. Two proteins, Chain A, Crystal Structure of Adzuki Bean 7s Globulin-3 and 7S globulin-2 had amino acid sequence similar to *Vigna angularis* while another two, Dehydrin and vicilin was similar to *Vigna unguiculata*. Only one protein type (Seed storage protein B) was similar to *Vigna luteola* specie. The identified peptide sequences showed that some proteins with low protscores had high percentage confidence level (Table 4).

Table 1: Proteins identified in malted bambara groundnut (*Vigna subterranea*)

No	Name	Species	Unused Score	Total Score	% Coverage	Accession
1	Seed storage protein B	<i>Vigna luteola</i>	52.03	52.03	43.94	gi 70672852
2	8S globulin alpha subunit	<i>Vigna radiata</i>	15.60	25.16	46.04	gi 158251953
3	8S globulin beta isoform precursor	<i>Vigna radiata</i>	9.18	29.23	32.89	gi 108743976
4	Chain A, crystal structure of adzuki bean 7s globulin-1	<i>Vigna angularis</i>	4.32	22.37	34.56	gi 167013178
5	10 kDa protein precursor (Clone PSAS10)	<i>Vigna radiata</i> var. radiata	4.16	4.16	38.67	gi 112671
6	Vicilin protein	<i>Vigna unguiculata</i>	2.00	19.68	37.41	gi 160332746
7	BV1 protein	Mungbean yellow mosaic virus Vigna [KA27]	0.32	0.32	7.42	gi 8272630
8	Chain I, crystal structure of the Bowman-Birk inhibitor from <i>V. unguiculata</i> seeds in complex with beta-trypsin at 1.55 angstroms resolution	<i>Vigna unguiculata</i>	0.28	2.77	25.30	gi 122920306
9	Heat shock protein 90	<i>Phytophthora melonis</i>	0.20	0.20	5.67	gi 156986798
10	8S globulin alpha' isoform precursor	<i>Vigna radiata</i>	0.06	22.71	34.44	gi 108743974

Table 2: Peptide sequence of malted *Vigna subterranea*

% Coverage	Accessions	Names	Contribution	% Confidence	Sequence
43.94	gi 70672852	Seed storage protein B [<i>Vigna luteola</i>]	2.00	99.0	QEEESWQVQR
			2.000	99.00	QIQNLENYRVVEFK
			2.000	99.00	QIQNLENYR
			2.000	99.00	QEEESWQVQR
			2.000	99.00	NQYGHIR
			2.000	99.00	NQKPIYSNKFGR
			2.000	99.00	NQKPIYSNK
			2.000	99.00	LQRFDQR
			2.000	99.00	IKKQSQSHFVDAQPDEQEKGFR
			2.000	99.00	LIKKQSQSHFVDAQPDEQEKGFR
			2.000	99.00	LIKKQSQSHFVDAQPDEQEKGFR
			2.000	99.00	KQSQSHFVDAQPDEQEKGFR
			2.000	99.00	KQSQSHFVDAQPDEQEKGFR
			2.000	99.00	KQSQSHFVDAQPDEQEKGFR
			2.000	99.00	KQSQSHFVDAQPDEQEKGFR
			2.000	99.00	KQSQSHFVDAQPDEQEKGFR
			2.000	99.00	KQSQSHFVDAQPDEQEKGFR
			2.000	99.00	KQSQSHFVDAQPDEQEKGFR
			2.000	99.00	KQSQSHFVDAQPDEQEKGFR
			1.155	93.00	LRNQKPIYSNK
0.638	77.00	QQEESWQVQR			
46.04	gi 158251953 gi 108743972	8S globulin alpha subunit [<i>Vigna radiata</i>]	2.000	99.00	VPVNNPHRF
			2.000	99.00	SKQMQLNENYR
			2.000	99.00	QQKQEQEEQEESEWVQR
			2.000	99.00	QMQLNENYR
			2.000	99.00	LAVPVNNPHRF
			2.000	99.00	GKNNPFYFNDR
			1.699	98.00	LAVPVNNPHR
1.398	96.00	AVPVNNPHR			
32.89	gi 108743976	8S globulin beta isoform precursor [<i>Vigna radiata</i>]	2.000	99.00	QSESHFVDAQPEQQQR
			2.000	99.00	KQSESHFVDAQPEQQQR
			2.000	99.00	KIPAGTTFFLVNPNNDNLR
			1.699	98.00	DSRGQNNPFYFNDR
1.398	96.00	SDSRGQNNPFYFNDR			
34.56	gi 167013178	Chain A, crystal structure of adzuki bean 7s globulin-1	2.000	99.00	QEEESLEVQR
			2.000	99.00	QMQLNENYRVVEFK
			0.276	99.00	EQIQELMK
38.67	gi 112671	10 kDa protein precursor (Clone PSAS10)	2.000	99.00	NKEHLLSGR
			2.000	99.00	CWCTRNC
37.41	gi 160332746	Vicilin protein [<i>Vigna unguiculata</i>]	2.000	99.00	LHEITPEKNPQLR
7.42	gi 8272630	BV1 protein mungbean yellow mosaic virus - <i>Vigna</i> [KA27]	0.319	52.00	GIDGDGRSR
25.30	gi 122920306	Chain I, crystal structure of the Bowman-Birk Inhibitor from <i>Vigna Unguiculata</i> seeds in complex with beta-trypsin at 1.55 angstroms resolution	0.277	83.00	CSDVRLNSCHSACK
5.67	gi 156986798	Heat shock protein 90 [<i>Phytophthora melonis</i>]	0.162	31.08	KEVLDDKVEK
34.44	gi 108743974	8S globulin alpha' isoform precursor [<i>Vigna radiata</i>]	0.054	11.73	EQQQQQQDER

Table 3: Proteins identified in dry bambara groundnut (*Vigna subterranea*)

No	Protein	Species	Accession	Unused Score	Total Score	% Coverage
1	Seed storage protein B	<i>Vigna luteola</i>	gi 70672852	38.06	38.06	48.74
2	Chain A, crystal structure of adzuki bean 7s globulin-3	<i>Vigna angularis</i>	gi 167013181	20.11	24.27	64.20
3	Dehydrin	<i>Vigna unguiculata</i>	gi 6358640	10.04	10.05	65.64
4	8S globulin alpha subunit	<i>Vigna radiata</i>	gi 158251953	7.46	14.5	51.76
5	Mung bean seed albumin	<i>Vigna radiata</i> var. radiata	gi 1000708	3.70	3.70	13.97
6	Em protein	<i>Vigna radiata</i>	gi 1141784	2.00	2.00	26.26
7	ARG10	<i>Vigna radiata</i>	gi 2970051	2.00	2.00	18.99
8	PDF1	<i>Vigna radiata</i>	gi 18146788	1.70	1.70	38.67
9	10 kDa protein precursor (Clone PSAS10)	<i>Vigna radiata</i> var. radiata	gi 112671	1.52	1.52	30.67
10	Vicilin protein	<i>Vigna unguiculata</i>	gi 160332746	1.50	19.44	64.43
11	Actin	<i>Vigna radiata</i>	gi 6934188	0.96	0.96	8.75
12	7S globulin-2	<i>Vigna angularis</i>	gi 145207915	0.32	14.28	60.60

Table 4: Peptide sequence of dry *Vigna subterranea*

% Coverage	Accessions	Name	Contribution	% Confidence	Sequence
48.74	gi 70672852	Seed storage protein B [<i>Vigna luteola</i>]	2.00	99.00	VNPNDNDNLR
			2.000	99.00	NQYGHIR
			2.000	99.00	NQKPIYSNKFGR
			2.000	99.00	NQKPIYSNK
			2.000	99.00	NFLAGEKDNVMSEIP
			2.000	99.00	LVNPNDNDNLR
			2.000	99.00	KQSQSHFVDAQPDEQEKG
			2.000	99.00	KQSQSHFVDAQPDEQEK
			2.000	99.00	IPAGTTFFLVNPNDNDNLR
			2.000	99.00	FQDFFLSSTEAAQQSYLQGF
			2.000	99.00	FLVNPNDNDNLR
			2.000	99.00	EQQQQQEESWQVQR
			2.000	99.00	DNVMSEIPTEVLDVTFPASGEK
			2.000	99.00	ALLTLVNPDGR
			1.301	95.00	TLSEDEPFNLR
			1.046	91.00	PNDNDNLR
			0.699	80.00	NFLAGEKDNVMSEIPTEVLDVT FPASGEKVEK
			0.477	66.67	PAGTTFFLVNPNDNDNLR

Table 4 (cont.): Peptide sequence of dry *Vigna subterranean*

% Coverage	Accessions	Name	Contribution	% Confidence	Sequence
64.20	gi 167013181	Chain A, crystal structure of adzuki bean 7s globulin-3	2.000	99.00	TNQYGHLR
			2.000	99.00	TNQYGHLR
			2.000	99.00	QQQQGEESR
			2.000	99.00	QESQEESSDR
			2.000	99.00	NILEASFDSDFK
			2.000	99.00	KQSESHFVDAQPEQQQR
			2.000	99.00	HREHQESQEESSDR
			2.000	99.00	GINAENNQR
			2.000	99.00	EHQESQEESSDR
			1.699	98.00	HQESQEESSDR
51.76	gi 158251953; gi 108743972	8S globulin alpha subunit of adzuki bean 7s globulin-3 precursor [<i>Vigna radiata</i>]	1.699	98.00	ELSSQDEPFNLR
			0.602	75.00	REHQESQEESSDR
			2.000	99.00	QMQLNENYR
			2.000	99.00	GKNNPFYFNSTR
			1.699	99.00	NQKPIYSNKL
64.43	gi 160332746	Vicilin protein [<i>Vigna unguiculata</i>]	0.886	87.00	SKQMQLNENYR
			0.456	65.00	WFEITPEKNPQLR
			0.357	56.00	SLSTQNEPFNLR
			0.301	99.00	RNQYGHLR
			0.292	49.00	EGLLMPNYNSK
60.60	gi 145207915	7S globulin-2 [<i>Vigna angularis</i>]	0.244	45.00	GQNNPFYFSDSR
			0.229	41.00	GHQESQEESEPR
			0.319	52.00	NEWGHIR
			2.000	99.00	LQYTPGKTEDKILTNR
13.97	gi 1000708	Mung bean seed albumin [<i>Vigna radiata</i> var. <i>radiata</i>]	1.699	98.00	SNLPYINAAFR
			2.000	99.00	TTGGFTGGDTGLGGPYVGA
65.64	gi 6358640	Dehydrin [<i>Vigna unguiculata</i>]	2.000	99.00	NTADTGTGPR
			2.000	99.00	TTGGFTGGDTGLGGPYVGA
			2.000	99.00	NTADTGTGPR
			2.000	99.00	QYGTGGFTGDTGR
			2.000	99.00	QHGTGGFTGDTGR
			2.000	99.00	QHGTIGDTGR
			2.000	99.00	GTTGGFTGGDTGLGGPYVGA
8.75	gi 6934188	Actin [<i>Vigna radiata</i>]	0.959	89.00	ANTADTGTGPR
18.99	gi 2970051	ARG10 [<i>Vigna radiata</i>]	2.000	99.00	AGFAGDDAPR
30.67	gi 112671	10 kDa protein precursor (Clone PSAS10)	1.523	97.00	CYENPKNK
38.67	gi 18146788 gi 145579401	PDF1 [<i>Vigna radiata</i>] Chain A, Nmr solution structure of <i>Vigna Radiata</i> defensin 2 (Vrd2)	1.523	97.00	GPCFTTGSCDDHCKNKEH
			1.699	98.00	LLSGR
26.26	gi 1141784	Em protein [<i>Vigna radiata</i>]	2.000	99.00	CENLANTYR
					QQNKQELDER

DISCUSSION

In this study, we investigated the types of proteins present in the mature malted and dry seeds of *Vigna subterrenea*. Bambara groundnut is an important legume that could become a major source of dietary protein in developing countries whose staple foods are predominantly carbohydrates. Unfortunately however, bambara nut is eaten mostly by a small proportion of the population especially, in the Eastern part of Nigeria.

Ten different types of proteins were identified from malted while twelve were found in the dry seeds. The difference in the number of proteins observed in the two states of the bambara seed may suggest that the dryseeds require additional proteins to cope with biological processes. In both conditions of the seed however, only five of the proteins were identical and these are Seed storage protein B, vicilin, beta and alpha isoforms of 8S globulin, and 10kDa protein precursors. The other proteins observed varied according to the condition of the seeds. Østergaard et al. (2002) observed that there were cultivar differences in the 2-D gel spot patterns of proteins at the mature barley seeds and the malt. Only five protein types were identical in the two conditions emphasising that each state is unique and develops the requirements that enables it to cope with biological demands.

A viral protein with homology to Mung bean yellow mosaic virus (KA27), identified in the malted sample might be from a virus known to infect seeds of the *vigna* species (Garrido-Ramirez et al., 2000). The BV1 gene of the bipartite Begomovirus genome which encodes a nuclear shuttle protein (NSP) is also an avirulence determinant present in common bean (Zhou et al., 2007). The virus infection might have occurred during the malting process of *Vigna subterrenea*. Heat shock proteins identified in this study had amino acid homology with that found in *Phytophthora melonis*. Heat shock proteins (HSPs) are expressed in response to heat and other forms of stress (Queitsch et al., 2000) and are known to be responsible

for protein folding, assembly, translocation and degradation in many normal cellular processes. They also assist in stabilizing proteins and membranes and under stress conditions, can assist in protein refolding (Wang et al., 2004). It is our thinking that during the malting process of bambara seeds, there was a form of stress which was expressed by the appearance of heat shock proteins. This stress could be in form of excess or shortage of water or infection. Infection is likely to be the case; therefore, this protein may be transient because it has a very small percentage coverage and low confidence level.

It is likely that the malted sample rather than increase or decrease its seed nitrogen, produced other protein types like Chain A, crystal structure of adzuki bean 7s globulin-1 and Chain I, crystal structure of the Bowman-Birk Inhibitor. Dehydrin present in the dry seeds may assist the seed in withstanding stress due to lack of water in that state.

There are reports that seeds of higher plants accumulate large quantities of seed proteins including storage proteins and lectins (Adachi et al., 2003). Seed storage proteins are deposited in protein bodies, specialized vacuoles known as protein storage vacuoles (PSVs) (Shimada et al., 2003). They serve as nitrogen, sulphur and carbon source for the developing embryo during seed germination (Adachi et al., 2003). They are also major sources of dietary protein for humans. Shimada et al., (2003), demonstrated that unique vesicles which are precursor-accumulating vesicles, mediate the mass transport of storage proteins to PSVs in maturing pumpkin seeds. They also showed that isolated precursor-accumulating vesicles contain large amount of the precursors of major storage proteins of seeds. These types of precursor proteins are 8S globulin beta and alpha isoforms and 10kDa protein precursors. Vicilins also sequenced in this study, is a globulin storage protein found in plants. The importance of vicilin proteins was reported in soyabeans where 7S globulin subunit of soyabeans was observed to lower plasma lipids and upregulates liver (β)-VLDL receptors in rats fed with hypercholesteromic diet (Duranti et al., 2004).

Vicilins of various legume seeds show very high degree of sequence homology but present different properties when treated with proteases (Deshpande and Damodaran, 1989). We observed that the condition of the seed (dry or malted), determined the type of proteins identified. Swire-Clark and Marcotte (1999) had observed that late embryogenesis-abundant (LEA) proteins are capable of binding large amounts of water and it is possible that this is the function of Em protein found in the dry bambara seed. Chain I, crystal structure of the Bowman-Birk Inhibitor from *Vigna unguiculata* widespread in leguminous plants are involved in diverse cellular processes such as cell cycle, apoptosis, intracellular protein breakdown (Barbosa et al., 2007). The reorganization of cellular processes in plants has been linked to Actin cytoskeleton (Aon et al., 1999). Although the protscore and sequence coverage of Actin protein is low, it is important in maintaining the cytoskeleton. The presence of ARG10 protein which down regulates auxin showed that the dry bambara seed is in a dormant state and the cells are not actively dividing.

These proteins, PDF1 and Em may be important for the seeds to fulfil other biological processes apart from storing protein. Em proteins function in the embryo as storage protein for nutrients that are necessary in the early period of the germination process. It has been suggested that Em protein may be a cryptobiotic protein analogue found in *Artemia*. It therefore provides a form of desiccation tolerance to the cytoplasm while it is dehydrated at seed maturation (McCubbin et al., 1985). PDF1 plays a defensive role as shown in *Arabidopsis* plant.

The study of this legume is important because of its potential nutritional and health benefits to the populations of the developing world. Seed proteins have been implicated in anti-hypertensive activities (Yamada et al., 2008), regulation of food intake and body weight in humans (Anderson and Moore, 2004) and the improvement of plasma and liver lipid profile in rats fed with high cholesterol diet (Yang et al., 2007).

This study reports the different types of proteins of bambara groundnut (*Vigna subterranea*) present in the malted and dry seeds, their similarities and various uses which may enhance the appreciation of the food, nutritional and health values of the seed. Bambara groundnut belongs to the group of seeds labelled under-utilized and neglected, it is therefore strongly recommended for mass production to fill the current food gaps. Furthermore, extensive research is suggested to fully exploit the nutritional and health benefits derivable from it for the overall benefit of the starving populations of the third world.

REFERENCES

- Adachi M, Kanamori J, Masuda T, Yagasaki K, Kitamura K, Mikani B, Utsumi S. Crystal structure of soybean 11S globulin: glycin A3B4 homohexamer. *Proc Natl Acad Sci USA* 2003;100:7395-7400.
- Adu-Dapaah HK, Sangwan RS. Improving bambara groundnut productivity using gamma irradiation and *in vitro* techniques. *Afr J Biotechnol* 2004;3:260-5.
- Anderson GH, Moore SE. Dietary proteins in the regulation of food intake and body weight in humans. *J Nutr* 2004;134:974S-9S.
- Aon MA, Cortassa S, Gomez CDF, Iglesias AA. Effects of stress on cellular infrastructure and metabolic organization in plant cells. *Int Rev Cytol* 1999;194:239-73.
- Anchirinah VM, Yiridoe EK, Bennett-Lartey SO. Enhancing sustainable production and genetic resource conservation of Bambara groundnut: A survey of indigenous agricultural knowledge systems. *Outlook on Agric* 2001;30:281-8.

- Azam-Ali SN, Sesay A, Karikari KS, Mas-sawe FJ, Aguilar-Manjarrez J, Bannayan M, Hampson KJ. Assessing the potential of an underutilized crop - a case study using bambara groundnut. *Exp Agric* 2001;37:433-72.
- Barbosa JARG, Silva LP, Teles CL, Esteves GF, Azevedo FB, Ventura MM, de-Freitas SM. Crystal structure of the Bowman-Birk Inhibitor from *Vigna unguiculata* seeds in complex with β -trypsin at 1.55 Å resolution and its structural properties in association with proteinases. *Biophys J* 2007;92:1638-50.
- Brough SH, Azam-Ali SN, Taylor AJ. The potential of bambara groundnut (*Vigna subterranea*) in vegetable milk production and basic protein functionality systems. *Food Chem* 1993;47:277-83.
- Deshpande SS, Damodaran S. Structure-digestibility relationships of legume 7S storage proteins. *J Food Sci* 1989;54:108-13.
- Duranti M, Lovati MR, Dani V, Barbiroli A, Scarafoni A, Castiglioni S, Ponzzone C, Morazzoni P. The α' subunit from soybean 7S globulin lowers plasma lipids and upregulates liver β -VLDL receptors in rats fed a hypercholesterolemic diet. *J Nutr* 2004;134: 1334-9.
- Garrido-Ramirez ER, Sudarshana MR, Lucas WJ, Gilbertson RL. Bean dwarf mosaic virus BV1 protein is a determinant of the hypersensitive response and avirulence in *Phaseolus vulgaris*. *Mol Plant Microbe Interact* 2000;13:1184-94.
- McCubbin WD, Kay CM, Lane BG. Hydrodynamic and optical properties of the wheat germ Em protein. *Can J Biochem Cell Biol* 1985;63:803-11.
- Minka SR, Bruneteau M. Partial chemical composition of bambara pea (*Vigna subterranea* (L. Verde). *Food Chem* 2000; 68:273-6.
- Østergaard O, Melchior S, Roepstorff P, Svensson B. Initial proteome analysis of mature barley seeds and malt. *Proteomics* 2002; 2:733-9.
- Padulosi S, Hodgkin T, Williams JT, Haq N. Underutilized crops: trends, challenges and opportunities in the 21st century. In: Engels JMM et al. (eds): *Managing plant genetic diversity* (pp 323-38). CABI-IPGRI, UK and Rome, 2002.
- Poulter NH, Caygill JC. Vegetable milk processing and rehydration characteristics of bambara groundnut [*Voandzeia subterranea* (L.) thouars]. *J Sci Food Agric* 2006; 31:1158-63.
- Queitsch C, Hong S, Vierling E, Lindquist S. Heat shock protein 101 plays a crucial role in thermotolerance in arabidopsis. *Plant Cell* 2000;12:479-92.
- Shimada T, Fuji K, Tamura K, Kondo M, Nishimura M, Hara-Nishimura I. Vacuolar sorting receptor for seed storage proteins in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 2003;100:16095-100.
- Swire-Clark GA, Marcotte Jr. WR. The wheat LEA protein Em functions as an osmoprotective molecule in *Saccharomyces cerevisiae*. *Plant Mol Biol* 1999;39:117-28.
- Wang W, Vinocur B, Shoseyov O, Altmar A. Role of plant heat-shock protein and molecular chaperons in the abiotic stress response. *Trends Plant Sci* 2004;9:244-52.
- Wazael HN, Inga CB, Jorgen LC, Sven BA. Analysis of genetic diversity in bambara groundnut [*Vigna subterranea* (L.) Verdc] landraces using amplified fragment length polymorphism (AFLP) markers. *Afri J Biotech* 2004;3:220-5.

Yamada Y, Nishizawa K, Yokoo M, Hui Z, Onishi K, Teraishi M, Utsumi S, Ishimoto M, Yoshikawa M. Anti-hypertensive activity of genetically modified soybean seeds accumulating novokinin. *Peptides* 2008;29:331-7.

Yang S-C, Liu S-M, Yang H-Y, Lin Y-H, Chen J-R. Soybean protein hydrolysate improves plasma and liver lipid profiles in rats fed high-cholesterol diet. *J Amer Coll Nutr* 2007;26:416-23.

Zhou Y-C, Garrido-Ramirez ER, Sudarshana MR, Yendluri S, Gilbertson RL. The N-terminus of the Begomovirus nuclear shuttle protein (BV1) determines virulence or avirulence in *Phaseolus vulgaris*. *Mol Plant Microbe Interact* 2007;20:1523-34.