

# **Travelling wave solutions for gravity fingering in porous media flows**

Koondanibha Mitra, Ben Schweizer, Andreas Rätz

Preprint 2020-05

Dezember 2020

# Travelling wave solutions for gravity fingering in porous media flows

K. Mitra, A. Rätz, and B. Schweizer

November 10, 2020

**Abstract:** We study an imbibition problem for porous media. When a wetted layer is above a dry medium, gravity leads to the propagation of the water downwards into the medium. In experiments, the occurrence of fingers was observed, a phenomenon that can be described with models that include hysteresis. In the present paper we describe a single finger in a moving frame and set up a free boundary problem to describe the shape and the motion of one finger that propagates with a constant speed. We show the existence of solutions to the travelling wave problem and investigate the system numerically.

**MSC:** 76S05, 35C07, 47J40

**Keywords:** porous media, travelling waves, hysteresis

## 1 Introduction

Standard models for flow in unsaturated porous media fail in the description of a fundamental process, namely the imbibition into a dry medium with gravity as the driving force. While standard Richards models predict the formation of uniform imbibition fronts, the experimentally observed fingers [11, 26] can only be described with a model that incorporates hysteresis.

Models for incompressible unsaturated porous media flow typically use the water pressure  $p$  and the water saturation  $s$  as primary variables. The Darcy law for the velocity together with the mass balance equation leads to

$$\partial_t s = \nabla \cdot (k(s)[\nabla p + g e_z]), \quad (1.1a)$$

we refer to [2, 13, 22, 25] for the modelling. In the Richards equation (1.1a), the function  $k : [0, 1] \rightarrow \mathbb{R}$  is the permeability function which has to be determined from experiments,  $g$  is the gravitational acceleration,  $e_z$  is the normal vector pointing upwards. It is always assumed that  $s$  takes only values in  $[0, 1]$ .

Equation (1.1a) must be accompanied by a relation between saturation  $s$  and pressure  $p$ . Models without hysteresis demand either the algebraic relation  $p = p_c(s)$  for some given function  $p_c : [0, 1] \rightarrow \bar{\mathbb{R}}$ , or they include the “ $\tau$ -correction” and demand, for some physical parameter  $\tau > 0$ , known as the dynamic capillary number, that  $p = p_c(s) + \tau \partial_t s$ ; this latter model takes inertia in the material law into account,

see [12]. If, additionally, hysteresis in an imbibition process shall be modelled, a possible simple law is

$$\partial_t s = \frac{1}{\tau} [p - p_c(s)]_+, \quad (1.1b)$$

where  $[\cdot]_+ := \max\{0, \cdot\}$  denotes the positive part. Our aim is a travelling wave analysis of equation (1.1). We recall that  $p_c : (0, 1) \rightarrow \mathbb{R}$  is a given imbibition capillary pressure function and  $\tau > 0$  is a given constant.

Regarding the modelling we note that, if both imbibition and drainage should be modelled, one replaces (1.1b) by the model of [4],

$$\partial_t s = \frac{1}{\tau} [p - p_c(s)]_+ + \frac{1}{\tau} [p - p_d(s)]_-. \quad (1.2)$$

Here,  $p_d : (0, 1) \rightarrow \mathbb{R}$  is a drainage capillary pressure function with  $p_d(s) \leq p_c(s)$  for all  $s \in (0, 1)$ , and  $[\cdot]_- := \min\{0, \cdot\}$  is the negative part function. Equation (1.2) is a hysteresis model since, pointwise in space and time, all pressure values in the closed interval  $[p_d(s), p_c(s)]$  are permitted for a fixed saturation  $s$ . The play-type hysteresis model with dynamic capillary pressure was analyzed in [4, 15–17, 19, 21, 23, 24, 27]. Since we are interested in an infiltration problem with  $\partial_t s \geq 0$ , we restrict ourselves to the case  $p_d(s) = -\infty$  as in [9], i.e., we study (1.1b) instead of (1.2).




Figure 1: Motivation for this contribution. Left: A snapshot of a solution to the time dependent system (1.1). Fingers are clearly visible; the solution is comparable to experimental observations [21]. Middle: With another choice of boundary values and  $p_c$ , a single finger is generated. A small squared region of size  $2 \times 2$  around the finger-tip is marked [15]. Right: Enlargement of the marked region. We see the typical shape of the single finger in time-dependent calculations. The aim of this contribution is to analyze the travelling wave equations corresponding to (1.1) in order to obtain the shape of the single finger without a time-dependent calculation.

Numerical results for the time dependent system (1.1) are shown in Figure 1, originally published in [15, 21]. The figure illustrates a gravity driven imbibition process into an originally dry medium. Several fingers evolve in the process. It is observed that each finger travels approximately with constant speed. This has also been verified experimentally [26]. The present work aims at the description of a single finger in a co-moving frame of coordinates.

**Travelling wave ansatz, domains and boundary conditions.** Since we are interested in imbibition fronts in columns of porous media, we choose a cylindrical spatial domain  $\Omega_\infty$ . Restricting to two dimensions for convenience and denoting the width of the cylinder by  $L > 0$ , we consider  $\Omega_\infty := (0, L) \times \mathbb{R} \subset \mathbb{R}^2$ . Points in  $\mathbb{R}^2$  are denoted as  $x = (y, z)$ . We seek time-dependent solutions to (1.1) that move with

a constant speed  $c > 0$  in negative  $z$ -direction, i.e., downwards. This motivates the travelling wave coordinates

$$\tilde{z} = z + ct, \quad p(y, z, t) = p(y, \tilde{z}), \quad s(y, z, t) = s(y, \tilde{z}). \quad (1.3)$$

In the following, we omit the tilde symbol and write  $z$  instead of  $\tilde{z}$ . The new coordinates transform system (1.1) into

$$c\partial_z s = \nabla \cdot (k(s)[\nabla p + ge_z]), \quad (1.4a)$$

$$c\tau\partial_z s = [p - p_c(s)]_+. \quad (1.4b)$$

Even though the physical interpretation of a travelling wave solution requires the study of domains  $\Omega_\infty$  that extend to  $z \rightarrow \pm\infty$ , we choose here to study problem (1.4) on the semi-infinite domain

$$\Omega := (0, L) \times \mathbb{R}_+ \quad \text{with bottom} \quad \Sigma := (0, L) \times \{0\} = \{(y, 0) : 0 < y < L\}.$$

Truncations of the domain are necessary for numerical calculations and facilitate the analysis. The problem is translation invariant; one should consider the bottom  $\Sigma = \{z = 0\}$  as being far below the finger.

The boundary data are given by a prescribed saturation  $s_0 > 0$  and a prescribed pressure  $p_0$  at the bottom  $\Sigma$  of the domain, and by a prescribed total influx  $F_\infty$  on the top of the domain. More precisely, we assume that we are given  $s_0 : [0, L] \rightarrow [0, 1]$ ,  $p_0 : [0, L] \rightarrow \mathbb{R}$ , and  $F_\infty \in \mathbb{R}_+ = (0, \infty)$ , and impose the boundary conditions

$$\int_0^L k(s(y, z))[\partial_z p(y, z) + g] dy \rightarrow F_\infty \quad \text{as } z \rightarrow +\infty, \quad (1.5a)$$

$$s = s_0 \quad \text{at } z = 0, \quad (1.5b)$$

$$p = p_0 \quad \text{at } z = 0. \quad (1.5c)$$

If the initial saturation of the medium is given by a number  $s_* \in (0, 1)$ , a natural choice for the boundary data is  $s_0 \equiv s_*$  and  $p_0 \equiv p_c(s_*)$ . Along the lateral boundaries of  $\Omega$  we impose homogeneous Neumann conditions (no flux).

**Main results.** We perform an analysis of the travelling wave problem (1.4)–(1.5) on  $\Omega$ . For the most part of this article, we prescribe the relaxation parameter  $\tau$ , the frame speed  $c$ , and the boundary data  $s_0$ ,  $p_0$ , and  $F_\infty$ . Only in our last result, Theorem 4.7, we choose  $c$  in dependence of the other parameters in order to satisfy a physically adequate flux condition on the lower boundary.

The first part of our results concerns the system (1.4)–(1.5) on the bounded truncated domain  $\Omega^H = (0, L) \times (0, H)$ . We choose boundary conditions on the upper boundary appropriately and show that the system has a solution. The solution can be found with a variational principle, the analysis is given in Section 3.

The numerical part of this paper deals with this truncated problem. One result is the calculation of a finger solution, see Figure 2. The numerical method and the results are described in Section 5.

The limit  $H \rightarrow \infty$  for the solutions on the bounded domain is studied in Section 4. We find that every sequence of solutions  $(s_H, p_H)$  to truncated domain problems possesses a subsequence and a limit  $(s, p)$  which is a solution of the original problem

(1.4). The limit process shows an interesting dichotomy: In one case, the flux boundary condition for  $z \rightarrow \infty$  as in (1.5) remains satisfied (“large solution”). In the other case (“small solution”), only a corresponding inequality is satisfied.

The two cases are analyzed further. We find that “large solutions” are of the type that we would like to see in the fingering process: they possess a free boundary, the pressure  $p$  tends to  $-\infty$  as  $z \rightarrow \infty$ , and the solution is “large” in the sense that the saturation exceed a certain threshold. In the second case, the properties are reverted: The solution has a bounded pressure and it is “small” in the same sense as the solution was “large” in the other case. Interestingly, both types of solutions are found numerically, see Section 5.

**Free boundary problem.** Let us emphasize that we treat a free boundary problem. By (1.4b), one has to distinguish between the subdomain  $\{x \in \Omega \mid \partial_z s(x) > 0\}$  (expected to be in the bottom) and the subdomain  $\{x \in \Omega \mid \partial_z s(x) = 0\}$  (expected in the top part). In physical terms, this means that an imbibition process occurs near and below the finger-tip, whereas, in the region around the developed finger, the saturation does not change any more. With reference to the hysteresis relation, we note that the  $z$ -independent saturation implies that the pressure can take arbitrary values (below  $\min p_c(s)$ ). Therefore, the pressure profile does not have to reflect the saturation profile and the fingers can remain stable in their upper part; no blurring by pressure differences occurs.




Figure 2: A numerical solution of the free boundary travelling wave problem. The gray scale indicates the values of the saturation  $s$  (left) and the pressure  $p$  (right). The level line  $\Gamma = \{x \mid p = p_c(s)\}$  is marked in the left image. The line  $\Gamma$  shows the free boundary: Below the line, the saturation is increasing, above the line, the saturation remains constant (increasing in vertical direction and, hence, increasing in time when interpreted as a time dependent solution).

With Theorem 4.7 we provide the result that, for every  $F_\infty$  within appropriate bounds, there exists a wave speed  $c$  such that a physical flux condition at the lower boundary is satisfied.

**Literature.** The classical porous media equation is obtained by setting  $\tau = 0$  and by replacing (1.1b) by the algebraic law  $p = p_c(s)$ . This classical equation is interesting when the permeability coefficient is degenerate  $k(0) = 0$ . For existence and uniqueness results in this classical case we refer to [1, 20]. The hysteresis model

(1.1b) was introduced in [3, 4, 12]. It combines dynamic effects ( $\tau > 0$ ) with a play-type hysteresis relation; the latter allows for an interval of pressure values  $p$  for a fixed saturation  $s$ . For a review of the modelling, we refer to [25].

For the model (1.1), well-posedness results have been obtained in one space dimension in [4], and in higher dimension in [15, 21]. Existence of solutions for an extension of the play-type model was shown in [16]. In [23], it was shown that the model does not define an  $L^1$ -contraction; in this sense, it can explain the fingering effect. The fingers were found numerically for unsaturated media in [15], for the two-phase flow in [14]. Fingers were also observed numerically in [5, 7], where a free-energy based approach is used for modelling the capillary pressure. For a result with a degenerate  $p_c$ -curve, see [24]. A uniqueness result was derived in [6].

Travelling waves for the model have been analyzed in [17, 19, 27]. An analysis for pure imbibition ( $\partial_t s \geq 0$  allows to set  $p_d(s) = -\infty$ ) was previously performed for one space dimension in [9]. The present work extends the results to two space dimensions. Let us note that the methods are independent of the dimension and that, up to notation, the results remain valid, e.g., in three space dimensions. The dimension enters only in Sobolev embeddings that are used for regularity statements in the appendix.

## 2 Preliminaries

The coefficient functions  $k$  and  $p_c$  are fixed throughout this work. We make assumptions that are quite common and consistent with experiments, see [13]. For an illustration see Figure 3.

**Assumption 2.1.** *The functions  $k : [0, 1] \rightarrow [0, \infty)$  and  $p_c : (0, 1) \rightarrow \mathbb{R}$  satisfy:*

**(Ass-pc)** *The function  $p_c$  is differentiable and for some  $\rho > 0$  holds  $p'_c \geq \rho$  on  $(0, 1)$ . Upon normalization of the pressure, we can set  $p_c(s_*) = 0$  for a given saturation value  $s_* \in \mathbb{R}$ . We assume  $p_c(s) \rightarrow -\infty$  as  $s \searrow 0$  and  $p_c(s) \rightarrow \infty$  as  $s \nearrow 1$ .*

**(Ass-k)** *The function  $k$  is differentiable,  $k|_{(0,1)} \in C^2$ , and  $k'(\cdot), k''(\cdot) > 0$  on  $(0, 1)$ .*




Figure 3: Typical functions  $p_c$  and  $k$ .

**The free boundary description.** What qualitative behavior can we expect for solutions of the travelling wave problem (1.4)–(1.5)? We expect that the pressure stabilizes, as  $z \rightarrow +\infty$ , to an affine function with  $\nabla p \approx -g_F e_z$ . If  $s$  (and hence  $k(s)$ ) does not depend on  $z$ , then both sides of (1.4a) can vanish. This is what we expect for solutions in the upper part of the domain. We will be interested in solutions  $(p, s)$  that satisfy, for some  $h \in \mathbb{R}_+$ ,

$$\partial_z s = 0 \text{ and } p \leq p_c(s) \text{ for all } (y, z) \text{ with } y \in (0, L) \text{ and } z > h. \quad (2.1)$$

For such a solution we can define a function  $\Psi : [0, L] \rightarrow [0, \infty)$  as

$$\Psi(y) := \inf \{z_0 > 0 \mid \partial_z s(y, z) = 0 \text{ for all } z \geq z_0\}. \quad (2.2)$$

The graph of  $\Psi$  is a part of the free-boundary,  $\{(y, \Psi(y)) \mid y \in (0, L)\} \subset \Gamma$ . For the rest of the paper, we define the function  $s^* : [0, L] \rightarrow [0, 1]$  as

$$s^*(y) := \lim_{z \rightarrow \infty} s(y, z). \quad (2.3)$$

By positivity  $\partial_z s(y, z) \geq 0$  and boundedness of  $s$ , the function  $s^*$  is well-defined for solutions  $(s, p)$  of (1.4). When a solution satisfies (2.1), there holds  $s(y, z) = s^*(y)$  for all  $z > h$ .




Figure 4: When interpreted as a solution of the time-dependent problem, the finger moves with a constant speed downwards. The dashed line represents the boundary of the finger; one may think of an isoline of the saturation. The graph at the top part of the Figure indicates a profile of the limiting saturation  $s^*$  as defined in (2.3).

We refer to Figure 4 for an illustration. It is important not to confuse the free boundary  $\Gamma$  with the shape of the finger (the region of high saturation). We emphasize that the saturation profile remains unchanged (independent of  $z$ ) above  $\Gamma$ ; in particular, the finger extends to  $z \rightarrow +\infty$ .

**Relations in the travelling wave formulation.** A fundamental problem in travelling wave analysis is the determination of free parameters, in our case the wave speed  $c$ . The other parameters are fixed:  $\tau, g > 0$  are physical constants,  $L > 0$  a geometrical constant, and the boundary conditions fix  $F_\infty > 0$  and  $s_* > 0$ . In the

travelling wave formulation,  $c \geq 0$  is a further unknown of the system. Nevertheless, for the most part of our analysis, we fix boundary values  $s_0$  and  $p_0$  and treat the problem with prescribed  $c$ . Only in our final result we determine  $c$  from an additional boundary condition for  $z \rightarrow -\infty$ .

Let us collect some properties of the real parameters.

**Lemma 2.2** (Wave speed and limiting pressure in the doubly infinite domain). *Let  $(s, p) \in C^1(\Omega_\infty) \times C^2(\Omega_\infty)$  be a classical solution to (1.4) on  $\Omega_\infty$  with the boundary condition (1.5a) and the two conditions  $s \rightarrow s_*$  and  $k(s)\nabla p \rightarrow 0$  as  $z \rightarrow -\infty$ . Then, with  $s^*$  as in (2.3), the wave speed satisfies*

$$c = (F_\infty - k(s_*)gL) / \left( \int_0^L (s^*(y) - s_*) dy \right). \quad (2.4)$$

If the solution possesses a free boundary, i.e. (2.1) holds for some  $h > 0$ , then

$$g_F := g - \left( F_\infty / \int_0^L k(s^*(y)) dy \right) \quad (2.5)$$

satisfies  $g_F > 0$  and there holds  $\nabla p(y, z) + g_F e_z \rightarrow 0$  as  $z \rightarrow \infty$  for every  $y \in (0, L)$ .

*Proof.* Integrating (1.4a) over  $(0, L) \times (-H, H)$  yields

$$c \int_0^L s(y, z) dy \Big|_{z=-H}^H = \int_0^L k(s(y, z)) [\partial_z p(y, z) + g] dy \Big|_{z=-H}^H.$$

Sending  $H \rightarrow \infty$  provides (2.4).

Relation (2.1) implies that  $s(y, z) = s^*(y)$  holds for  $z > h$ . Therefore, the elliptic equation reduces to

$$\nabla \cdot (k(s^*)\nabla p) = 0 \text{ in } (0, L) \times (h, \infty). \quad (2.6)$$

In particular, the flux quantity  $\int_0^L k(s^*) \partial_z p(y, z) dy$  is independent of  $z$  for  $z > h$ . The boundary condition (1.5a) allows to evaluate this flux for  $z \rightarrow \infty$ ; we find

$$\int_0^L k(s^*(y)) \partial_z p(y, z) dy = F_\infty - g \int_0^L k(s^*(y)) dy = -g_F \int_0^L k(s^*(y)) dy. \quad (2.7)$$

This provides that, for  $z > h$ , the weighted average of  $\partial_z p$  coincides with  $-g_F$ .

Solutions  $p$  of the elliptic equation (2.6) with homogeneous Neumann boundary conditions on unbounded domains have the property that  $\nabla p$  stabilizes to a constant as  $z \rightarrow \infty$  (a consequence of the strong maximum principle for  $\partial_z p$ ). Relation (2.7) shows that this constant is  $-g_F e_z$ .

Let us assume for a contradiction  $g_F < 0$ . Then  $p$  is a growing function for  $z \rightarrow \infty$ . This is in contradiction with (1.4b), in which the left hand side vanishes for  $z > h$  and  $p_c(s)$  is independent of  $z$  for  $z > h$ .

Let us now assume  $g_F = 0$  in order to exclude also this case. We use a maximum principle for  $p$  in the interior of the set  $\{(y, z) | \partial_z s = 0\} = \{(y, z) | p \leq p_c(s)\}$ . The minimum of  $p$  is attained at the boundary. At the lower boundary of this set, there holds  $p = p_c(s)$ . This implies that the minimum is attained in a point of the form  $(y, z) = (y, \Psi(y))$ . We now use, for any  $\varepsilon > 0$ , the strong maximum principle:  $p(y, \Psi(y) + \varepsilon) > p(y, \Psi(y)) = p_c(s(y, \Psi(y))) = p_c(s(y, \Psi(y) + \varepsilon))$ . This implies  $p > p_c(s)$  in  $(y, \Psi(y) + \varepsilon)$  and hence  $\partial_z s(y, \Psi(y) + \varepsilon) > 0$ , in contradiction to the construction of  $\Psi$ .  $\square$

**Notation.** Together with the domain  $\Omega = (0, L) \times \mathbb{R}_+$  with bottom boundary  $\Sigma = (0, L) \times \{0\}$  we also use, for any  $H > 0$ , the bounded domain  $\Omega^H := (0, L) \times (0, H)$  with the top boundary  $\Sigma^H := (0, L) \times \{H\}$ . We recall that we always impose homogeneous Neumann conditions at the lateral boundaries  $\{0\} \times \mathbb{R}_+$  and  $\{L\} \times \mathbb{R}_+$  (accordingly for the truncated domain).

The function  $\text{sign} : \mathbb{R} \rightarrow \{0, 1\}$  is defined as  $\text{sign}(u) := 0$  for  $u \leq 0$ , and  $\text{sign}(u) := 1$  otherwise. The letter  $C$  denotes a generic positive constant and the value may change from one line to the next in calculations. We already introduced  $[q]_+ = \max\{0, q\} = (q + |q|)/2$  and  $[q]_- = \min\{0, q\} = -[-q]_+$ .

### 3 Existence result for bounded domains

Let  $\tau > 0$ ,  $s_* \in (0, 1)$ , and two functions  $p_0 \in H^{\frac{1}{2}}(\Sigma) \cap C^0(\bar{\Sigma})$  and  $s_0 \in H^1(\Sigma)$  be given. We assume  $s_* \leq s_0 \leq 1$  and  $p_0 \geq p_c(s_0)$ . For a height parameter  $H > 0$  we introduce the following truncated problem.

**Definition 3.1** (Truncated domain travelling wave problem). *Let  $c, F_\infty > 0$  be given. A pair  $(s, p) \in H^1(\Omega^H) \times H^2(\Omega^H)$  on the domain  $\Omega^H = (0, L) \times (0, H)$  with upper boundary  $\Sigma^H$  and lower boundary  $\Sigma$  is a truncated domain travelling wave solution (TW<sub>H</sub>-solution) if there holds*

$$c\partial_z s = \nabla \cdot (k(s)[\nabla p + ge_z]) \quad \text{in } \Omega^H, \quad (3.1a)$$

$$c\tau\partial_z s = [p - p_c(s)]_+ \quad \text{in } \Omega^H, \quad (3.1b)$$

$$s = s_0, \quad p = p_0 \quad \text{on } \Sigma, \quad (3.1c)$$

$$p \equiv p^* \in \mathbb{R} \quad \text{on } \Sigma^H, \quad (3.1d)$$

$$\int_{\Sigma^H} k(s)[\partial_z p + g] = F_\infty. \quad (3.1e)$$

We emphasize that the constant pressure value  $p^* \in \mathbb{R}$  is a free parameter and part of the solution of the problem.

We note that for every TW<sub>H</sub>-solution  $(s, p)$ , the flux quantity

$$F_c(z) := \int_0^L k(s(y, z))[\partial_z p(y, z) + g] - cs(y, z) \, dy \quad (3.2)$$

is independent of  $z \in (0, H)$  by (3.1a). Evaluating this flux in the upper and in the lower boundary provides, by (3.1e),

$$\int_{\Sigma} k(s_0)\partial_z p + \int_{\Sigma} (k(s_0)g - cs_0) = F_\infty - \int_{\Sigma^H} cs. \quad (3.3)$$

**Remark 3.2.** *Let us give a sloppy description of the consequences of (3.3) for small boundary data  $s_0$ . There is the possibility that  $\partial_z p$  is large at  $\Sigma$ . This means that a sharp transition occurs near the lower boundary. In the opposite case (without boundary layer), the left hand side of (3.3) is small. In this case, a moderate flux  $F_\infty > 0$  forces the system that  $s$  is not small at  $\Sigma^H$ . This is the desired behavior for finger-like travelling wave solutions; they should connect a small saturation at  $z = 0$  with a moderate or large saturation at  $z = H$ .*

**Remark 3.3** (A condition for the wave speed  $c$ ). *Let us highlight another consequence of the fact that  $F_c$  of (3.2) is independent of  $z$ . When  $(s, p)$  is a solution on the doubly unbounded domain  $\Omega_\infty$  then we expect, in the limit  $z \rightarrow -\infty$ , that  $s \rightarrow s_*$ ,  $p \rightarrow p_c(s_*)$ , and  $\partial_z p \rightarrow 0$ . In this situation, the constant flux quantity is necessarily  $F_c = (gk(s_*) - cs_*)L$ .*

*We use this observation in order to choose a closure condition for the case when the speed  $c$  is treated as an unknown: Even when we solve a Dirichlet problem in the truncated domain  $\Omega$  with boundary conditions  $s_0$  and  $p_0$  at the lower boundary  $\Sigma$ , we will seek for  $c$  and solutions to the Dirichlet problem that satisfy the additional relation*

$$F_c = \int_{\Sigma} (k(s_0)[\partial_z p + g] - cs_0) = (gk(s_*) - cs_*)L. \quad (3.4)$$

*Theorem 4.7 yields that, given  $s_0$ ,  $p_0$ ,  $s_*$ , and  $F_\infty$ , we find a speed  $c$  such that (3.4) is satisfied.*

In the remainder of this section, we seek for  $TW_H$ -solutions  $(s, p)$ . We use the space of functions

$$H_{\#}^1(\Omega^H) := \{u \in W^{1,2}(\Omega^H) \mid \text{tr}(u) = 0 \text{ on } \Sigma, \exists u^* \in \mathbb{R} : u = u^* \text{ on } \Sigma^H\}. \quad (3.5)$$

The weak formulation of (3.1a) and (3.1e) is:

$$\int_{\Omega^H} c \partial_z s \phi + \int_{\Omega^H} k(s)[\nabla p + g e_z] \cdot \nabla \phi = \int_{\Sigma^H} F_\infty \phi \quad \text{for all } \phi \in H_{\#}^1(\Omega^H). \quad (3.6)$$

**Theorem 3.4** (Existence of  $TW_H$ -solutions to prescribed data). *Let  $H, c, \tau, F_\infty > 0$  and  $s_* \in (0, 1)$  be given, let  $p_0 \in H^{\frac{1}{2}}(\Sigma) \cap C^0(\bar{\Sigma})$  and  $s_0 \in H^1(\Sigma)$  satisfy*

$$s_* \leq s_0 < 1, \quad \text{and} \quad 0 < p_0 - p_c(s_0) \text{ on } \Sigma.$$

*Then there exists a  $TW_H$ -solution  $(s, p)$  with  $s, \partial_z s \in L^2(\Omega^H)$ ,  $p \in H^1(\Omega^H) \cap H_{\text{loc}}^2(\Omega^H)$ .*

*Proof.* We use an iteration over saturation fields.

*Definition of the iteration.* Let there be given a saturation field

$$s^{i-1} \in Y := \{s \in L^2(\Omega^H) \mid s_* \leq s \leq 1\}.$$

We define the coefficient functions  $a := k(s^{i-1})$  and  $b := p_c(s^{i-1})$  on  $\Omega^H$ . We seek a solution  $p$  of

$$\frac{1}{\tau} [p - b]_+ = \nabla \cdot (a [\nabla p + g e_z]) \text{ in } \Omega^H, \quad (3.7)$$

with the boundary conditions  $p = p_0$  on  $\Sigma$  and (3.1d)–(3.1e). This solution can be found with a variational method. We define the space of admissible functions as  $X_{p_0} := \{u \in H^1(\Omega^H) \mid u = p_0 \text{ on } \Sigma, \exists u^* \in \mathbb{R} : u = u^* \text{ on } \Sigma^H\}$  and minimize the functional

$$A : X_{p_0} \rightarrow \mathbb{R}, \quad A(p) := \int_{\Omega^H} \frac{1}{2\tau} [p - b]_+^2 + \frac{1}{2} a |\nabla p + g e_z|^2 - F_\infty \int_{\Sigma^H} p. \quad (3.8)$$

The functional is convex and coercive, which implies that a minimizer  $p$  exists. The Euler-Lagrange equation for  $p$  reads

$$\int_{\Omega^H} \frac{1}{\tau} [p - b]_+ \varphi + a [\nabla p + g e_z] \cdot \nabla \varphi = F_\infty \int_{\Sigma^H} \varphi \quad \forall \varphi \in H_{\#}^1(\Omega^H).$$

Since arbitrary compactly supported test-functions  $\varphi$  can be inserted, equation (3.7) holds for  $p$ . The Euler-Lagrange equation additionally encodes the boundary condition  $\int_{\Sigma^H} a(\partial_z p + g) = F_\infty$ . Given  $p^i = p$ , we can solve the family of ordinary differential equations

$$c\tau\partial_z s = [p^i - p_c(s)]_+, \quad (3.9)$$

with initial data  $s = s_0$  on  $z = 0$ ; this system is related to (3.1b) together with the first equation in (3.1c). We denote the solution of this system by  $s =: s^i$ .

*Fixed point of the iteration.* We claim that, for some constant  $C = C(H, c, \tau)$  independent of  $s^{i-1}$ , the pressure  $p = p^i$  satisfies

$$\|p\|_{L^2(\Omega^H)}^2 + \|\nabla p\|_{L^2(\Omega^H)}^2 \leq C. \quad (3.10)$$

In order to show this estimate, we first choose an  $H^1$ -extension  $\hat{p}_0$  of the data  $p_0$ , vanishing at the upper boundary. We can now multiply equation (3.7) with  $p - \hat{p}_0$  and integrate to obtain

$$\int_{\Omega^H} \frac{1}{\tau} [p - b]_+ ([p - b] - \hat{p}_0 + b) + \int_{\Omega^H} (a[\nabla p + g e_z]) \cdot \nabla(p - \hat{p}_0) = \int_{\Sigma^H} F_\infty p.$$

One of the integrals on the left hand side is an upper bound for  $k(s_*)\|\nabla p\|_{L^2(\Omega^H)}^2$ , the other term with quadratic growth in  $p$  is on the left hand side and positive because of  $[p - b]_+[p - b] \geq 0$ . The remaining terms have linear growth in  $p$  and can therefore be estimated with Youngs inequality and with the Poincaré inequality.

The corresponding solutions  $s^i = s$  of the ordinary differential equation satisfy  $0 \leq s \leq 1$  by the growth assumption on  $p_c$ . In particular, there holds  $s^i \in Y$ . With  $R := |\Omega^H|^{1/2} = |LH|^{1/2}$ , we find that the above construction provides a map

$$\mathcal{T} : Y \supset B_R(0) \rightarrow B_R(0) \subset Y, \quad s^{i-1} \mapsto s^i.$$

We claim that the map  $\mathcal{T}$  is compact. We will show the compactness below with the characterization of compact subsets of  $L^2(\Omega^H)$  by Kolmogorov-Riesz. An application of Schauder's fixed point theorem yields the existence of the desired solution  $s$ .

Let us turn to compactness of  $\mathcal{T}$ . We consider the family  $p = p^i$  of solutions for  $s = s^{i-1} \in B_R(0)$ . This family of solutions is bounded in  $H^1(\Omega^H)$ , hence the finite differences  $p(y, \cdot) - p(y + \delta, \cdot) \in L^2((0, H); \mathbb{R})$  are small for  $\delta > 0$  small, independent of  $s$ . More precisely,

$$\int_0^{L-\delta} \int_0^H |p(y, z) - p(y + \delta, z)|^2 dz dy \leq \eta(\delta),$$

with  $\eta(\delta) \rightarrow 0$  as  $\delta \rightarrow 0$ , independent of  $s$ . We now consider two solutions of the ordinary differential equation (3.9),  $s(y, \cdot)$  and  $s(y + h, \cdot)$  to inputs  $p(y, \cdot)$  and  $p(y + h, \cdot)$ . The solutions differ only as much as their right hand sides and their initial values differ. Because of our assumption  $s_0 \in H^1(\Sigma)$ , we therefore find also for the solutions

$$\int_0^{L-\delta} \int_0^H |s(y, z) - s(y + \delta, z)|^2 dz dy \leq C\eta(\delta).$$

On the other hand, since  $\partial_z s$  is bounded in  $L^2(\Omega^H)$ , the corresponding estimate  $\int_0^L \int_0^{H-\delta} |s(y, z + \delta) - s(y, z)|^2 dz dy \leq C\eta(\delta)$  is clear. This shows compactness of the image set of  $s$ -fields.  $\square$

## 4 Unbounded domain solutions for $H \rightarrow \infty$

In this section we analyze the solutions  $(s_H, p_H)$  in the limit  $H \rightarrow \infty$ . Again, for the larger part of this section, we keep  $\tau > 0$ ,  $s_* \in (0, 1)$ ,  $F_\infty$ , and  $c > 0$  fixed; only in Theorem 4.7 we determine  $c$  from the other parameters. The main result of this section is the following: Let  $(s_H, p_H)$  denote the  $TW_H$ -solution as discussed in Theorem 3.4. Then, for  $H \rightarrow \infty$ , there holds  $(s_H, p_H) \rightarrow (s, p)$  in an appropriate sense for some limit pair  $(s, p)$ , which is defined on the unbounded domain  $\Omega$ . The pair  $(s, p)$  is a travelling wave solution for the semi-infinite domain  $\Omega$ .

It turns out that two different limiting solution types are possible. Type I is the “large solution”. It is characterized by the following properties: 1) The solution is large in the sense that  $\int_0^L gk(s(y, z_0)) dy \geq F_\infty$  for some  $z_0$ . This means that a certain  $F_\infty$ -dependent threshold is exceeded by the saturation variable. 2) The solution has a free boundary: For some  $h > 0$  there holds  $\partial_z s(y, z) = 0$  for every  $z \geq h$ . 3) The solution has an unbounded pressure,  $p \rightarrow -\infty$  as  $z \rightarrow \infty$ .

Accordingly, Type II solutions are the “small solutions”. They have a bounded pressure and no free boundary.

To proceed with the analysis, we consider different assumptions.

**Assumption 4.1.** *The following properties can be considered for the solution sequence  $(s_H, p_H)$  of (3.1), obtained in Theorem 3.4.*

**Bounds for parameters** *The limiting saturation  $s_* \in (0, 1)$ , the wave speed  $c$ , and the flux  $F_\infty$  satisfy*

$$gk'(s_*) < c < g(k(1) - k(s_*))/(1 - s_*), \quad (4.1a)$$

$$gL[k(s_*) + k'(s_*)(1 - s_*)] < F_\infty < gLk(1). \quad (4.1b)$$

**Bound for the pressure** *For a real number  $\bar{p} < \infty$  independent of  $H$  holds*

$$p_H \leq \bar{p} \quad \text{in } \Omega^H. \quad (4.2)$$

**Local bound for the gradient** *There exists  $C_P > 0$  such that, for every  $H > 0$ ,*

$$\|\nabla p_H\|_{L^\infty(\Omega^H)} \leq C_P. \quad (4.3)$$

**Regularity** *The saturation has the regularity properties*

$$s_H, \partial_z s_H \in H^1(\Omega^H). \quad (4.4)$$

The assumptions have a quite different character. Inequalities (4.1) are ranges for the physical parameters; we expect the existence of travelling waves in this parameter regime. The uniform upper bound of (4.2) is expected to hold, but it should be derived from the system of equations, which we did not succeed to do. The regularity estimate (4.3) and the local regularity (4.4) can be shown with the tools of elliptic regularity theory, see [10]. We formulate them here as assumptions, since the regularity theory is not the focus of this contribution.

We note that the relations (4.2)–(4.3) imply three further estimates:

$$\|s_H\|_{L^\infty(\Omega^H)} \leq \bar{s} := p_c^{-1}(\bar{p}) < 1. \quad (4.5a)$$

In Lemma A.3 we prove that, for a constant  $C_s = C_s(C_P, s_0, p_0)$ ,

$$\|\nabla s_H\|_{L^\infty(\Omega^H)} \leq C_s. \quad (4.5b)$$

Since (4.5b) provides  $\|\partial_z s_H\|_{L^\infty(\Omega^H)} < C_s$ , one also has from (3.1b) that

$$p_H \leq p_c(s_H) + c\tau C_s \quad \text{in } \Omega^H. \quad (4.5c)$$

Our main result on unbounded domains is the following.

**Theorem 4.2** (Limits of  $TW_H$ -solutions). *Let  $c, F_\infty, \tau > 0$ ,  $s_* \in (0, 1)$ , and boundary data  $s_0, p_0 \in C^1(\Sigma)$  with  $s_* \leq s_0 < 1$  and  $p_c(s_0) < p_0$  be given. Let all the properties of Assumption 4.1 be satisfied. For a sequence  $H \rightarrow \infty$ , let  $(s_H, p_H)$  be solutions to (3.1). Then, for a limiting pair  $(s, p)$ , there holds  $(s_H, p_H) \rightarrow (s, p)$  locally in  $L^2(\Omega)$ . The limits satisfy  $s \in C_b^0(\Omega)$ ,  $\partial_z s \in L^2(\Omega)$ ,  $p \in H_{\text{loc}}^2(\Omega) \cap H_{\text{loc}}^1(\Omega \cup \Sigma)$ ,  $(s, p) = (s_0, p_0)$  on  $\Sigma$ , and (1.4). The solution  $(s, p)$  is either of Type I or of Type II:*

**Type I: “Large solution”** *The solution has a free-boundary: There exists  $h \in \mathbb{R}_+$  such that  $\partial_z s = 0$  for all  $y \in (0, L)$  and  $z \geq h$ . The solution is large in the sense that, with  $s^*(y) := \lim_{z \rightarrow \infty} s(y, z)$ , there holds  $g \int_0^L k(s^*(y)) dy \geq F_\infty$ , with strict inequality if  $p, s$ , and  $\partial_z s$  are continuous. Furthermore,  $p(y, z) \rightarrow -\infty$  as  $z \rightarrow \infty$  in this case.*

**Type II: “Small solution”** *The solution has a bounded pressure, there holds  $p \in L^\infty(\Omega)$ . Furthermore,  $\nabla p \in L^2(\Omega)$ . The solution is “small” in the sense that  $g \int_0^L k(s^*(y)) dy \leq F_\infty$ .*

Type I solutions satisfy additionally the boundary condition (1.5a).

The theorem follows from Propositions 4.4 and 4.5. Before we can prove these results, we have to establish an a priori estimate, which is the basis for both propositions.

**Lemma 4.3** (A priori estimate for  $TW_H$ -solutions). *Let  $F_\infty, c, \tau, s_* > 0$  and  $s_0, p_0 \in C^1(\Sigma)$  with  $s_* \leq s_0(y) < 1$  and  $p_c(s_0) < p_0$ . For a sequence  $0 < H \rightarrow \infty$ , let  $(s_H, p_H)$  be solutions to (3.1). We assume that the solution sequence satisfies relations (4.3) and (4.4). We use the characteristic functions  $\mathbb{1}_> := \mathbb{1}_{\{\partial_z s_H > 0\}}$  and  $\mathbb{1}_0 := \mathbb{1}_{\{\partial_z s_H = 0\}}$  on  $\Omega^H$ . There exists a constant  $C_1 := C_1(c, \tau, s_0, p_0, C_P)$ , independent of  $H$ , such that*

$$\int_{\Omega^H} \mathbb{1}_> p_c'(s_H) |\nabla s_H|^2 + \int_{\Omega^H} \mathbb{1}_0 \frac{1}{p_c'(p_c^{-1}(p_H))} |\nabla p_H|^2 + c\tau \int_{\Sigma^H} |\nabla s_H|^2 \leq C_1. \quad (4.6a)$$

If, additionally, (4.2) is satisfied, there exists  $C_2 := C_2(c, \tau, s_0, p_0, C_P, \bar{p})$  such that

$$\int_{\Omega^H} \mathbb{1}_> |\nabla p_H|^2 + \int_{\Omega^H} |\nabla(\partial_z s_H)|^2 \leq C_2. \quad (4.6b)$$

*Proof.* Within this proof, we write  $(s, p)$  instead of  $(s_H, p_H)$  to have shorter formulas. With  $C > 0$  we refer to generic constants that may depend on  $c, \tau, s_0, p_0, C_P, \bar{p}$ , but not on  $H$ .

*Step 1: Test function  $K(s)$ .* We use  $K : [0, 1] \rightarrow [0, \infty)$ , defined as  $K(s) := \int_0^s k(\varrho)^{-1} d\varrho$ . Equivalently, we may say that  $K$  is the primitive of  $k^{-1}$ , satisfying

$$K'(s) = \frac{1}{k(s)}, \quad K(0) = 0.$$

Below, we will use additionally the primitive of  $K$ ; we denote by  $\tilde{K}$  the function that satisfies  $\tilde{K}'(s) = K(s)$  and  $\tilde{K}(0) = 0$ .

We use  $K(s)(y, z) = K(s(y, z))$  as a test function in (3.1a) and study

$$c \int_{\Omega^H} K(s) \partial_z s = \int_{\Omega^H} K(s) \nabla \cdot (k(s) [\nabla p + g e_z]).$$

Using an integration by parts, we may write this relation as

$$\begin{aligned} c \int_{\Omega^H} \partial_z \tilde{K}(s) + \int_{\Omega^H} k(s) [\nabla p + g e_z] \cdot \nabla K(s) \\ = \int_{\Sigma^H} K(s) k(s) [\partial_z p + g] - \int_{\Sigma} K(s_0) k(s_0) [\partial_z p + g]. \end{aligned} \quad (4.7)$$

We have constructed  $K$  such that  $\nabla K(s) = k(s)^{-1} \nabla s$ . This gives a simple formula for the second integral. With another integration by parts and with  $\mathbb{1} = \mathbb{1}_> + \mathbb{1}_0$  we find

$$\begin{aligned} c \int_{\Sigma^H} \tilde{K}(s) - c \int_{\Sigma} \tilde{K}(s_0) + \int_{\Omega^H} \mathbb{1}_> \nabla p \cdot \nabla s + \int_{\Omega^H} \mathbb{1}_0 \partial_y p \partial_y s + \int_{\Omega^H} g e_z \cdot \nabla s \\ = \int_{\Sigma^H} K(s) k(s) \partial_z p - \int_{\Sigma} K(s_0) k(s_0) \partial_z p + g \int_{\Sigma^H} K(s) k(s) - g \int_{\Sigma} K(s_0) k(s_0). \end{aligned} \quad (4.8)$$

We note that the last two integrals on the right hand side and the first two integrals on the left hand side are bounded. Since we assumed (4.3), actually the entire right hand side of (4.8) is bounded. The last integral of the left hand side can be integrated, which shows that also this term is bounded. We therefore find

$$\int_{\Omega^H} \mathbb{1}_> \nabla p \cdot \nabla s + \int_{\Omega^H} \mathbb{1}_0 \partial_y p \partial_y s \leq C. \quad (4.9)$$

We want to rewrite the first integral. With this aim, we observe that  $c\tau \partial_z s = [p - p_c(s)]_+$  in  $\Omega^H$  implies  $c\tau \nabla \partial_z s = (\nabla p - p'_c(s) \nabla s) \mathbb{1}_>$  (we recall that we assumed  $\partial_z s \in H^1(\Omega)$ ). This yields

$$\int_{\Omega^H} \mathbb{1}_> \nabla p \cdot \nabla s = c\tau \int_{\Omega^H} \nabla s \cdot \nabla \partial_z s + \int_{\Omega^H} \mathbb{1}_> p'_c(s) |\nabla s|^2. \quad (4.10)$$

The first term on the right hand side of (4.10) is

$$\begin{aligned} c\tau \int_{\Omega^H} \nabla s \cdot \nabla \partial_z s &= c\tau \int_{\Omega^H} \partial_z \left( \frac{1}{2} |\nabla s|^2 \right) = \frac{c\tau}{2} \int_{\Sigma^H} |\nabla s|^2 - \frac{c\tau}{2} \int_{\Sigma} |\nabla s|^2 \\ &= \frac{c\tau}{2} \int_{\Sigma^H} |\nabla s|^2 - \frac{1}{2c\tau} \int_{\Sigma} [p_0 - p_c(s_0)]_+^2 - \frac{c\tau}{2} \int_{\Sigma} |\partial_y s_0|^2. \end{aligned} \quad (4.11)$$

At this point, we obtained from (4.9)

$$\int_{\Omega^H} \mathbb{1}_0 \partial_y s \partial_y p + \int_{\Omega^H} \mathbb{1}_> p'_c(s) |\nabla s|^2 + \frac{c\tau}{2} \int_{\Sigma^H} |\nabla s|^2 \leq C. \quad (4.12)$$

*Step 2: Test function  $\Phi$ .* We next consider the new test function

$$\Phi := [K(s) - K(p_c^{-1}(p))]_{+} \in H^1(\Omega^H).$$

Note that  $\partial_z s > 0 \iff p > p_c(s) \iff p_c^{-1}(p) > s \iff K(p_c^{-1}(p)) > K(s)$ . This shows

$$\Phi = [K(s) - K(p_c^{-1}(p))] \mathbb{1}_0.$$

Using  $\Phi$  as a test function for (3.1a) and exploiting that  $\Phi \neq 0$  only when  $\partial_z s = 0$ , we find

$$\int_{\Omega^H} \Phi \nabla \cdot (k(s)[\nabla p + g e_z]) = c \int_{\Omega^H} \Phi \partial_z s = 0. \quad (4.13)$$

Also on the left hand side, the term  $\Phi \nabla \cdot [k(s)g e_z] = \Phi k'(s)g \partial_z s$  vanishes identically. Integration by parts in (4.13) yields, using  $\Phi = 0$  on  $\Sigma$ ,

$$\int_{\Omega^H} k(s) \nabla \Phi \cdot \nabla p = \int_{\Sigma^H} \Phi k(s) \partial_z p. \quad (4.14)$$

Because of  $\nabla \Phi = \left( \frac{1}{k(s)} \nabla s - \frac{1}{k(p_c^{-1}(p))} \frac{1}{p'_c(p_c^{-1}(p))} \nabla p \right) \mathbb{1}_0$ , we find

$$\int_{\Omega^H} \nabla s \cdot \nabla p \mathbb{1}_0 - \int_{\Omega^H} \frac{k(s)}{k(p_c^{-1}(p))} \frac{|\nabla p|^2}{p'_c(p_c^{-1}(p))} \mathbb{1}_0 = \int_{\Sigma^H} \Phi k(s) \partial_z p. \quad (4.15)$$

The first integral is  $\int_{\Omega^H} \nabla s \cdot \nabla p \mathbb{1}_0 = \int_{\Omega^H} \partial_y s \partial_y p \mathbb{1}_0$ , hence it coincides with the first term in (4.12). Since  $k(s) \mathbb{1}_0 > k(p_c^{-1}(p)) \mathbb{1}_0$ , from (4.12) we arrive at

$$\int_{\Omega^H} \frac{1}{p'_c(p_c^{-1}(p))} |\nabla p|^2 \mathbb{1}_0 + \int_{\Omega^H} \mathbb{1}_{>} p'_c(s) |\nabla s|^2 + \frac{c\tau}{2} \int_{\Sigma^H} |\nabla s|^2 \leq C, \quad (4.16)$$

where we exploited once more (4.3). At this point, we have shown (4.6a).

*Step 3: Test function  $\partial_z s$ .* To show (4.6b), we use the test function  $\partial_z s = \frac{1}{c\tau} [p - p_c(s)]_{+} \in H^1(\Omega^H)$  in (3.1a). With an integration by parts we obtain

$$\begin{aligned} & \frac{1}{c\tau} \int_{\Omega^H} k(s) \nabla p \cdot \nabla [p - p_c(s)]_{+} \\ &= \int_{\Sigma^H} \partial_z s k(s) \partial_z p - \int_{\Sigma} \partial_z s k(s_0) \partial_z p + \int_{\Omega^H} (gk'(s) - c) |\partial_z s|^2. \end{aligned} \quad (4.17)$$

We observe that, by (4.3) and (4.5b), the first two integrals on the right hand side are bounded. Furthermore, the middle term of (4.16) shows that also the last integral is bounded.

Using the algebraic manipulation  $2a(a-b) = a^2 - b^2 + (a-b)^2$ , the left hand side of (4.17) is written as

$$\begin{aligned} & \frac{1}{c\tau} \int_{\Omega^H} k(s) \nabla p \cdot \nabla [p - p_c(s)]_{+} = \frac{1}{c\tau} \int_{\Omega^H} k(s) \nabla p \cdot (\nabla p - \nabla p_c(s)) \mathbb{1}_0 \\ &= \frac{1}{2c\tau} \int_{\Omega^H} k(s) [|\nabla p|^2 + |\nabla(p - p_c(s))|^2 - |\nabla p_c(s)|^2] \mathbb{1}_0 \\ &= \frac{1}{2c\tau} \int_{\Omega^H} k(s) [\mathbb{1}_{>} |\nabla p|^2 + (c\tau)^2 |\nabla(\partial_z s)|^2 - \mathbb{1}_{>} (p'_c(s))^2 |\nabla s|^2]. \end{aligned}$$

Inequality (4.16) along with (4.5a) shows that the negative term has a bounded integral. This shows (4.6b) and concludes the proof.  $\square$

To investigate the free-boundary structured solution described in Theorem 4.2 we define the function  $h : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  with (2.1) in mind: For  $H > 0$  and  $(s_H, p_H)$  solving (3.1),  $h = h(H)$  is defined as

$$h(H) := \inf\{z_0 \in [0, H] : \partial_z s_H = 0 \text{ a.e. in } (0, L) \times (z_0, H)\}. \quad (4.18)$$

The height  $h$  marks a horizontal line such that, above that line,  $\partial_z s$  vanishes. We note that  $h \in [0, H]$  is well-defined and that  $h = H$  is possible.

**Proposition 4.4** (Free-boundary solutions). *We consider the situation of Theorem 4.2 with a sequence  $(s_H, p_H)$  of  $TW_H$ -solutions for  $H \rightarrow \infty$ . Additionally, we assume for the sequence  $H \rightarrow \infty$  that the height*

$$h(H) \text{ is bounded.} \quad (4.19)$$

*Under this assumption, a free-boundary travelling wave solution  $(s, p)$  exists. More precisely, there exists a pair  $(s, p)$  with  $s \in C_b^0(\Omega)$ ,  $\partial_z s \in L^2(\Omega)$ ,  $p \in H_{\text{loc}}^2(\Omega) \cap H_{\text{loc}}^1(\Omega \cup \Sigma)$ , satisfying (1.4)–(1.5). The solution is of free boundary type in the sense that there exists  $h^* > 0$  such that  $\partial_z s = 0$  for all  $y \in (0, L)$  and  $z \geq h^*$ . The flux satisfies*

$$F_\infty \leq g \int_0^L k(s^*(y)) dy. \quad (4.20)$$

*Under the additional regularity assumptions  $s, \partial_z s, p \in C^0(\Omega)$ , the strict inequality holds in (4.20).*

*Proof.* Let  $h^* > 0$  denote an upper bound of the function  $h(H)$ , i.e.

$$h(H) \leq h^* \text{ for all } H. \quad (4.21)$$

*Step 1: An additional a priori estimate.* We consider once more the function

$$s_H^*(y) = s_H(y, h^*) \text{ for all } y \in (0, L). \quad (4.22)$$

Let  $g_{F,H} \in \mathbb{R}$  be the number

$$g_{F,H} := g - \left( F_\infty / \int_0^L k(s_H^*(y)) dy \right), \quad (4.23)$$

and let  $\tilde{p}_H \in H^1(\Omega^H)$  be the function

$$\tilde{p}_H(y, z) := p_H(y, z) + g_{F,H} z \text{ for } (y, z) \in \Omega^H. \quad (4.24)$$

We note that these definitions reflect the observations of Lemma 2.2. We finally define  $\varphi_H \in C^2([0, 1])$  as the function

$$\varphi_H(s) := cs - (g - g_{F,H})k(s). \quad (4.25)$$

This allows to write (3.1a) in the form

$$\nabla \cdot [k(s_H) \nabla \tilde{p}_H] = \partial_z \varphi_H(s_H). \quad (4.26)$$

We observe that, by (3.1e) and the choice of  $g_{F,H}$  in (4.23),

$$\int_{\Sigma^H} k(s_H) \partial_z \tilde{p}_H = F_\infty + (g_{F,H} - g) \int_{\Sigma^H} k(s_H) = 0. \quad (4.27)$$

The test function  $\tilde{p}_H$  in (4.26) provides the identity

$$\int_{\Omega^H} \tilde{p}_H \nabla \cdot [k(s_H) \nabla \tilde{p}_H] = \int_{\Omega^H} \tilde{p}_H \partial_z \varphi_H(s_H). \quad (4.28)$$

The left hand side of (4.28) is calculated with an integration by parts, exploiting the fact that  $p_H$  on the upper boundary is constant,  $p_H \equiv p_H^*$  on  $\Sigma_H$ . In the last line of the calculation we use (4.27).

$$\begin{aligned} & \int_{\Omega^H} \tilde{p}_H \nabla \cdot [k(s_H) \nabla \tilde{p}_H] \\ &= - \int_{\Omega^H} k(s_H) |\nabla \tilde{p}_H|^2 + \int_{\Sigma^H} \tilde{p}_H k(s_H) \partial_z \tilde{p}_H - \int_{\Sigma} \tilde{p}_H k(s_H) \partial_z \tilde{p}_H \\ &= - \int_{\Omega^H} k(s_H) |\nabla \tilde{p}_H|^2 + (p_H^* + g_{F,H} H) \int_{\Sigma^H} k(s_H) \partial_z \tilde{p}_H - \int_{\Sigma} p_0 k(s_0) [\partial_z p_H + g_{F,H}] \\ &= - \int_{\Omega^H} k(s_H) |\nabla \tilde{p}_H|^2 - \int_{\Sigma} p_0 k(s_0) [\partial_z p_H + g_{F,H}]. \end{aligned}$$

The right hand side of (4.28) is treated with two integrations by parts,

$$\begin{aligned} & \int_{\Omega^H} \tilde{p}_H \partial_z \varphi_H(s_H) \\ &= - \int_{\Omega^H} \varphi_H(s_H) \partial_z \tilde{p}_H + \int_{\Sigma^H} \tilde{p}_H \varphi_H(s_H^*) - \int_{\Sigma} \tilde{p}_H \varphi_H(s_H) \\ &= - \int_{\Omega^H} \varphi_H(s_H) \partial_z \tilde{p}_H + \left[ \int_{\Omega^H} \partial_z \tilde{p}_H \varphi_H(s_H^*) + \int_0^L p_0 \varphi_H(s_H^*) \right] - \int_{\Sigma} p_0 \varphi_H(s_0) \\ &= \int_{\Omega^H} (\varphi_H(s_H^*) - \varphi_H(s_H)) \partial_z \tilde{p}_H + \int_0^L (\varphi_H(s_H^*) - \varphi_H(s_0)) p_0. \end{aligned}$$

Boundedness of many of the above terms can be concluded from the facts that  $g_{F,H}$  is bounded,  $\varphi_H \in C^1([0, 1])$ , and boundedness of  $\partial_z p$  from (4.3). From (4.28) and Young's inequality we obtain

$$\begin{aligned} & \int_{\Omega^H} k(s_H) |\nabla \tilde{p}_H|^2 \leq C - \int_{\Omega^H} (\varphi_H(s_H^*) - \varphi_H(s_H)) \partial_z \tilde{p}_H \\ & \leq C + \int_{\Omega^H} \frac{1}{2k(s_H)} |\varphi_H(s_H^*) - \varphi_H(s_H)|^2 + \int_{\Omega^H} \frac{k(s_H)}{2} |\partial_z \tilde{p}_H|^2. \end{aligned}$$

We have applied Young's inequality in such a way that the last term on the right hand side can be subtracted from both sides. Since  $\varphi_H(s_H^*) - \varphi_H(s_H) = 0$  holds for  $z \geq h^*$ , the first integral on the right hand side is bounded. We conclude

$$\int_{\Omega^H} k(s_H) |\nabla \tilde{p}_H|^2 \leq C(1 + h^*).$$

Recalling additionally the estimates from Lemma 4.3, we have the following estimates for the solution sequence:

$$\int_{\Omega^H} [|\nabla \tilde{p}_H|^2 + |\partial_z s_H|^2 + |\nabla \partial_z s_H|^2] \leq C. \quad (4.29)$$

*Step 2: Limit equations.* It remains to exploit the bounds of (4.29) to construct the limit solution for  $H \rightarrow \infty$ . Since the sequence  $g_{F,H}$  is bounded, we can choose a subsequence  $\{H_i\}_{i \in \mathbb{N}}$  with  $\lim H_i = \infty$  and  $g_F \in \mathbb{R}$  such that  $g_{F,H_i} \rightarrow g_F$ . In the following, we only use this subsequence. The estimate (4.29) allows to choose a further subsequence and a pair  $(s, p)$  with  $s \in H^1_{\text{loc}}(\Omega) \cap L^\infty(\Omega)$  and  $p \in H^1_{\text{loc}}(\Omega)$  such that, for any bounded compact subset  $\Omega' \subset \Omega$ , there holds

$$s_H \rightarrow s \text{ and } \partial_z s_H \rightarrow \partial_z s \text{ strongly in } L^2(\Omega'), \quad (4.30a)$$

$$p_H \rightarrow p \text{ weakly in } H^1(\Omega') \text{ and } p_H \rightarrow p \text{ strongly in } L^2(\Omega'). \quad (4.30b)$$

These convergences imply that also the limit  $(s, p)$  satisfies (1.4) in  $\Omega$  and the boundary conditions at the lower boundary. Furthermore,  $\partial_z s_H \equiv 0$  for all  $H$  on  $\{z \geq h^*\}$  implies  $\partial_z s \equiv 0$  on  $\{z \geq h^*\}$ .

*Step 3: Flux relations.* Regarding the flux we use that the quantity

$$F_c^H(z) := \int_0^L k(s_H(y, z))[\partial_z p_H(y, z) + g] - c s_H(y, z) dy \quad (4.31)$$

is independent of  $z \geq 0$  (compare  $F_c$  in (3.2)). Since the saturation  $s_H$  is independent of  $z$  for  $z \geq h^*$ , also the quantity

$$F^H(z) := \int_0^L k(s_H(y, z))[\partial_z p_H(y, z) + g] dy \quad (4.32)$$

is independent of  $z$  for  $z \geq h^*$ . Because of this independence and because of  $F^H(H) = F_\infty$ , we find, as  $H \rightarrow \infty$ , for every  $z \geq h^*$ ,

$$F_\infty = F^H(z) \rightarrow \int_0^L k(s(y, z))[\partial_z p(y, z) + g] dy. \quad (4.33)$$

This shows that the boundary condition (1.5a) is satisfied by the limit functions.

We have found a free boundary solution on an unbounded domain. As in Lemma 2.2, there follows  $g_F \geq 0$  and, under the regularity assumptions  $s, \partial_z s, p \in C^0(\Omega)$ , the strict inequality  $g_F > 0$ . This implies  $F_\infty = (g - g_F) \int_0^L k(s^*(y)) dy \leq g \int_0^L k(s^*(y)) dy$ , and hence (4.20).  $\square$

**Proposition 4.5** (Bounded pressure solutions). *Let the situation be that of Theorem 4.2, with  $TW_H$ -solutions  $(s_H, p_H)$  along a sequence  $H \rightarrow \infty$ . We assume here that the sequence of heights  $h(H)$  diverges,*

$$h(H) \rightarrow \infty \quad \text{as } H \rightarrow \infty. \quad (4.34)$$

*Then, a bounded pressure travelling wave solution  $(s, p)$  exists. More precisely, there exists a pair  $(s, p)$  with  $s \in C_b^0(\Omega)$ ,  $\partial_z s \in L^2(\Omega)$ ,  $p \in H^2_{\text{loc}}(\Omega) \cap H^1_{\text{loc}}(\Omega \cup \Sigma)$  satisfying (1.4). For  $C > 0$  there holds*

$$\|p\|_{L^\infty(\Omega)} + \|\nabla p\|_{L^2(\Omega)} + \|\partial_z s\|_{H^1(\Omega)} \leq C.$$

*The solution satisfies*

$$g \int_0^L k(s^*(y)) dy \leq F_\infty. \quad (4.35)$$

We note that we do not obtain the flux condition (1.5).

*Proof.* In this proof, we only write  $H \rightarrow \infty$  and  $h \rightarrow \infty$  for the two sequences. We furthermore use  $\Omega^h = [0, L] \times (0, h)$ .

*Step 1:  $L^\infty$ -bound for the pressure.* The upper bound for the pressure was assumed in (4.2),  $p_H \leq \bar{p}$  in  $\Omega^H$ . Our aim in this step is to show a lower bound for the pressure.

On the lower boundary  $\Sigma$  there holds  $p_H = p_0 \geq 0$ . We claim that there is a lower bound also along the upper boundary  $\Sigma^h$  of  $\Omega^h$ . Indeed, by definition of  $h$  in (4.18), there is a subset of non-vanishing measure in  $(0, L) \times (h-1, h)$  on which  $\partial_z s_H > 0$  holds, i.e.  $p_H > p_c(s_H) \geq 0$ . The Lipschitz bound (4.3) implies that  $p_H \geq -C_L$  holds on  $\Sigma^h$  for  $C_L = C_P \sqrt{1 + L^2}$ .

We can now exploit a maximum principle to obtain

$$-C_L \leq p_H \leq \bar{p} \quad \text{a.e. in } \Omega^h. \quad (4.36)$$

The maximum principle is derived by using  $[p_H + C_L]_-$  as a test function in (3.1a), which results in

$$\int_{\Omega^h} [p_H + C_L]_- \nabla \cdot [k(s_H) \nabla p_H] = \int_{\Omega^h} [p_H + C_L]_- (c - gk'(s_H)) \partial_z s_H.$$

An integration by parts yields

$$\begin{aligned} \int_{\Omega^h} k(s_H) |\nabla [p_H + C_L]_-|^2 &= \int_{\Sigma^h} [p_H + C_L]_- k(s_H) \partial_z p_H \\ &\quad - \int_{\Sigma} [p_H + C_L]_- k(s_H) \partial_z p_H + \int_{\Omega^h} [p_H + C_L]_- (c - gk'(s_H)) \partial_z s_H. \end{aligned}$$

As analyzed before, the boundary terms vanish because of  $p_H + C_L \geq 0$  along  $\Sigma$  and along  $\Sigma^h$ . Regarding the last integral we note that in every point  $x$  with  $\partial_z s(x) > 0$ , there holds  $p_H(x) \geq p_c(s_H(x)) \geq p_c(s_*) = 0$ , and hence  $[p_H + C_L]_- = 0$ . This shows that all terms on the right hand side vanish. We obtain (4.36).

*Step 2: A further a priori estimate.* From the uniform pressure bound (4.36) we conclude that  $p_c^{-1}(p_H)$  is bounded away from 1. With this information, the bound of (4.6a) provides, with a constant  $C > 0$  independent of  $H$ , the inequality

$$\int_{\Omega^h} \mathbb{1}_0 |\nabla p_H|^2 + \int_{\Omega^H} \mathbb{1}_> p_c'(s_H) |\nabla s_H|^2 \leq C.$$

Similarly, (4.6b) implies

$$\int_{\Omega^H} \mathbb{1}_> |\nabla p_H|^2 + \int_{\Omega^H} |\nabla (\partial_z s_H)|^2 \leq C.$$

Combining both of these inequalities with (4.36), and recalling  $\partial_z s_H = 0$  in  $\Omega^H \setminus \Omega^h$ , we obtain

$$\max_{\Omega^h} |p_H|^2 + \int_{\Omega^h} |\nabla p_H|^2 + \int_{\Omega^H} [|\partial_z s_H|^2 + |\nabla \partial_z s_H|^2] \leq C. \quad (4.37)$$

*Step 3: Limit  $H \rightarrow \infty$ .* Because of  $h \rightarrow \infty$ , we find a limiting pair  $(s, p)$  such that the local convergences of (4.30) hold for any compact subset  $\Omega'$  of  $\Omega$ . It is straightforward to verify that  $(s, p)$  solves (1.4). Moreover, (4.37) together with  $h \rightarrow \infty$  implies the additional properties  $\nabla p \in L^2(\Omega)$  (as a bounded solution to an elliptic equation) and  $\partial_z s \in L^2(\Omega)$ .

Regarding the limiting flux, we start from the relation  $\int_{\Sigma^H} k(s_H)(\partial_z p_H + g) = F_\infty$ . In order to calculate limits, we once more use the quantity  $F_c^H(z)$  of (4.31), which is independent of  $z$ . The local strong convergence of  $s_H$  and the local weak convergence of  $\nabla p_H$  yield, for almost every  $z$ , as  $H \rightarrow \infty$ ,

$$\begin{aligned} & \int_0^L k(s(y, z))[\partial_z p(y, z) + g] - cs(y, z) dy =: F_c(z) \\ & \leftarrow F_c^H(z) := \int_0^L k(s_H(y, z))[\partial_z p_H(y, z) + g] - cs_H(y, z) dy \\ & = F_c^H(H) = \int_0^L k(s_H(y, H))[\partial_z p_H(y, H) + g] - cs_H(y, H) dy \\ & = F_\infty - c \int_0^L s_H(y, H) dy. \end{aligned}$$

Because of  $s_H(y, H) \geq s_H(y, z)$  for every  $z$ , and  $s_H \rightarrow s$ , there holds

$$\lim_{z \rightarrow \infty} \int_0^L s(y, z) dy \leq \lim_{H \rightarrow \infty} \int_0^L s_H(y, H) dy.$$

Taking in the above calculation both limits,  $z \rightarrow \infty$  and  $H \rightarrow \infty$ , exploiting  $\nabla p \in L^2(\Omega)$ , we find

$$\int_0^L gk(s^*(y)) dy = \lim_{z \rightarrow \infty} \int_0^L gk(s(y, z)) dy \leq F_\infty.$$

This concludes the proof.  $\square$

**Remark 4.6** (Both solution types occur). *The one-dimensional travelling wave results in [9] indicate that both solution types exist for a given  $s_* \in (0, 1)$  and  $F_\infty$  satisfying (4.1). Type I (large) solutions occur in the one-dimensional model when  $\tau$  is large. On the other hand, if  $\|p_0 - p_c(s_*)\|_{L^\infty(\Sigma)}$  is small, then Type II (small) solutions are expected to occur for small  $\tau$  values. Our numerical results confirm that both solution types occur.*

We finally want to show that, for a given flux  $F_\infty$ , it is possible to find a wave-speed  $c$  such that condition (3.4) is satisfied.

**Theorem 4.7** (Selecting a wave-speed  $c$  in dependence of  $F_\infty$  and  $s_*$ ). *Let  $\tau > 0$ ,  $s_* \in (0, 1)$ , and boundary data  $s_0, p_0 \in C^1(\Sigma)$  be given,  $p_c(s_*) \leq p_c(s_0) < p_0$  on  $\Sigma$ , furthermore  $F_\infty$  in the bounds of (4.1). We assume that, for all  $c \in [c_1, c_2]$  with  $c_1 := k'(s_*)g$  and  $c_2 := g(k(1) - k(s_*))/(1 - s_*)$ , a sequence  $(s_H, p_H)$  of solutions to (3.1) satisfying Assumption 4.1 exists. We consider the corresponding limit solutions  $(s, p)$  and their fluxes*

$$F_c = \int_\Sigma (k(s_0)[\partial_z p + g] - cs_0), \quad (4.38)$$

and assume that  $F_c$  depends continuously on  $c$ . Then there exists a wave-speed  $\bar{c} \in (c_1, c_2)$  such that the corresponding pair  $(s, p)$  satisfies (3.4),  $F_c = (gk(s_*) - cs_*)L$ .

*Proof.* We consider the continuous function  $G : [c_1, c_2] \rightarrow \mathbb{R}$

$$G(c) := F_c - (gk(s_*) - cs_*)L. \quad (4.39)$$

We recall that  $G$  depends in an explicit way on  $c$ , but also implicitly, since  $s$  and  $p$  (and hence  $F_c$ ) depend on  $c$ . The flux quantity  $F_c$  is independent of  $z$ , we choose to evaluate it at  $z \rightarrow \infty$ . We denote the limit of the first two terms as

$$F_0 := F_0(c) := \lim_{z \rightarrow \infty} \int_0^L k(s(y, z)) [\partial_z p(y, z) + g] dy.$$

We observe that, by Theorem 4.2,

$$F_0 = \begin{cases} F_\infty & \text{if } g \int_0^L k(s^*) > F_\infty, \\ g \int_0^L k(s^*) & \text{if } g \int_0^L k(s^*) \leq F_\infty. \end{cases}$$

In both cases holds  $F_0 \leq F_\infty$  and  $F_0 \leq g \int_0^L k(s^*)$ . The function  $G$  can be written as

$$G(c) = F_0 - k(s_*)gL - c \int_0^L (s^*(y) - s_*) dy.$$

Showing  $G(c) > 0$  as  $c \rightarrow c_1$ . If the solution is of Type II (small solution, second case in the above distinction), then

$$G(c) = \int_0^L (s^*(y) - s_*) dy \left( g \frac{\int_0^L (k(s^*(y)) - k(s_*)) dy}{\int_0^L (s^*(y) - s_*) dy} - c \right).$$

We exploit that  $s^* > s_0 \geq s_*$  implies, for every  $y \in (0, L)$ , that  $k(s^*(y)) - k(s_*) > k'(s_*)(s^*(y) - s_*)$ . This implies that, for  $c$  close to  $c_1 = k'(s_*)g$ , there holds  $G(c) > 0$ . On the other hand, if  $(s, p)$  is of Type I (large solutions), then

$$\begin{aligned} G(c) &= F_\infty - k(s_*)gL - c \int_0^L (s^*(y) - s_*) dy \\ &\geq F_\infty - k(s_*)gL - c(1 - s_*)L \\ &\geq gLk'(s_*)(1 - s_*) - c(1 - s_*)L + \varepsilon \\ &= (gk'(s_*) - c)(1 - s_*)L + \varepsilon, \end{aligned}$$

where we exploited the lower bound  $gL[k(s_*) + k'(s_*)(1 - s_*)] + \varepsilon \leq F_\infty$  for some  $\varepsilon > 0$ . We see that, also in this case, for  $c$  close to  $c_1 = k'(s_*)g$ , there holds  $G(c) > 0$ .

Showing  $G(c) < 0$  as  $c \rightarrow c_2$ . Consider solutions of Type II (small solutions). For  $\mu := (k(1) - k(s_*))/(1 - s_*)$ , we show that in this case, there exists  $\varepsilon > 0$  independent of  $c \in [c_1, c_2]$  such that

$$\int_0^L (k(s^*) - k(s_*)) \leq (\mu - \varepsilon) \int_0^L (s^* - s_*). \quad (4.40)$$

Since  $(k(s) - k(s_*))/(s - s_*)$  is a strictly increasing function for  $s > s_*$ ,  $\int_0^L (k(s^*) - k(s_*)) = \mu \int_0^L (s^* - s_*)$  if and only if  $s^*(y) \in \{s_*, 1\}$  for all  $y \in (0, L)$ . From Jensen's inequality, one has

$$k\left(\frac{1}{L} \int_0^L s^*\right) \leq \frac{1}{L} \int_0^L k(s^*) \leq \frac{F_\infty}{gL} < k(1), \quad (4.41)$$

implying  $\frac{1}{L} \int_0^L s^* < 1$ . Hence, the possibility  $s^* \equiv 1$  in  $(0, L)$  is ruled out. Moreover, since  $s^* > s_0 \geq s_*$ , the possibility  $s^* \equiv s_*$  in  $(0, L)$  is also ruled out. From Lemma A.3,

$\|\nabla s\|_{L^\infty(\Omega)}$  is bounded. Hence  $s^*$  cannot take both the values  $s_*$  and 1 without transitioning through the intermediate values. Thus (4.40) holds.

If the solution is of Type II, then, for  $c$  close enough to  $g\mu = g(k(1) - k(s_*))/(1 - s_*)$ , we obtain from (4.40),

$$\begin{aligned} G(c) &= F_0 - k(s_*)gL - c \int_0^L (s^* - s_*) = \int_0^L [g(k(s^*) - k(s_*)) - c(s^* - s_*)] \\ &\leq \int_0^L (s^* - s_*) [g\mu - g\varepsilon - c] \leq 0. \end{aligned}$$

If the solution is of Type I, then for  $g_F > 0$  as defined in (2.5) (see also Proposition 4.4), one has

$$\begin{aligned} G(c) &= F_\infty - k(s_*)gL - c \int_0^L (s^* - s_*) \\ &= \int_0^L [g(k(s^*) - k(s_*)) - c(s^* - s_*)] - g_F \int_0^L k(s^*) \\ &\leq (g\mu - c) \int_0^L (s^* - s_*) - g_F \int_0^L k(s_0). \end{aligned}$$

Consequently,  $G(c) < 0$  for  $c$  close enough to  $c_2 = g\mu$ . Hence, there exists a zero  $\bar{c}$  of  $G(\cdot)$  in  $(c_1, c_2)$ . This was the claim.  $\square$

## 5 Numerics

### 5.1 Numerical solution of system (3.1)

The primary numerical task is to solve system (3.1) for  $s$  and  $p$ , where the speed  $c$  and the total influx  $F_\infty$  are given. The existence of a solution was established in Theorem 3.4. We use an iterative method in order to deal with the nonlinearities. With a positive number  $M > 0$ , we use the iteration  $(s^{i-1}, p^{i-1}) \mapsto (s^i, p^i)$  that is given by

$$Mp^i - \nabla \cdot [k(s^{i-1})(\nabla p^i + ge_z)] = Mp^{i-1} - \frac{1}{\tau} [p^{i-1} - p_c(s^{i-1})]_+, \quad (5.1a)$$

$$\partial_z s^i - \varepsilon \Delta s^i = \frac{1}{c\tau} [p^i - p_c(s^{i-1})]_+. \quad (5.1b)$$

The equations are solved in the rectangular computational domain  $\Omega^H$  for some initial guess  $(s^0, p^0)$ . They are supplemented by the boundary conditions (3.1c)–(3.1e) and no-flux conditions at the lateral boundaries.

For  $\varepsilon = 0$ , a fixed point of the iteration scheme (5.1) provides a solution of (3.1). The set-up is such that the equations can be solved subsequently: One can solve the first equation for  $p^i$ , then the second equation for  $s^i$ . The iteration strategy is based on the L-scheme [18], the iteration is expected to converge for  $M \geq \tau^{-1}$ , irrespective of the initial guess. We introduce an elliptic regularization in the second equation (which is first order in  $s$ ), numerical experiments are run with a small number  $\varepsilon > 0$ .

In order to discretize (5.1), we introduce a uniform triangulation  $\Omega_h^H$  of the domain  $\Omega^H$  and apply linear finite elements. In this sense, the discretization is based on the weak formulation in (3.5)–(3.6). The resulting scheme has been implemented

in the adaptive finite element tool box AMDiS [28]. The linear equations arising from the discretization are treated with the direct solver UMFPAK, [8].

The physical parameters of the problem are chosen as in [15],

$$p_c(s) = s, \quad k(s) = \begin{cases} \kappa & \text{for } s < a, \\ \kappa + (s - a)^2 & \text{for } s \geq a, \end{cases} \quad (5.2)$$

and

$$g = 1, \quad \tau = 2, \quad \kappa = 0.001, \quad a = 0.32, \quad F_\infty = 0.056.$$

The domain is  $\Omega^H = (-1, 1)^2$ ; up to a shift of the domain, this coincides with  $L = 2$  and  $H = 2$  in analytical results. The parameters for the numerical code are

$$M = 4, \quad \varepsilon = 0.0008.$$

The initial values for the iteration have been chosen as

$$p^0 = 4.5, \quad s^0 = 10^{-5}.$$

Regarding the lower boundary, we use the constant function  $s_0 = 10^{-5}$  and the slightly perturbed pressure boundary condition

$$p_0(y) = p_c(s_0) + \delta e^{-(y/d)^2}.$$

The positive parameter  $\delta = 0.078$  measures the amplitude of the perturbation and the scaling factor  $d = 0.25$  measures the width of the perturbation.

Figure 5 shows results for four different values of the speed  $c$ . We see a remarkable difference between the solution for  $c = 0.04785$  and the solution for  $c = 0.04786$ . The abrupt change finds its counterpart in Theorem 4.2 (we recall that the theorem is treating unbounded domains while the numerical results are for a fixed bounded domain): The two images on the left show Type I solutions, i.e., “large solutions” with a free boundary. The two images on the right show “small solutions”.




Figure 5: The discrete solutions  $s_h$  of the iteration scheme for (from left to right)  $c = 0.04$ ,  $c = 0.04785$ ,  $c = 0.04786$ ,  $c = 0.05013166020$ .

In the above experiments, we have solved system (3.1) for different values of  $c$ . We now ask: What is the correct wave speed  $c$  in the sense of (3.4)? We use the following finite domain approximations:  $s_*$  can be neglected, hence, in particular,  $k(s_*) = \kappa$ . Furthermore,  $s_0$  is constant and so small that also  $k(s_0)$  can be replaced by  $\kappa$ . Condition (3.4) then reads  $G_1(c) := c - (\kappa \int_\Sigma \partial_z p_h) / (L s_0) \stackrel{!}{=} 0$ . We find the values as displayed in Table 1. We conclude that  $G_1(\bar{c}_1) = 0$  is satisfied for some  $\bar{c}_1 \in [0.0476, 0.0477]$ . Up to the above finite domain approximations, we expect the travelling wave speed to be about 0.0477. This is remarkably close to the jump point, compare Figure 5. We furthermore note that the value is not far from the value  $c = 0.053$  that can be extracted from simulation results reported in [15].


$c$	0.0476	0.0477
$G_1(c)$	$0.0476 + 0.0218 > 0$	$0.0477 - 0.3304 < 0$

Table 1: Values for  $G_1(c)$  for various  $c$ -values.

## 5.2 Path-following algorithm to adjust $c$

So far, for each value of  $c$ , we started the iterative scheme (5.1) with constant functions  $s^0$  and  $p^0$  as initial guess. Since we are interested in solutions for a whole range of  $c$ -values, there is a very natural idea to speed up calculations: After having changed the value of  $c$ , instead of starting the iterative scheme from scratch, we start the iteration with the solution of the last value of  $c$ . Thereby, we increased  $c$  in every iteration step by  $10^{-4}$  in some experiments, by  $10^{-11}$  in others.

Interestingly, it turns out that this scheme produces results that are different from those reported in Section 5.1. Results are displayed in Figure 6, and once more, we observe that, below a critical value for  $c$ , solutions are “large solutions”, above the critical value, we find “small solutions”. This feature is as in the sequence of Figure 5, but the critical value of  $c$  is now different: It is about  $\bar{c}_2 = 0.050$  and no longer about  $\bar{c}_1 = 0.048$ . For values of  $c$  below  $\bar{c}_1$  and for values above  $\bar{c}_2$ , the results of the two schemes coincide.

Figure 6: Plots of the discrete solutions  $s_h$  of the path-following iteration scheme for (from left to right)  $c = 0.04$ ,  $c = 0.05013166020$ ,  $c = 0.05013166023$ ,  $c = 0.0625$ .

We conclude with an evaluation of the integral condition in Theorem 4.2, where the criterion for a “large solution” was  $g \int_0^L k(s^*(y)) dy \geq F_\infty$  for the saturation values  $s^*(y) := \lim_{z \rightarrow \infty} s(y, z)$  at infinity. With the approximation  $s^* \approx s|_{\Sigma_H}$  and with (3.1e), the criterion for a “large solution” reads

$$G_2(c) := \int_{\Sigma_H} k(s) \partial_z p \leq 0. \quad (5.3)$$

Our simulations yield the values in Table 2. We observe that the change of sign of  $G_2$  occurs only after the point that the solution switched to the “small solution”.

$c$	0.04	0.05013166020	0.05013166023	0.0625
$G_2(c)$	-0.1948	-0.1630	-0.1492	0.0183

Table 2: Values for  $G_2(c)$  for various  $c$ -values.

Our observations may be interpreted as follows: For a range of values of  $c$ , there are two solutions of system (3.1). This is not in contradiction with our analysis, since Theorem 3.4 provides the existence, but not the uniqueness of solutions. A numerical

scheme has the tendency to find the “stable” solution (“stable” has to be interpreted appropriately). In a path-following code as described here (in Section 5.2), due to numerical stabilization aspects, the code can follow one path beyond the point where it loses stability. We conjecture that this is what is visible in the observation  $\bar{c}_2 > \bar{c}_1$ .

## Conclusions

We studied the travelling wave equations for a porous media imbibition problem with hysteresis. Denoting by  $c$  the unknown speed of the travelling wave, we treat a free boundary problem with an additional parameter. Our analysis shows that, after a domain truncation and for boundary conditions within physically reasonable limits: (i) For a prescribed speed  $c$ , travelling wave solutions exist. In the limit of infinite domains, different types of limit solutions can occur. (ii) A critical wave speed  $c$  can be selected by a flux condition. (iii) Numerical experiments provide solutions with the shape of a finger. We find values of  $c$  that are in good agreement with time-dependent calculations. Different numerical algorithms yield slightly different values for  $c$ , an effect that may be related to non-uniqueness of solutions.

## A Appendix

The following result on solution sequences  $(s_H, p_H)$  does not rely on Assumption 4.1, but follows directly from the variational principle.

**Lemma A.1** (Large solution sequences have unbounded pressure). *For a sequence  $0 < H \rightarrow \infty$ , let  $(s_H, p_H)$  be solutions to (3.1). We assume that, for some height parameter  $z_0 > 0$  and some bound  $C_k > 0$ , every solution  $s_H$  satisfies the integral condition*

$$\int_0^L gk(s_H(y, z_0)) dy \geq C_k > F_\infty. \quad (\text{A.1})$$

*In this situation, the sequence of pressure functions is unbounded,*

$$\|p_H\|_{L^\infty} \rightarrow \infty. \quad (\text{A.2})$$

*In particular, it generates a “large” Type I solution.*

*Proof.* For a contradiction argument we assume that, for some  $\bar{p} > 0$ , the pressure functions are bounded,  $|p_H| \leq \bar{p}$  on  $\Omega^H$ . We recall that  $p_H$  is the minimizer for the functional  $A$  of (3.8), for given  $s = s_H$ . This provides a lower bound for  $A$ : For any function  $\varphi \in X_{p_0}$ , there holds, by Lemma A.2,

$$\begin{aligned} A(\varphi) &\geq A(p_H) \geq \int_{\Omega^H} \frac{1}{2} k(s_H) |\nabla p_H + g e_z|^2 - F_\infty L \bar{p} \\ &\geq \frac{1}{2} g^2 \left( \int_{\Omega^H} k(s_H) \right) - C_1(\bar{p}) - F_\infty L \bar{p}. \end{aligned}$$

Our aim is to find a contradiction, which we obtain by constructing a comparison function with lower energy. We choose a function  $\tilde{p}_H$  that connects, in the domain  $\{z \in (0, 1)\}$ , the boundary data  $p_0$  in a smooth way with  $\tilde{p}_H \equiv 0$  for  $z = 1$ . For larger

$z$ , we set  $\tilde{p}_H(y, z) = -g_F(z - 1)$ , where the coefficient  $g_F \in (0, g)$  is chosen below. We calculate for the energy

$$A(\tilde{p}_H) \leq C_2 + \frac{1}{2}|g - g_F|^2 \left( \int_{\Omega^H} k(s_H) \right) + F_\infty H g_F.$$

Combining the two inequalities and using  $\bar{C}_k := \left( g \int_0^L \int_{z_0}^H k(s_H) \right) / (H - z_0)$ , we find

$$\frac{1}{2} g \bar{C}_k H \leq C_3 + F_\infty H g_F + \frac{H \bar{C}_k}{2g} |g - g_F|^2. \quad (\text{A.3})$$

Optimizing in  $g_F$  leads to the choice  $g_F := g - q$  with  $q := (g F_\infty) / \bar{C}_k < g$ . In order to compare the prefactors of  $H$  on both sides we study

$$\begin{aligned} \frac{1}{2} \bar{C}_k g - \frac{\bar{C}_k}{2g} |g - g_F|^2 - F_\infty g_F &= \frac{1}{2} \bar{C}_k g - \frac{\bar{C}_k}{2g} |q|^2 - F_\infty (g - q) \\ &= \frac{\bar{C}_k}{2g} (g^2 - q^2 - 2q(g - q)) = \frac{\bar{C}_k}{2g} (g^2 + q^2 - 2qg) = \frac{\bar{C}_k}{2g} (g - q)^2 > 0. \end{aligned}$$

For large  $H$ , this yields a contradiction in (A.3).  $\square$

**Lemma A.2** (A Jensen type inequality). *For  $\Omega^H = (0, L) \times (0, H)$  with points  $x = (y, z)$ ,  $k : \Omega^H \rightarrow [0, k_0]$  monotonically increasing in  $z$ , and  $u : \Omega^H \rightarrow \mathbb{R}$  with the uniform bound  $\|u\|_{L^\infty} \leq \bar{u}$ , there exists a constant  $C_1 = C_1(\bar{u}, k_0)$ , independent of  $H$ , such that*

$$\int_{\Omega^H} k |\nabla u + g e_z|^2 \geq -C_1 + g^2 \int_{\Omega^H} k. \quad (\text{A.4})$$

*Proof.* We use the averaging operator  $M : L^2(\Omega^H) \rightarrow \mathbb{R}$ , defined by

$$M(v) := \left( \int_{\Omega^H} k v \right) / \left( \int_{\Omega^H} k \right).$$

This operator is linear and maps the constant function  $v \equiv a \in \mathbb{R}$  to  $M(v) = a$ . We furthermore use the convex function  $\psi : \mathbb{R} \rightarrow \mathbb{R}$ ,  $\xi \mapsto |\xi + g|^2$ . Jensen's inequality provides

$$M(\psi(\partial_z u)) \geq \psi(M(\partial_z u)).$$

In our setting and with  $m := \int_{\Omega^H} k$ , this yields

$$\int_{\Omega^H} k |\nabla u + g e_z|^2 \geq \int_{\Omega^H} k |\partial_z u + g|^2 = m M(\psi(\partial_z u)) \geq m \psi(M(\partial_z u)).$$

We calculate, using that  $k$  is increasing in  $z$ ,

$$\begin{aligned} |M(\partial_z u)| &= \left| \frac{1}{m} \int_{\Omega^H} k \partial_z u \right| \leq \frac{1}{m} \left| \int_0^L k u \Big|_0^H - \int_{\Omega^H} \partial_z k u \right| \\ &\leq \frac{1}{m} \left( 2k_0 \bar{u} + \bar{u} \int_{\Omega^H} \partial_z k \right) \leq \frac{3k_0 \bar{u}}{m}. \end{aligned}$$

Inserting above we obtain

$$\int_{\Omega^H} k |\nabla u + g e_z|^2 \geq m \psi(M(\partial_z u)) = m |g + M(\partial_z u)|^2 \geq m g^2 - 6g k_0 \bar{u}.$$

This shows the claim.  $\square$

**Lemma A.3** (Lipschitz continuity of  $s_H$ ). *Let  $F_\infty, c, \tau, s_* > 0$  and  $s_0, p_0 \in C^1(\Sigma)$  with  $s_* \leq s_0 < 1$  and  $p_0 \geq p_c(s_0)$  be fixed. For  $H > 0$ , let  $(s_H, p_H)$  be the  $TW_H$ -solution to (3.1) satisfying (4.3) and (4.4). Then, for  $\rho = \min\{p_c'\} > 0$ , there holds*

$$\|\partial_z s_H\|_{L^\infty(\Omega^H)}, \|\partial_y s_H\|_{L^\infty(\Omega^H)} \leq C_P/\rho + \|\partial_y s_0\|_{L^\infty(\Sigma)} + \frac{1}{c\tau} \|p_0 - p_c(s_0)\|_{L^\infty(\Sigma)} =: C_s. \quad (\text{A.5})$$

*Proof.* To prove the lemma, we consider a regularization of the signum function, denoted as  $\text{sign}_\varepsilon : \mathbb{R} \rightarrow [-1, 1]$ . A possible choice is  $\text{sign}_\varepsilon(\eta) := \eta/\varepsilon$  for  $\eta \in [-\varepsilon, \varepsilon]$ ,  $\text{sign}_\varepsilon(\eta) := -1$  for  $\eta < -\varepsilon$  and  $\text{sign}_\varepsilon(\eta) := 1$  for  $\eta > \varepsilon$ . We also introduce the primitive  $H_\varepsilon(\eta) = \int_0^\eta \text{sign}_\varepsilon(\rho) d\rho$ . We demand that, as  $\varepsilon \rightarrow 0$ , there holds  $\text{sign}_\varepsilon(\eta) \rightarrow \text{sign}(\eta)$ ,  $H_\varepsilon(\eta) \rightarrow |\eta|$ , and  $\eta \text{sign}_\varepsilon(\eta) \rightarrow |\eta|$ .

We differentiate relation (3.1b), in the sense of distributions, with respect to  $x_j$  for  $x_j = y$  and for  $x_j = z$ . The regularity assumption (4.4) on  $s_H$  allows to write

$$c\tau \partial_z \partial_{x_j} s_H + p_c'(s_H) \partial_{x_j} s_H \text{sign}(\partial_z s_H) = \partial_{x_j} p_H \text{sign}(\partial_z s_H).$$

Multiplying both sides with  $\text{sign}_\varepsilon(\partial_{x_j} s_H)$  yields

$$\begin{aligned} c\tau \partial_z H_\varepsilon(\partial_{x_j} s_H) + p_c'(s_H) \text{sign}_\varepsilon(\partial_{x_j} s_H) \partial_{x_j} s_H \text{sign}(\partial_z s_H) \\ = \text{sign}(\partial_z s_H) \text{sign}_\varepsilon(\partial_{x_j} s_H) \partial_{x_j} p_H. \end{aligned} \quad (\text{A.6})$$

Passing to the limit  $\varepsilon \rightarrow 0$ , we obtain for  $x_j = z$  the relation

$$c\tau \partial_z |\partial_z s_H| + p_c'(s_H) |\partial_z s_H| \leq |\partial_z p_H| \leq C_P, \quad (\text{A.7})$$

where we used (4.3) in the last inequality. We exploit that, for  $z = 0$ , there holds  $\partial_z s_H = \frac{1}{c\tau} [p_0 - p_c(s_0)]_+$ , and hence also  $|\partial_z s_H| \leq \frac{1}{c\tau} \|p_0 - p_c(s_0)\|_{L^\infty(\Sigma)}$ . Inequality (A.7) implies that  $|\partial_z s_H|$  cannot exceed the value  $C_s$  of (A.5).

We now study  $x_j = y$  in (A.6). In the limit  $\varepsilon \rightarrow 0$ , exploiting  $\partial_z s_H \geq 0$ , we find

$$c\tau \partial_z |\partial_y s_H| + p_c'(s_H) |\partial_y s_H| \text{sign}(\partial_z s_H) \leq |\partial_y p_H| \text{sign}(\partial_z s_H). \quad (\text{A.8})$$

With the uniform bound  $|\nabla p_H| \leq C_P$  of (4.3) we can write

$$c\tau \partial_z |\partial_y s_H| \leq (C_P - \rho |\partial_y s_H|) \text{sign}(\partial_z s_H). \quad (\text{A.9})$$

For  $z = 0$ , there holds  $\partial_y s_H = \partial_y s_0$ . Inequality (A.9) implies that  $|\partial_y s_H|$  cannot exceed the value  $C_s$  of (A.5).  $\square$

## References

- [1] H.-W. Alt and S. Luckhaus. Quasilinear elliptic-parabolic differential equations. *Math. Z.*, 183(3):311–341, 1983.
- [2] J. Bear. *Hydraulics of groundwater*. McGraw-Hill International Book Co., 1979.
- [3] A.Y. Beliaev and S.M. Hassanizadeh. A theoretical model of hysteresis and dynamic effects in the capillary relation for two-phase flow in porous media. *Transport in Porous Media*, 43(3):487–510, 2001.

- [4] A.Y. Beliaev and R.J. Schotting. Analysis of a new model for unsaturated flow in porous media including hysteresis and dynamic effects. *Computational Geosciences*, 5(4):345–368 (2002), 2001.
- [5] A. Beljadid, L. Cueto-Felgueroso, and R. Juanes. A continuum model of unstable infiltration in porous media endowed with an entropy function. *Advances in Water Resources*, page 103684, 2020.
- [6] X. Cao and I.S. Pop. Two-phase porous media flows with dynamic capillary effects and hysteresis: uniqueness of weak solutions. *Comput. Math. Appl.*, 69(7):688–695, 2015.
- [7] L. Cueto-Felgueroso and R. Juanes. Nonlocal interface dynamics and pattern formation in gravity-driven unsaturated flow through porous media. *Physical Review Letters*, 101(24):244504, 2008.
- [8] T. A. Davis. Algorithm 832: UMFPACK V4.3—an unsymmetric-pattern multifrontal method. *ACM Transactions on Mathematical Software*, 30(2):196–199, 2004.
- [9] E. El Behi-Gornostaeva, K. Mitra, and B. Schweizer. Traveling wave solutions for the Richards equation with hysteresis. *IMA Journal of Applied Mathematics*, 84(4):797–812, 2019.
- [10] D. Gilbarg and N.S. Trudinger. *Elliptic partial differential equations of second order*. springer, 2015.
- [11] R.J. Glass, T.S. Steenhuis, and J.Y. Parlange. Mechanism for finger persistence in homogeneous, unsaturated, porous media: Theory and verification. *Soil Science*, 148(1):60–70, 1989.
- [12] S.M. Hassanizadeh and W.G. Gray. Thermodynamic basis of capillary pressure in porous media. *Water Resources Research*, 29(10):3389–3405, 1993.
- [13] R. Helmig. *Multiphase flow and transport processes in the subsurface: a contribution to the modeling of hydrosystems*. Springer-Verlag, 1997.
- [14] J. Koch, A. Rätz, and B. Schweizer. Two-phase flow equations with a dynamic capillary pressure. *European Journal of Applied Mathematics*, 24(1):49–75, 2013.
- [15] A. Lamacz, A. Rätz, and B. Schweizer. A well-posed hysteresis model for flows in porous media and applications to fingering effects. *Advances in Mathematical Sciences and Applications*, 21(1):33–64, 2011.
- [16] K. Mitra. Existence and properties of solutions of extended play-type hysteresis model. *arXiv*, arXiv:2009.03209, 2020.
- [17] K. Mitra, T. Köppl, C.J. van Duijn, I.S. Pop, and R. Helmig. Fronts in two-phase porous media flow problems: the effects of hysteresis and dynamic capillarity. *arXiv preprint arXiv:1906.08134*, 2019.

- [18] K. Mitra and I.S. Pop. A modified L-scheme to solve nonlinear diffusion problems. *Computers & Mathematics with Applications*, 77(6):1722 – 1738, 2019. 7th International Conference on Advanced Computational Methods in Engineering (ACOMEN 2017).
- [19] K. Mitra and C.J. van Duijn. Wetting fronts in unsaturated porous media: The combined case of hysteresis and dynamic capillary pressure. *Nonlinear Analysis: Real World Applications*, 50:316 – 341, 2019.
- [20] F. Otto.  $L^1$ -contraction and uniqueness for unstationary saturated-unsaturated porous media flow. *Advances in Mathematical Sciences and Applications*, 7(2):537–553, 1997.
- [21] A. Rätz and B. Schweizer. Hysteresis models and gravity fingering in porous media. *Zeitschrift für Angewandte Mathematik und Mechanik*, 94(7-8):645–654, 2014.
- [22] L.A. Richards. Capillary conduction of liquids through porous mediums. *Journal of Applied Physics*, 1(5):318–333, 1931.
- [23] B. Schweizer. Instability of gravity wetting fronts for Richards equations with hysteresis. *Interfaces and Free Boundaries*, 14(1):37–64, 2012.
- [24] B. Schweizer. The Richards equation with hysteresis and degenerate capillary pressure. *Journal of Differential Equations*, 252(10):5594 – 5612, 2012.
- [25] B. Schweizer. Hysteresis in porous media: Modelling and analysis. *Interfaces and Free Boundaries*, 19(3):417–447, 2017.
- [26] J. Selker, J-Y. Parlange, and T. Steenhuis. Fingered flow in two dimensions: 2. Predicting finger moisture profile. *Water Resources Research*, 28(9):2523–2528, 1992.
- [27] C.J. van Duijn, K. Mitra, and I.S. Pop. Travelling wave solutions for the Richards equation incorporating non-equilibrium effects in the capillarity pressure. *Nonlinear Analysis: Real World Applications*, 41(Supplement C):232 – 268, 2018.
- [28] S. Vey and A. Voigt. AMDiS: adaptive multidimensional simulations. *Computing and Visualization in Science*, 10(1):57–67, 2007.

## Preprints ab 2015/06

- 2020-05 **Koondanibha Mitra, Ben Schweizer, Andreas Rätz**  
Travelling wave solutions for gravity fingering in porous media flows
- 2020-04 **Sergio Andraus, Kilian Hermann and Michael Voit**  
Limit theorems and soft edge of freezing random matrix models via dual orthogonal polynomials
- 2020-03 **Michael Voit and Jeannette H.C. Woerner**  
Limit theorems for Bessel and Dunkl processes of large dimensions and free convolutions
- 2020-02 **Patrizia Donato, Agnes Lamacz, Ben Schweizer**  
Sound absorption by perforated walls along boundaries
- 2020-01 **Christoph Schumacher**  
Habilitationsschrift: Concentration inequalities in random Schrödinger operators
- 2019-06 **Matthias Röger and Ben Schweizer**  
Relaxation analysis in a data driven problem with a single outlier
- 2019-05 **Klaas Hendrik Poelstra, Ben Schweizer and Maik Urban**  
The geometric average of curl-free fields in periodic geometries
- 2019-04 **Agnes Lamacz and Ben Schweizer**  
Representation of solutions to wave equations with profile functions
- 2019-03 **Ben Schweizer**  
Existence results for the Helmholtz equation in periodic wave-guides with energy methods
- 2019-02 **Ben Schweizer and Maik Urban**  
On a limiting absorption principle for sesquilinear forms with an application to the Helmholtz equation in a waveguide
- 2019-01 **Michael Voit and Jeannette H.C. Woerner**  
Functional central limit theorems for multivariate Bessel processes in the freezing regime
- 2018-08 **Ben Schweizer**  
Effective Helmholtz problem in a domain with a Neumann sieve perforation
- 2018-07 **Sergio Andraus and Michael Voit**  
Limit theorems for multivariate Bessel processes in the freezing regime
- 2018-06 **Michael Voit**  
Central limit theorems for multivariate Bessel processes in the freezing regime
- 2018-05 **Elena El Behi-Gornostaeva, Koondanibha Mitra and Ben Schweizer**  
Traveling wave solutions for the Richards equation with hysteresis
- 2018-04 **Mario Ohlberger, Ben Schweizer, Maik Urban and Barbara Verfürth**  
Mathematical analysis of transmission properties of electromagnetic meta-materials
- 2018-03 **Margit Rösler and Michael Voit**  
Beta distributions and Sonine integrals for Bessel functions on symmetric cones
- 2018-02 **Michael Voit**  
Continuous Association Schemes and Hypergroups

- 2018-01 **Merdan Artykov and Michael Voit**  
Some central limit theorems for random walks associated with hypergeometric functions of type BC
- 2017-05 **Ben Schweizer and Florian Theil**  
Lattice dynamics on large time scales and dispersive effective equations
- 2017-04 **Frank Klinker and Christoph Reineke**  
A note on the regularity of matrices with uniform polynomial entries
- 2017-03 **Tomáš Dohnal and Ben Schweizer**  
A Bloch wave numerical scheme for scattering problems in periodic wave-guides
- 2017-02 **Matthias Röger and Ben Schweizer**  
Strain gradient visco-plasticity with dislocation densities contributing to the energy
- 2017-01 **Ben Schweizer and Maik Urban**  
Effective Maxwell's equations in general periodic microstructures
- 2016-05 **Robert Lipton and Ben Schweizer**  
Effective Maxwell's equations for perfectly conducting split ring resonators
- 2016-04 **Ben Schweizer**  
Resonance meets homogenization - Construction of meta-materials with astonishing properties
- 2016-03 **Ben Schweizer**  
On Friedrichs inequality, Helmholtz decomposition, vector potentials, and the div-curl lemma
- 2016-02 **Michael Voit**  
Generalized commutative association schemes, hypergroups, and positive product formulas
- 2016-01 **Agnes Lamacz and Ben Schweizer**  
Effective acoustic properties of a meta-material consisting of small Helmholtz resonators
- 2015-13 **Christian Eggert, Ralf Gäer, Frank Klinker**  
The general treatment of non-symmetric, non-balanced star circuits: On the geometrization of problems in electrical metrology
- 2015-12 **Daniel Kobe and Jeannette H.C. Woerner**  
Oscillating Ornstein-Uhlenbeck processes and modelling electricity prices
- 2015-11 **Sven Glaser**  
A distributional limit theorem for the realized power variation of linear fractional stable motions
- 2015-10 **Herold Dehling, Brice Franke and Jeannette H.C. Woerner**  
Estimating drift parameters in a fractional Ornstein-Uhlenbeck process with periodic mean
- 2015-09 **Harald Garcke, Johannes Kammann, Andreas Rätz and Matthias Röger**  
A coupled surface-Cahn-Hilliard bulk-diffusion system modeling lipid raft formation in cell membranes
- 2015-08 **Agnes Lamacz and Ben Schweizer**  
Outgoing wave conditions in photonic crystals and transmission properties at interfaces
- 2015-07 **Manh Hong Duong, Agnes Lamacz, Mark A. Peletier and Upanshu Sharma**  
Variational approach to coarse-graining of generalized gradient flows
- 2015-06 **Agnes Lamacz and Ben Schweizer**  
A negative index meta-material for Maxwell's equations