



Γ -Convergence for Plane to Wrinkles Transition Problem

Peter Bella¹ · Roberta Marziani²

Received: 23 January 2024 / Accepted: 12 November 2024 / Published online: 9 December 2024
© The Author(s) 2024

Abstract

We consider a variational problem modeling transition between flat and wrinkled region in a thin elastic sheet and identify the Γ -limit as the sheet thickness goes to 0, thus extending the previous work of the first author (Bella et al. in Arch Ration Mech Anal 218(1):163–217, 2015). The limiting problem is scalar and convex, but constrained and posed for measures. For the $\Gamma - \liminf$ inequality, we first pass to quadratic variables so that the constraint becomes linear, and then obtain the lower bound using Reshetnyak's theorem. The construction of the recovery sequence for the $\Gamma - \limsup$ inequality relies on mollification of quadratic variables and careful choice of multiple construction parameters. Eventually for the limiting problem, we show existence of a minimizer and equipartition of the energy for each frequency.

Keywords Microstructure · Wrinkling of elastic sheets · Nonconvex variation problem · Γ -convergence · Constrained minimization

Mathematics Subject Classification 74N15 · 49N

1 Introduction

This paper is about fine analysis of minimizers of a non-convex variational problem which describes wrinkling of thin elastic sheets.

Motivated by physical experiments with thin elastic sheets (Davidovitch et al. 2012, 2011; Géminard et al. 2004), the first author, in his PhD thesis (Bella 2012) (see also Bella and Kohn 2014), considered a specific variational problem describing deformations of a thin elastic sheet of thickness h and cross section of annular shape

Communicated by Peter Sternberg.

✉ Peter Bella
peter.bella@udo.edu

Roberta Marziani
roberta.marziani@unisi.it

¹ Technische Universität Dortmund Fakultät für Mathematik, 44227 Dortmund, Germany

² Dipartimento di Ingegneria dell' Informazione e Scienze Matematiche, 53100 Siena, Italy

$\Omega = \{x \in \mathbb{R}^2 : R_{\text{in}} < |x| < R_{\text{out}}\}$. If dead loads are applied on the inner and outer boundary of Ω in the radial direction with magnitude T_{in} pointing inwards and T_{out} pointing outwards, the membrane will stretch mainly in the radial direction. Moreover, if the inner loads T_{in} are much larger than the outer T_{out} , the material close to the inner boundary will move inwards and will occupy much less space than favored. To relieve the compression in the angular direction, the sheet becomes unstable and wrinkles out-of-plane (Fig. 1 left). Analysis of this wrinkled region is the main object of the present study.

In the reduced two-dimensional Kirchhoff–Love setting, the elastic energy, corresponding to a deformation of the cross section $v : \Omega \rightarrow \mathbb{R}^3$, consists of a membrane term, measuring stretching and compression of the sheet, a bending term, which penalizes curvature, and a boundary term representing the boundary loads. As a proxy for the energy, one can think of

$$E_h(v) = \int_{\Omega} W(\nabla v) \, dx + h^2 \int_{\Omega} \mathcal{Q}(\nabla v) \, dx + B(v), \quad (1.1)$$

where v is the normal to the deformed surface $v(\Omega)$, W is a nonlinear energy density, $\mathcal{Q}$ a quadratic function, and

$$B(v) = T_{\text{in}} \int_{|x|=R_{\text{in}}} v \cdot \frac{x}{R_{\text{in}}} \, dS - T_{\text{out}} \int_{|x|=R_{\text{out}}} v \cdot \frac{x}{R_{\text{out}}} \, dS$$

describing the dead loads applied in the radial direction on the inner and outer boundary. The membrane part is non-convex, possibly giving rise to oscillations. In contrast, the bending part is convex and of higher order, thus regularizing the problem. Since the bending resistance is related to the sheet thickness h , the magnitude of this contribution asymptotically vanishes in the limit $h \searrow 0$.

The physics approach to tackle these problems consists of a specific choice of an ansatz (guess) for the form of a minimizer. In other words, one restricts the analysis to a class of competitors having specific characteristics and looks for a minimizer of the energy within that class. On the other hand, the rigorous analytical approach does not make any assumptions on the form of a minimizer, i.e., the energy is minimized over all possible deformations. The problem in (1.1) being non-convex, hence possibly possessing many (local) minimizers or critical point, the first step is to understand the minimal value of the energy, with possibly learning some clues by which deformation is this minimal value, at least approximately, achieved.

Hence, we first try to identify the minimal value of the energy. Precisely, in the present situation, the goal is to understand its dependence on the (small) sheets thickness h . It turns out that the minimal value $\min_v E_h(v)$ consists of a leading zeroth-order term $\mathcal{E}_0$ (coming from the stretching of the sheet) plus a linear correction in h , which corresponds to the cost of *wrinkling* of the sheet (Bella 2012; Bella and Kohn 2014). More precisely, there exist two constants $0 < C_0 < C_1 < \infty$ such that for any thickness $0 < h < 1$ there holds

$$\mathcal{E}_0 + C_0 h \leq \min_v E_h(v) \leq \mathcal{E}_0 + C_1 h. \quad (1.2)$$




Fig. 1 Left: Elastic annular membrane stretched in the radial direction. The blue dotted curve represents the free boundary that separates the outer stretched region from the inner wrinkled one. Right: Rectangular piece of the membrane intersecting the free boundary curve (Color figure online)

The wrinkling serves as a mechanism to relieve compressive stresses in the inner region of the annulus (see Fig. 1), which are caused by specific geometrical effects. An alternative to wrinkling would be simply compression, which contributes to the membrane part at the order $O(1)$. Hence, in the case of small thickness (present situation) compression is much less energetically favorable ($O(1)$ vs $O(h^2)$) and thus not expected.

The identified linear scaling law (1.2) in h for the minimal value of the energy raised a lot of discussion among the physics community, having improved their ansatz-based prediction by a factor of $\log h$ (i.e., h in Bella and Kohn (2014) vs $h(|\log h| + 1)$ in Davidovitch et al. (2012)). It turns out that this discrepancy is related to a sub-optimal choice of the ansatz close to the interface between the wrinkled and the flat region. Moreover, the upper bound in (1.2) is achieved through a complex construction involving branching effects at the transition between the wrinkled and the flat region, a pattern which was not observed experimentally.

There might be several reasons for the discrepancy between the upper bound construction in Bella and Kohn (2014) and the experiments. Maybe the experimental sheet is not thin enough for the additional $\log h$ factor to lose the battle against (possibly large) prefactor C_1 in (1.2)? Maybe there are some additional effects which are neglected in the mathematical study? Or this discrepancy is related to dynamics leading to the deformation, i.e., one can be trapped in a local minima of the energy. Finally, due to non-convexity of the energy there is no reason for the construction achieving the upper bound (1.2) being unique. Actually, inspired by the study of wrinkling in twisted ribbons (Kohn and O'Brien 2018), in Bella and Kohn (<https://math.nyu.edu/~kohn/pcmi/annulus-problem-notes-for-pcmi.pdf>) a different fold-like construction achieving the optimal scaling without branching is sketched.

To understand which of possibly many different constructions is *energetically optimal* as well as which of the terms neglected in the force balance approach Davidovitch et al. (2012) should not be ignored, the first author considered a variational toy problem

modeling the transition region (Bella 2015) with the aim of better understanding the behavior of the minimizer in that region. Precisely, in Bella (2015) the author restricts the analysis to a rectangular piece of the elastic annulus (see Fig. 1), thus neglecting some terms due to circular geometry. In fact, wrinkles are caused by the compression in the angular direction, which is related to movement of the inner circles toward the center. To avoid the need to work in radial coordinates, for example to avoid the need to include that the circles with different radii have different lengths, in Bella (2015) the first author instead decided to phrase the problem in rectangular domain while prescribing a non-Euclidean metric together with periodic boundary conditions in the y -variable. Let us point out that the metric we prescribe is not coming from the change from radial to rectangular setting, but rather from the fact that the circles move inwards and are therefore compressed. In particular, already the rectangular model includes essential features of the problem, while being less technical to be analyzed compared to the original radial geometry. We believe that the present arguments can be used with minor modifications also in the original circular setting.

Then, working at the level of the energy, one considers the quantity $\frac{\min_v E_h(v) - \mathcal{E}_0}{h}$ (where now E_h is a simplified version of (1.2) coming from Föppl-von Kármán theory, and v is the displacement, see formula (2.1)), which is not only bounded away from 0 and ∞ (see (1.2)), but as $h \searrow 0$ it actually converges to some value σ (as proven in Bella (2015)). Even though the value σ is characterized as a limit of minima of simpler scalar and convex variational problems, it *does not* provide any information on the form of sequence of minimizers.

In that respect, the goal of this paper is to overcome this shortcoming by identifying the Γ -lim of $\frac{E_h - \mathcal{E}_0}{h}$ as $h \searrow 0$ combined with a compactness result. As usual, as a consequence we obtain convergence of minimizers u_h of E_h to a minimizer of the limiting problem, hence providing information on u_h , at least for $0 < h \ll 1$. Denoting by $\mathcal{F}_\infty$ the Γ -limit functional (see (2.9) below), it turns out that as expected from Bella (2015), $\mathcal{F}_\infty$ is scalar and convex, thus possibly much easier to analyze than the original E_h . Nevertheless, the study of minimizers of $\mathcal{F}_\infty$ is still far from trivial and we postpone it to a future work—except for some preliminary results collected in Sect. 6. As actually mentioned above, by learning the form of (the) minimizer of $\mathcal{F}_\infty$, hence also asymptotic form of minimizers of E_h , we hope to understand how is the transition between the planar and the wrinkled region precisely achieved, and which terms in the force balance cannot be neglected.

While analysis of the form of minimizer(s) of $\mathcal{F}_\infty$ will be pursued elsewhere, in the present article we directly show existence of a minimizer (which alternatively also follow from compactness of minimizing sequence and Γ -convergence) and equipartition between membrane and bending energies. While equipartition of the energy had been observed in several problems on energy-driven pattern formation with competing energetic parts, here we show much stronger statement: it actually holds for *every frequency* separately. This observation should later be important for the analysis of the form of minimizer (e.g., its asymptotic self-similarity), since it is usually easier to control one part of the energy than the other: here the bending part should be easier to handle since it depends on the values of the coefficients and not its derivative. As a

first implication of this fact, we show that any minimizer of $\mathcal{F}_\infty$ will not possess any long frequencies.

There are many areas of material science, most of them falling within a class of energy-driven pattern formation (Kohn 2007), where the idea to study energy scaling laws for variational problems turned out to be very fruitful. The common feature of these problems is the presence of a non-convex term in the energy, which is regularized by a higher-order term with a small prefactor. This small parameter (for now denoted ε) has different meanings: thickness in the case of elastic films, inverse Ginzburg–Landau parameter in the theory of superconductors, strength of the interfacial energy for models of shape memory alloys or micromagnetics, to name just few. As $\varepsilon \searrow 0$, the oscillations caused by the non-convexity are less penalized, giving the energy more freedom to form patterns/microstructure.

The first paper in this direction, in the context of shape memory alloys, is a seminal work of Kohn and Müller (1994), where they studied a toy problem to model the interface in the austenite–martensite phase transformation. They showed that the energy minimum scales like $\varepsilon^{2/3}$, which was in contrast with the scaling $\varepsilon^{1/2}$, widely accepted in the physics community. More precisely, the physics arguments were based on an ansatz of “one-dimensional” structure of minimizers, whereas Kohn and Müller used a branching construction to achieve lower energy. While they did not show the form of minimizers, they provided localized (in one direction) estimates on the energy distribution for the minimizer—thus providing hints on scales used for branching. Subsequently, Conti (2000) used an intricate upper bound construction to show localized energy bounds (in both directions), which in particular implies asymptotical self-similarity of the minimizer close to the interface. The analysis of the toy model was later generalized in several directions, for example analysis based on energy scalings laws for the cubic-to-tetragonal phase transformation—e.g., rigidity of the microstructure (Capella and Otto 2009, 2012) or study of the energy barrier for the nucleation in the bulk (Knüpfer et al. 2013) and at the boundary (Bella and Goldman 2015). In that respect it is worth to also mention recent works of Rüländ and Tribuzio (2023b, 2023a), where a novel use of Fourier Analysis allows to obtain sharp lower bounds on the energy on a more advanced model.

The work of Kohn and Müller initiated many developments in other areas of material science to study pattern formation driven by the energy minimization, for example in micromagnetics (Otto 2002; Brietzke and Knüpfer 2023; Ried and Román 2024), island growth on epitaxially strained films (Bella et al. 2015), diblock copolymers (Choksi 2001), optimal design (Kohn and Wirth 2016, 2014), superconductors (Serfaty 2015), dislocations in crystals (see, e.g., Conti et al. 2023; Garroni et al. 2021), fractures in solids (see, e.g., Bach et al. 2023, 2022, 2023; Marziani and Solombrino 2024) and phase separation (Marziani 2023). Picking one of them as an example, the Ginzburg–Landau model describes behavior of superconductors in different regimes of the applied magnetic field. While for extreme values of the magnetic field (very small or very large) there is only one (normal or superconducting) phase, for intermediate values of the field the mixed states consisting of many vortices are observed. There the leading order energy characterizes the number of vortices, and the next order in the energy describes interaction between them (see Serfaty 2015 for a survey,

Román et al. 2023 for analysis in three spatial dimensions, and Petrache and Serfaty 2017 for a similar work in the context of $2d$ Coulomb gases).

The models for wrinkling of thin elastic films have similar feature, with the leading order term in the energy expansion encoding the wrinkled regions while the next term in the energy expansion being related to the form (e.g., length scale) of wrinkling. The relevant physical object being a two-dimensional (thickened) surface in $\mathbb{R}^3$, the local energy expense of a deformation v is encoded using two principal values of a 3×2 matrix ∇v —heuristically, singular value greater or smaller than 1 corresponds to a tension or a compression, respectively. Wrinkling being an energetically efficient alternative to a compression, we expect it to appear in the case of (at least) one singular value being less than one.

A compressed elastic sheet can feel the compression in one (“tensile wrinkling”) or both directions (“compressive wrinkling”). A class of problems falling into the latter category for which the energy scaling laws were identified include, for instance, blistering/delamination problem (with Kohn and Nguyen 2013; Bedrossian and Kohn 2015; Bourne et al. 2016; Padilla-Garza 2020 or without Jin and Sternberg 2001; Ben Belgacem et al. 2002, 2000 substrate effects), crumpling of elastic sheets (Conti and Maggi 2008; Venkataramani 2004), or analysis of conical singularities in elastic sheets (Olbermann 2019, 2018). A common feature of this problem is degeneracy of the relaxed energy: the minimum of the relaxed energy $\mathcal{E}_0$ equals zero, and more importantly, it is achieved by many different minimizers, making the analysis of the next order expansion of the energy often difficult.

In contrast, tensile wrinkling problems usually have relaxed problem with unique minimizer, making the analysis of the next order term (which describes wrinkling) more accessible. The need for compression usually comes from the prescribed boundary conditions (as, for example, in the raft problem Brandman et al. 2013; Huang et al. 2010, twisted ribbon Kohn and O’Brien 2018, hanging drapes Vandeparre et al. 2011; Bella and Kohn 2017, or compressed cylinder Tobasco 2018), through prescribed incompatible strain (Bella and Kohn 2014; Maor and Shachar 2019) or curvature effects (Bella and Kohn 2017; Bella and Román 2023; Tobasco 2021).

The model we consider here is a mixture of the first and the second case, i.e., it is driven by both the boundary conditions and prescribed nontrivial metric (prestrain). The latter should mimic the need to “waste the length” in one direction, this need coming from geometric effects in our original motivation (Bella and Kohn 2014). More precisely, in Bella and Kohn (2014) an elastic annulus is stretched radially with stronger inner loads, forcing some of the concentric circles of material to move closer to the center. Pushing some circles into less space naturally forces compression or wrinkling out of plane, while the circles toward the outer boundary stay planar, and are actually stretched in the azimuthal direction. Phrased differently, the *excess length* of circles (i.e., the amount of material which needs to be “wasted”) is positive close to the inner boundary (compression) and negative on the outer edge (tension). As a continuous (smooth) function, it passes through 0, which is exactly the region which we analyze in this paper (blue dotted circle in Fig. 1 left). The excess length can be obtained as a minimizer of a one-dimensional variational problem (Bella and Kohn 2014), in particular is non-degenerate close to its 0—for simplicity we approximate it with a linear function (its first-order Taylor polynomial).

As we will see, it is crucial that the amount of arclength grows linearly in the distance from the free boundary (between the wrinkled and planar region) and not slower (e.g., quadratically)—the latter case is expected to be quite boring with the minimizer using only *one* frequency. In contrast, the present problem requires infinitely many frequencies, in particular near the transition higher and higher frequencies are needed.

The rest of this section will provide an overview of our results and organization of the paper. As in Bella (2015), we consider a specific thickness-dependent energy E_h (see (2.1) for its precise definition), a model problem describing transition between planar and wrinkled region in thin elastic sheet and are interested to understand structure of minimizers of the energies as $h \searrow 0$. We consider a thin elastic sheet of thickness h and cross section of rectangular shape $[-1, 1] \times [-1, 1] \subset \mathbb{R}^2$, which represents a piece of the elastic annulus depicted in Fig. 1 by a green region, and assume the sheet is i) stretched in the x -direction, and ii) stretched/compressed in the y -direction proportional to x (i.e., it is unstrained for $x = 0$, stretched in the y -direction in the left half and compressed in the right half of the domain). The stretching/compression in the y -direction is modeled via prescribed metric together with periodic boundary conditions at the top and bottom boundary.

To relax the compression in the region $\{x > 0\}$, we expect the sheet to wrinkle, with the length scale of wrinkles of order $h^{1/2}$ (Bella and Kohn 2014). In order to analyze the limit of $\frac{E_h - \mathcal{E}_0}{h}$ as $h \searrow 0$, we rescale the y -variable by $h^{-1/2}$ so that the wrinkles length scales stay of order 1, and the out-of-plane displacement has chance to converge to some limiting shape. A consequence of the rescaling is a change of the reference domain to the domain $[-1, 1] \times [-L, L]$ which is getting larger and larger as $L := h^{-1/2} \rightarrow \infty$, and so it is not clear how to perform the Γ -convergence analysis of the corresponding functionals. In order to avoid these complications we pass to the Fourier space. More precisely, we rewrite the energy using Fourier expansion in y , with L appearing through the summation set $\frac{\pi\mathbb{Z}}{L}$. Heuristically, as $L \rightarrow \infty$ the Fourier sum will turn into an integral; hence, there is a hope for a limiting functional to make sense.

An alternative to the rescaling in the y -variable would be to keep the domain fixed, and after passing to the Fourier space to perform a blow-up of the coefficients in the k -variable—this being necessary since we know that the wrinkles live on length scale $h^{1/2}$, i.e., in the limit $h \rightarrow 0$ the Fourier coefficients will concentrate on higher and higher frequencies. Naturally, this would be nothing else than what we actually do here—we will only commute the passage to Fourier and rescaling in the $y(k)$ -variable.

Another natural question is whether it would be possible, after rescaling in y and sending $L \rightarrow \infty$, to avoid the need to work on changing (growing) domains. The motivation for this is an expectation guided by our intuition that minimizers (low-energy configurations) should be relatively homogeneous in the sense that the wrinkles are spread evenly in the y -variable. This homogeneity being plausible, since the constraint measures the deformation along the whole interval, it would be very difficult, if not impossible, to relate the global behavior (constraint) with a local one (form of wrinkles on a fixed interval).

The rescaling was successfully pursued by the first author in Bella (2015), by observing i) the out-of-plane displacement u being the only relevant quantity to be monitored in this limit, and ii) for fixed (large) $L = h^{-1/2}$ the minimum of the excess energy $\frac{E_h - \mathcal{E}_0}{h}$ is well approximated by minimum of a *scalar, convex, and constrained* variational problem for u of the form

$$\begin{aligned} \mathcal{S}_L(u) := \int_0^1 \int_{-L}^L u_{,x}^2 + u_{,yy}^2 \, dx \, dy \quad \text{subject to} \\ \int_{-L}^L u_{,y}^2(x, y) \, dy = 2x + o(1) \quad \text{for a.e. } x \in (0, 1) \end{aligned} \quad (1.3)$$

Denoting by $a_k(x)$ the Fourier coefficients in y of $u(x, \cdot)$, we can rewrite

$$\mathcal{S}_L(u) = \int_0^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L}} (\dot{a}_k^2(x) + a_k^2(x)k^4) \, dx, \quad \text{and} \quad \int_{-L}^L u_{,y}^2(x, y) \, dy = \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a_k^2(x)k^2, \quad (1.4)$$

where “dot” denotes the derivative. The main achievement of Bella (2015) was to show that minima of $\mathcal{S}_L$ converges, and then to construct a recovery sequence for the original energy E_h , including construction of the in-plane displacement. Since the elastic energy E_h includes all second derivatives of u , and not only $u_{,yy}$ which appears in $\mathcal{S}_L$, regularity statement for the minimizers of $\mathcal{S}_L$ ’s played a crucial role for the construction of the recovery sequence.

The analysis of minima of $\mathcal{S}_L$ from Bella (2015) completely avoided the notion of convergence of minimizers, which needs to be an integral part of a Γ -convergence which we study here. To avoid the issue of nonlinear constraint, we use quadratic variables (i.e., monitoring $b_k := a_k^2$ instead of a_k), which turns the constraint into a linear one. The second term in the energy $\mathcal{S}_L$ becomes also linear, while the first term can be rewritten as $\frac{(\dot{b}_k)^2}{4b_k}$. One disadvantage of this approach is the L^1 -framework, which naturally leads the limit functional to be defined on the space of measures. However, the constraint provides a good pointwise control in x , in particular the limiting measure can be written as a product of dx and x -dependent measures in k . The lower bound argument (Proposition 4.2) is obtain using Reshetnyak Theorem.

The upper bound (construction of a recovery sequence) is much more tricky since it needs to be done for any “limiting” measure with finite energy, in contrast with Bella (2015), where it was done just for one (more regular) minimizer. The proof of the upper bound (Proposition 5.1) consists of several steps:

1. Given a limiting measure, to obtain a_k ’s we will “discretize” the measure in the k -variable (Lemma 5.3). Moreover, using smoothing of a_k ’s (more precisely of a_k^2), for which we need to extend the coefficients a_k from $[0, 1]$ via dilation into larger interval $[0, 1+]$, we get a good starting point for the construction.
2. Careful choice of the smoothing scale $\varepsilon(L)$ together with few other parameters allows for definition of the out-of-plane displacement (see Lemma 5.4), which is then basis for the construction of the in-plane displacement as well as estimates on

the excess energy (Proposition 5.1). While relatively complicated, the idea behind the definition of the in-plane displacement and the corresponding estimates are very similar to the one used in Bella (2015).

The paper is organized as follows: in the next chapter we provide a derivation of the energy, including the functional-analytical framework in form of measures with well-behaved distributional derivatives in x , as well as rewriting the energy to a form compatible also with this framework.

Afterward, we state the main result (Theorem 2.5), which consists of two, somehow independent, parts. The first is compactness result, coupling notion of convergence (Definition 2.3) and coercivity of the functionals. Said differently, it shows that the notion of convergence is a good one in the sense that any sequence (w_L, u_L) with equibounded “energies” $\mathcal{F}_L$ will possess a subsequence which converges with respect to that convergence. In particular, the notion of convergence is *weak* enough to follow from the boundedness of energies. The second part of the result (Γ -convergence) shows that at the same time the choice of convergence is *strong* enough to allow the passage to the limiting functional $\mathcal{F}_\infty$, including preservation the constraint in the limit $L \rightarrow \infty$.

After stating the main result, in the subsequent Chapter 3 we show how to disintegrate the limiting measures, and then continue with the arguments for the compactness (Proposition 4.1). With the help of Reshetnyak’s Theorem, we then show the lower bound (Proposition 4.2). The upper bound construction is content of Chapter 5. Eventually, in the last chapter we state and prove existence of minimizer (as a measure) for the limiting energy as well as pointwise (in k) equipartition of the energy for this minimizer (see Theorem 6.2). A finer analysis of this minimizer will be pursued in a future work of the authors.

2 Setting of the Problem and Main Results

We start by collecting some notation we will use throughout the paper.

Notation.

- (a) $a \lesssim b$ denotes $a \leq Cb$ for some constant $C > 0$;
- (b) χ_A denotes the characteristic function of the set A ;
- (c) $\mathcal{L}^1$ denotes the one-dimensional Lebesgue measure;
- (d) δ_k denotes the Dirac measure on $k \in \mathbb{R}$;
- (e) $\mathcal{M}_b(A)$ denotes the space of bounded Radon measures on A with $A \subset \mathbb{R}^2$ Borel measurable;
- (f) $\mathcal{M}_b^+(A)$ denotes the subspace of $\mathcal{M}_b(A)$ of positive bounded Radon measures;
- (g) For a function $f: A \subset \mathbb{R} \rightarrow \mathbb{R}$ we denote by $\dot{f}(x)$ and $\ddot{f}(x)$ the first and the second derivative, respectively;
- (h) For a function $u: A \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ we denote by $u_{\underbrace{x \dots x}_i \underbrace{y \dots y}_j}$ its partial derivative

$$D^{i+j}u(x, y) = \frac{d^j}{dy^j} \frac{d^i}{dx^i} u(x, y), \quad i, j \in \mathbb{N}, \quad 1 \leq i + j \leq 3;$$




Fig. 2 Domain with the boundary conditions

- (i) For a measure $\mu \in \mathcal{M}_b(A)$ we denote by $\mu_{,x}$ its distributional derivative with respect to the first variable;
- (j) For a measure $\mu \in \mathcal{M}_b(A)$ we denote by $|\mu| \in \mathcal{M}_b^+(A)$ its total variation;
- (k) For $\tilde{\mu} = (\mu_1, \mu_2) \in (\mathcal{M}_b(A))^2$ we analogously denote by $|\tilde{\mu}| \in \mathcal{M}_b^+(A)$ its total variation;
- (l) For $\mu_1 \in \mathcal{M}_b(A)$, $\mu_2 \in \mathcal{M}_b^+(A)$ we write $\mu_1 \ll \mu_2$ if μ_1 is absolute continuous with respect to μ_2 and we indicate by $\frac{d\mu_1}{d\mu_2} \in L^1(A, \mu_2)$ the associated density (Radon–Nikodým derivative);
- (m) $f * g(x)$ denotes the convolution between two functions f and g .

The Model. Let us now describe the model (energy) for the transition between the flat and wrinkled region, which the first author started analyzing in Bella (2015). Instead of considering the annular elastic sheet as in Bella and Kohn (2014), we consider only a rectangular piece (cut off from the sheet) near the transition region, represented by the domain $[-1, 1] \times [-1, 1] \subset \mathbb{R}^2$, in particular simplifying the problem by avoiding the need to work in the radial geometry. The annular sheet in Bella and Kohn (2014) is stretched in the radial direction and the concentric circles close to the transition region are stretched/compressed proportional to the distance from the free boundary. We will model the radial stretching by dead tension loads in the horizontal direction with magnitude $T = 1$, while the stretching/compression in the vertical direction will be modeled by prescribing a non-Euclidean metric of the form $dx + (1 + \delta x) dy$ for some $\delta > 0$.¹ Moreover, for the rectangle modeling part of the annulus, we prescribe periodic boundary conditions in the vertical direction (see Fig. 2). The choice of rectangular domain and the metric should prevent the arguments becoming too technical, but such that this simplified setting already includes the essential features of the problem. In particular, we believe that our arguments could be extended also to the original setting of an annular domain and metric with linear growth near the transition.

It is physically natural (Davidovitch et al. 2012) and mathematically convenient to use “small-slope” geometrically linear Föppl–von Kármán theory. In the membrane

¹ The parameter δ is exactly value of the derivative of the excess length at the transition.

part of the energy the in-plane displacement is represented via the *linear* strain, while the out-of-plane displacement is kept nonlinear (quadratic). The bending part is modeled by simply L^2 norm of the Hessian of the out-of-plane displacement instead of L^2 norm of the second fundamental form. Denoting by $w = (w_1, w_2)$ and u the in-plane and out-of-plane displacement, respectively, the elastic energy E_h (normalized per unit thickness) has the form

$$E_h(w, u) := \frac{1}{2} \int_{-1}^1 \int_{-1}^1 |e(w) + \frac{1}{2} \nabla u \otimes \nabla u - \delta x e_2 \otimes e_2|^2 dx dy + \frac{1}{2} \int_{-1}^1 \int_{-1}^1 h^2 |\nabla^2 u|^2 dx dy - \int_{-1}^1 (w_1(1, y) - w_1(-1, y)) dy. \quad (2.1)$$

Here $e(w) := (\nabla w + \nabla^T w)/2$ denotes the symmetric gradient of w and $\delta x e_2 \otimes e_2$ is the deviation of the prescribed metric from the Euclidean one. To simplify the notation, we will w.l.o.g. assume $\delta = 1$. Indeed, as will be seen from the subsequent arguments for both the lower and upper bounds, the limiting problem depends linearly on the value of $\delta > 0$,² and hence, this value can be simply scaled out. The third integral models the applied tensile dead loads in the horizontal direction. The factor $1/2$ in front of the elastic energy is chosen for convenience and can be changed to any factor using simple rescaling of w and u , at the expense of changing the metric prefactor as well. Finally, we assume the displacement (w, u) is 2-periodic in the second variable.

The behavior of E_h as $h \rightarrow 0$ at the leading order is well understood using relaxation techniques (Pipkin 1994) (also called tension-field theory in the mechanics community). Applied to E_h from (2.1), in the limit $h \rightarrow 0$ the bending term simply disappears, and the integrand in the membrane term gets relaxed to

$$(e(w) - x e_2 \otimes e_2)_+^2 := \min_{A \geq 0} |e(w) - x e_2 \otimes e_2 + A|^2,$$

where " $A \geq 0$ " stays for positive semi-definite 2×2 matrix. Hence, one can explicitly compute the (unique) minimizer of the relaxed energy ($w_2 = 0$ and $w_1 = x$) and its minimum $-2 + \frac{1}{3} = -\frac{5}{3} =: \mathcal{E}_0$.

From Bella and Kohn (2014) we know that the next term in the energy E_h scales linearly in h ; hence, the right quantity to look at is the rescaled *excess energy* $\frac{E_h - \mathcal{E}_0}{h}$. For $x > 0$ one expects that the sheet wrinkles out-of-plane in the y -direction, in order to offset $-x e_2 \otimes e_2$ with $u_{,yy}^2$. The linear scaling in h predicts $h^2 |\nabla^2 u|^2 \sim h$, in particular $u_{,yy}$ (its largest component) to be of order $h^{-1/2}$. As a consequence, the scale of wrinkles in the bulk should be reciprocal of this value, i.e., $h^{1/2}$. Not surprisingly, this is also the scale used in the upper bound construction in Bella and Kohn (2014).

In order to analyze the limiting form of the wrinkles as $h \rightarrow 0$, we rescale the y -variable by a factor $L := h^{-1/2}$, so that the characteristic length scale of wrinkles becomes 1. Precisely, after performing the change of variables

$$\hat{w}_1(x, y) := w_1(x, L^{-1}y), \quad \hat{w}_2(x, y) := L w_2(x, L^{-1}y), \quad \hat{u}(x, y) := L u(x, L^{-1}y),$$

² In the definition (2.8) the constraint simply becomes $2\delta x$.

the energy E_h becomes (see Bella 2015, page 630) for a straightforward algebraic manipulation)

$$\begin{aligned}\mathcal{E}_L(w, u) := & \int_{-L}^L \int_{-1}^1 \left(\left(w_{1,x} + \frac{u_{,x}^2}{2L^2} - 1 \right)^2 - 1 \right) dx dy \\ & + \int_{-L}^L \int_{-1}^1 \left(w_{2,y} + \frac{u_{,y}^2}{2} - x \right)^2 dx dy \\ & + \frac{1}{L^2} \int_{-L}^L \int_{-1}^1 \left(L^2 w_{1,y} + w_{2,x} + u_{,x} u_{,y} \right)^2 dx dy \\ & + \frac{1}{L^2} \int_{-L}^L \int_{-1}^1 (u_{,x}^2 + u_{,yy}^2) dx dy\end{aligned}\quad (2.2)$$

$$+ \frac{1}{L^4} \int_{-L}^L \int_{-1}^1 \left(2u_{,xy}^2 + \frac{1}{L^2} u_{,xx}^2 \right) dx dy. \quad (2.3)$$

Thus, the functional is defined as $\mathcal{E}_L: \mathcal{A}_L^{\text{in}} \times \mathcal{A}_L^{\text{out}} \rightarrow [0, +\infty]$, where the function spaces describing admissible deformations have the form

$$\begin{aligned}\mathcal{A}_L^{\text{in}} := & \left\{ w = (w_1, w_2) \in W_{\text{loc}}^{1,2}((-1, 1) \times \mathbb{R}; \mathbb{R}^2): \right. \\ & \left. w(x, \cdot) \text{ is } 2L - \text{periodic } \forall x \in (-1, 1) \right\},\end{aligned}\quad (2.4)$$

$$\mathcal{A}_L^{\text{out}} := \left\{ u \in W_{\text{loc}}^{2,2}((-1, 1) \times \mathbb{R}): u(x, \cdot) \text{ is } 2L\text{-periodic } \forall x \in (-1, 1) \right\}. \quad (2.5)$$

Furthermore, $\frac{E_h - \mathcal{E}_0}{h}$ turns into $\mathcal{F}_L: \mathcal{A}_L^{\text{in}} \times \mathcal{A}_L^{\text{out}} \rightarrow \mathbb{R}$ defined as

$$\mathcal{F}_L(w, u) := L^2(\mathcal{E}_L(w, u) - \mathcal{E}_0), \quad (2.6)$$

where $\mathcal{E}_0 = -\frac{5}{3}$ is as above the minimum of the relaxed energy, so that

$$\begin{aligned}\mathcal{F}_L(w, u) = & L^2 \int_{-L}^L \int_{-1}^1 \left(w_{1,x} + \frac{u_{,x}^2}{2L^2} - 1 \right)^2 dx dy - \frac{L^2}{3} \\ & + L^2 \int_{-L}^L \int_{-1}^1 \left(w_{2,y} + \frac{u_{,y}^2}{2} - x \right)^2 dx dy \\ & + \int_{-L}^L \int_{-1}^1 \left(L^2 w_{1,y} + w_{2,x} + u_{,x} u_{,y} \right)^2 dx dy \\ & + \int_{-L}^L \int_{-1}^1 (u_{,x}^2 + u_{,yy}^2) dx dy \\ & + \frac{1}{L^2} \int_{-L}^L \int_{-1}^1 \left(2u_{,xy}^2 + \frac{1}{L^2} u_{,xx}^2 \right) dx dy.\end{aligned}$$

Before we rigorously proceed further, let us discuss heuristically the form of functional $\mathcal{F}_L$ and its implications. Most of the terms in the energy are of quadratic nature, and since in addition we are dealing with oscillatory objects defined on longer and longer intervals, it is natural to look at the problem in the Fourier space.

Expecting the limit of $\mathcal{F}_L$ to exist (in particular having the minimizing sequence bounded as $L \rightarrow \infty$), both integrals on the first line need to (quickly) converge to 0. The first integral can easily achieve that by simply choosing $w_1 \sim x + o(L^{-1})$ and $u_{,x}$ not too big, the smallness of the second integral (after integration in y and using periodicity of w) implies the constraint $\int_{-L}^L u_{,y}^2 dy = 2x + o(L^{-1})$.

In order to have continuity of the constraint in the limit $L \rightarrow \infty$, and also for other reasons which will be apparent later, we will work with squares of the Fourier coefficients and suitably defined measures as primary objects of studies. In the following we denote by $k \in \mathbb{R}$ the variable corresponding to the Fourier transform in the y -variable. Moreover, we use the same notation to denote the second variable when working with measures.

Definition 2.1 (Measures μ^L and $\mu_{,x}^L$) Let $u \in \mathcal{A}_L^{\text{out}}$. We denote by $\mu^L(u) \in \mathcal{M}_b^+((-1, 1) \times \mathbb{R})$ the measure given by

$$\mu^L(u) := \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a(x, k) \mathbb{L}^1 \llcorner (-1, 1) \times \delta_k,$$

with

$$a(x, k) := k^2 a_k^2(x),$$

and $a_k \in W^{2,2}(-1, 1)$ being the k -th Fourier coefficient of $u(x, \cdot)$ for all $x \in (-1, 1)$, that is

$$a_k(x) := \begin{cases} \sqrt{2} \int_{-L}^L u(x, y) \sin(ky) dy & k \in \frac{\pi\mathbb{Z}}{L}, k > 0, \\ \sqrt{2} \int_{-L}^L u(x, y) \cos(ky) dy & k \in \frac{\pi\mathbb{Z}}{L}, k < 0, \\ \int_{-L}^L u(x, y) dy & k = 0. \end{cases} \quad (2.7)$$

Moreover, we denote by $\mu_{,x}^L(u)$ the distributional x -derivative of $\mu^L(u)$.

Remark 2.2 (i) The distributional x -derivative of a measure $\mu \in \mathcal{M}_b^+((-1, 1) \times \mathbb{R})$ is defined as follows: for all $\varphi \in C_c^\infty((-1, 1) \times \mathbb{R})$ we have

$$\langle \mu_{,x}, \varphi \rangle := - \int_{(-1, 1) \times \mathbb{R}} \varphi_{,x} d\mu.$$

Moreover, by a density argument $\mu_{,x}$ can be extended to functions $\varphi(x, k) = \phi(x)\chi_A(k)$ with $\phi \in C_c^\infty(-1, 1)$ and $A \subset \mathbb{R}$ bounded and measurable as

$$\langle \mu_{,x}, \phi(x)\chi_A(k) \rangle := - \int_{(-1,1) \times A} \dot{\phi}(x) \, d\mu;$$

(ii) Let $\mu \in \mathcal{M}_b^+((-1, 1) \times \mathbb{R})$ be of the form

$$\mu = \sum_{k \in K} a(x, k) \mathcal{L}^1 \llcorner (-1, 1) \times \delta_k,$$

with $K \subset \mathbb{R}$ countable and $a(\cdot, k) \in W^{1,1}(-1, 1)$ for all $k \in K$. Then

$$\mu_{,x} = \sum_{k \in K} a_{,x}(x, k) \mathcal{L}^1 \llcorner (-1, 1) \times \delta_k.$$

Moreover, as $a_{,x}(\cdot, k) = 0$ a.e. in $\{x \in (-1, 1) : a(x, k) = 0\}$ it follows $\mu_{,x} \in \mathcal{M}((-1, 1) \times \mathbb{R})$ and $\mu_{,x} \ll \mu$.

Definition 2.3 (Convergence) For $L > 0$ let $(w^L, u^L) \in \mathcal{A}_L^{\text{in}} \times \mathcal{A}_L^{\text{out}}$. We say a sequence (w^L, u^L) converges as $L \rightarrow \infty$ to $\mu \in \mathcal{M}_b^+((-1, 1) \times \mathbb{R})$, if $(\mu^L(u^L), \mu_{,x}^L(u^L))$ weakly-* converge to $(\mu, \mu_{,x})$.

We introduce the class of measures

$$\begin{aligned} \mathcal{M}_\infty := & \left\{ \mu \in \mathcal{M}_b^+((-1, 1) \times \mathbb{R}) : \mu((-1, 0] \times \mathbb{R}) = 0, \mu_{,x} \in \mathcal{M}_b((-1, 1) \times \mathbb{R}), \right. \\ & \left. \mu_{,x} \ll \mu, \int_{(0,1) \times \mathbb{R}} \phi(x) \, d\mu(x, k) = \int_0^1 2x\phi(x) \, dx \quad \forall \phi \in C_c^\infty(0, 1) \right\}, \end{aligned} \quad (2.8)$$

and the functional $\mathcal{F}_\infty : \mathcal{M}_\infty \rightarrow [0, +\infty]$

$$\mathcal{F}_\infty(\mu) = \int_{(0,1) \times \mathbb{R}} \left[k^2 + \frac{1}{4k^2} \left(\frac{d\mu_{,x}}{d\mu}(x, k) \right)^2 \right] d\mu(x, k). \quad (2.9)$$

Remark 2.4 (i) Above $\frac{d\mu_{,x}}{d\mu}$ denotes the Radon–Nikodym derivative, existence of which follows from absolute continuity of $\mu_{,x}$ w.r.t. μ . Moreover, if μ_L from Definition 2.1 is supported in $(0, 1] \times \mathbb{R}$, then

$$\mathcal{F}_\infty(\mu_L) = \sum_{k \in \frac{\pi\mathbb{Z}}{L}} \int_0^1 (\dot{a}_k^2(x) + a_k^2(x)k^4) \, dx,$$

i.e., via Plancherel equality (see equation (3.3)) it is equal to S_L from (1.3). We farther observe that (1.4) combined with the above heuristic expectation

$\int_{-L}^L u_{,y}^2 dy = 2x + o(L^{-1})$ imply in the limit $L \rightarrow \infty$ the integral constraint in (2.8).

(ii) When convenient we will identify the class $\mathcal{M}_\infty$ with the class of measures

$$\left\{ \mu \in \mathcal{M}_b^+((0, 1) \times \mathbb{R}) : \mu_{,x} \in \mathcal{M}_b((0, 1) \times \mathbb{R}), \mu_{,x} \ll \mu, \right. \\ \left. \int_{(0,1) \times \mathbb{R}} \phi(x) d\mu(x, k) = \int_0^1 2x \phi(x) dx \quad \forall \phi \in C_c^\infty(0, 1) \right\}; \quad (2.10)$$

(iii) For later convenience we observe that $\mathcal{F}_\infty$ can be rewritten as follows

$$\mathcal{F}_\infty(\mu) = \int_{(0,1) \times \mathbb{R}} k^2 d\mu + \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu}{d|\tilde{\mu}|} \right)^{-1} \left(\frac{d\mu_{,x}}{d|\tilde{\mu}|} \right)^2 d|\tilde{\mu}|, \quad (2.11)$$

where $\tilde{\mu} = (\mu, \mu_{,x})$ and $|\tilde{\mu}|$ denote its total variation. Indeed, since $\mu_{,x} \ll \mu \ll |\tilde{\mu}|$, we have

$$\frac{d\mu_{,x}}{d|\tilde{\mu}|} = \frac{d\mu_{,x}}{d\mu} \frac{d\mu}{d|\tilde{\mu}|},$$

from which we deduce

$$\begin{aligned} \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu_{,x}}{d\mu} \right)^2 d\mu &= \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu_{,x}}{d\mu} \right)^2 \frac{d\mu}{d|\tilde{\mu}|} d|\tilde{\mu}| \\ &= \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu}{d|\tilde{\mu}|} \right)^{-2} \left(\frac{d\mu_{,x}}{d|\tilde{\mu}|} \right)^2 \frac{d\mu}{d|\tilde{\mu}|} d|\tilde{\mu}| \\ &= \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu}{d|\tilde{\mu}|} \right)^{-1} \left(\frac{d\mu_{,x}}{d|\tilde{\mu}|} \right)^2 d|\tilde{\mu}|. \end{aligned}$$

We are now ready to state our main result.

Theorem 2.5 *Let $\mathcal{F}_L$ and $\mathcal{F}_\infty$ be as in (2.6) and (2.9) respectively. Then the following holds:*

- (Compactness). For $L > 0$ let $(w^L, u^L) \in \mathcal{A}_L^{\text{in}} \times \mathcal{A}_L^{\text{out}}$ be such that $\sup_L \mathcal{F}_L(w^L, u^L) < +\infty$. Then there exists a subsequence (not relabeled) and $\mu \in \mathcal{M}_\infty$ such that (w^L, u^L) converges as $L \rightarrow +\infty$ in the sense of Definition 2.3 to μ .
- (Γ -convergence). As $L \rightarrow +\infty$ the functionals $\mathcal{F}_L$ Γ -converge, with respect to the convergence in Definition 2.3, to the functional $\mathcal{F}_\infty$.

3 Preliminaries

Let $u \in \mathcal{A}_L^{\text{out}}$ and let $a_k(x)$ be defined as in (2.7). Then we have

$$\begin{aligned} u(x, y) &= a_0(x) + \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k > 0} a_k(x) \sqrt{2} \sin(ky) + \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k < 0} a_k(x) \sqrt{2} \cos(ky) \\ &= a_0(x) + \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k > 0} \text{sign}(a_k(x)) \frac{\sqrt{a(x, k)}}{k} \sqrt{2} \sin(ky) \\ &\quad + \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k < 0} \text{sign}(a_k(x)) \frac{\sqrt{a(x, k)}}{-k} \sqrt{2} \cos(ky). \end{aligned} \quad (3.1)$$

Then Plancherel equality yields

$$\int_{-L}^L u^2 dy = a_0^2(x) + \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} a_k^2(x) = a_0^2(x) + \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \frac{a(x, k)}{k^2}. \quad (3.2)$$

The same holds for partial derivatives of u , that is

$$\begin{aligned} \int_{-L}^L (D^\alpha u)^2 dy &= (D^\alpha a_0(x))^2 + \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \left(\frac{d^{\alpha_1}}{dx^{\alpha_1}} a_k(x) k^{\alpha_2} \right)^2 \\ &= (D^\alpha a_0(x))^2 + \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \left(\frac{\partial^{\alpha_1}}{\partial x^{\alpha_1}} (\sqrt{a(x, k)}) k^{\alpha_2-1} \right)^2, \end{aligned} \quad (3.3)$$

with $\alpha = (\alpha_1, \alpha_2)$ multi-index with $|\alpha| \leq 2$. In case u has higher regularity, i.e., $u \in W^{k,2}((-1, 1) \times \mathbb{R})$ with $k > 2$, then the same applies for the higher derivatives, i.e., for $|\alpha| \leq k$. For later convenience we also note that

$$\frac{\partial}{\partial x} (\sqrt{a(x, k)}) = \frac{a_{,x}(x, k)}{2\sqrt{a(x, k)}}, \quad \frac{\partial^2}{\partial x^2} (\sqrt{a(x, k)}) = \frac{a_{,xx}(x, k)}{2\sqrt{a(x, k)}} - \frac{(a_{,x}(x, k))^2}{4\sqrt{a^3(x, k)}}. \quad (3.4)$$

We now recall the definition of disintegration of measures only in a specific case that we will be used throughout the paper, and we refer to Ambrosio et al. (2000) for a complete treatment of the subject.

Definition 3.1 [Disintegration of measures in the x -variable] Let $I \subset \mathbb{R}$ be an interval and let $\mu \in \mathcal{M}_b(I \times \mathbb{R})$. We say that the family

$$(\nu_x, g(x))_{x \in I} \subset \mathcal{M}_b(\mathbb{R}) \times \mathbb{R}$$

is a disintegration of μ (in the x -variable) if $x \mapsto \nu_x$ is Lebesgue measurable, $|\nu_x|(\mathbb{R}) = 1$ for every $x \in I$, $g \in L^1(I)$, and

$$\int_{I \times \mathbb{R}} f(x, k) d\mu = \int_I \int_{\mathbb{R}} f(x, k) d\nu_x(k) g(x) dx, \quad (3.5)$$

for every $f \in L^1(I \times \mathbb{R}; |\mu|)$.

Formally it simply means $d\mu(x, k) = d\nu_x(k)g(x) dx$.

Lemma 3.2 *Let $I \subset \mathbb{R}$ be an interval and let $\mu \in \mathcal{M}_b^+(I \times \mathbb{R})$. Then*

$$\int_{I \times \mathbb{R}} \phi(x) d\mu = \int_I g(x)\phi(x) dx \quad \forall \phi \in C_c^\infty(I), \quad (3.6)$$

for some nonnegative $g \in L^1(I)$, if and only if there exists $x \mapsto \nu_x \in \mathcal{M}_b^+(\mathbb{R})$ Lebesgue measurable such that $(\nu_x, g(x))_{x \in I}$ is a disintegration of μ .

Proof Let $(\nu_x, g(x))_{x \in I} \subset \mathcal{M}_b^+(\mathbb{R}) \times \mathbb{R}^+$ be a disintegration of μ . Then (3.5) holds with $f(x, k) = \phi(x) \in C_c^\infty(I)$ and since $|\nu_x|(\mathbb{R}) = \nu_x(\mathbb{R}) = 1$ we readily deduce (3.6).

Assume instead that (3.6) holds true. Let $\pi_1: I \times \mathbb{R} \rightarrow I$ be the canonical projection and let $(\pi_1)_\# \mu \in \mathcal{M}_b^+(I)$ be the push-forward of μ with respect to π_1 . By the Disintegration Theorem (cf. Ambrosio et al. 2000, Theorem 2.28) there exists $x \mapsto \nu_x \in \mathcal{M}_b^+(\mathbb{R})$ measurable with $\nu_x(\mathbb{R}) = 1$ such that

$$\int_{I \times \mathbb{R}} f(x, k) d\mu(x, k) = \int_I \int_{\mathbb{R}} f(x, k) d\nu_x(k) d(\pi_1)_\# \mu(x),$$

for all $f \in L^1(I \times \mathbb{R}; \mu)$. On the other hand (3.6) implies that $(\pi_1)_\# \mu(x) = g(x)\mathbb{L}^1 \llcorner I$ and therefore $(\nu_x, g(x))_{x \in I}$ is a disintegration of μ . $\square$

Corollary 3.3 *(Disintegration of $\mu \in \mathcal{M}_\infty$ in the x -variable) Let $\mu \in \mathcal{M}_\infty$. Then there exists $x \mapsto \nu_x \in \mathcal{M}_b^+(\mathbb{R})$ measurable such that $(\nu_x, 2x)_{x \in (0,1)}$ is a disintegration of μ .*

Proof The proof follows by Lemma 3.2 and from the fact that

$$\int_{(0,1) \times \mathbb{R}} \phi(x) d\mu = \int_0^1 2x\phi(x) dx \quad \forall \phi \in C_c^\infty(0, 1). \quad (3.7)$$

$\square$

4 Compactness and Lower Bound

In this section we prove compactness and the $\Gamma - \lim \inf$ inequality.

Proposition 4.1 (Compactness) *Let for $L > 0$ be $(w^L, u^L) \in \mathcal{A}_L^{\text{in}} \times \mathcal{A}_L^{\text{out}}$ such that $\sup_L \mathcal{F}_L(w^L, u^L) < +\infty$. Then there exist a, not relabeled, subsequence and $\mu \in \mathcal{M}_\infty$ such that (w^L, u^L) converges to μ , as $L \rightarrow +\infty$, in the sense of Definition 2.3.*

Proof Let (w^L, u^L) be as in the statement. Let $\mu^L := \mu^L(u^L)$ and $\mu_{,x}^L := \mu_{,x}^L(u^L)$ be defined accordingly to Definition 2.1, i.e., there exist $a^L(x, k)$ such that $a(\cdot, k) \in W^{1,1}(-1, 1)$ and

$$\mu^L = \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) \mathbb{L}^1 \llcorner (-1, 1) \times \delta_k, \quad \mu_{,x}^L = \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a_{,x}^L(x, k) \mathbb{L}^1 \llcorner (-1, 1) \times \delta_k.$$

Step 1: we show that there exists $\mu \in \mathcal{M}_b^+((-1, 1) \times \mathbb{R})$ with $\mu_{,x} \in \mathcal{M}_b((-1, 1) \times \mathbb{R})$ and such that $(\mu^L, \mu_{,x}^L) \xrightarrow{*} (\mu, \mu_{,x})$. To this aim, we observe that by taking $0 < C_0 := \sup_L \mathcal{F}_L(w^L, u^L) < +\infty$ we have

$$\mathcal{F}_L(w^L, u^L) \leq C_0,$$

so that in particular

$$\begin{aligned} C_0 \geq \mathcal{F}_L(w^L, u^L) &\geq L^2 \int_{-L}^L \int_{-1}^1 \left(w_{2,y}^L + \frac{(u_{,y}^L)^2}{2} - x \right)^2 dx dy - \frac{L^2}{3} \\ &\quad + \int_{-L}^L \int_{-1}^1 (u_{,x}^L)^2 + (u_{,yy}^L)^2 dx dy. \end{aligned} \quad (4.1)$$

By Fubini's theorem, Jensen's inequality and the fact that $w^L(x, \cdot)$ is $2L$ -periodic we get

$$\begin{aligned} &L^2 \int_{-L}^L \int_{-1}^1 \left(w_{2,y}^L + \frac{(u_{,y}^L)^2}{2} - x \right)^2 dx dy \\ &\geq L^2 \int_{-1}^1 \left(\int_{-L}^L \left(w_{2,y}^L + \frac{(u_{,y}^L)^2}{2} - x \right) dy \right)^2 dx \\ &= L^2 \int_0^1 \left(\int_{-L}^L \frac{(u_{,y}^L)^2}{2} dy - x \right)^2 dx \\ &\quad + L^2 \int_{-1}^0 \left(\int_{-L}^L \frac{(u_{,y}^L)^2}{2} dy - x \right)^2 dx \\ &\geq L^2 \int_0^1 \left(\int_{-L}^L \frac{(u_{,y}^L)^2}{2} dy - x \right)^2 dx + L^2 \\ &\quad \int_{-1}^0 \left(\int_{-L}^L \frac{(u_{,y}^L)^2}{2} dy \right)^2 dx + L^2 \int_{-1}^0 x^2 dx, \end{aligned} \quad (4.2)$$

where the last inequality follows by using that $(a + b)^2 \geq a^2 + b^2$ provided that $ab > 0$ with $a = \frac{1}{2} \int_{-L}^L (u_{,x}^L)^2 dy$ and $b = -x$ for $x \in (-1, 0)$. Combining (4.1) with (4.2) and using that $\int_{-1}^0 x^2 dx = \frac{1}{3}$, we find

$$\begin{aligned} C_0 \geq \mathcal{F}_L(w^L, u^L) &\geq L^2 \int_0^1 \left(\int_{-L}^L \frac{(u_{,y}^L)^2}{2} dy - x \right)^2 dx \\ &\quad + L^2 \int_{-1}^0 \left(\int_{-L}^L \frac{(u_{,y}^L)^2}{2} dy \right)^2 dx \\ &\quad + \int_{-L}^L \int_{-1}^1 (u_{,x}^L)^2 + (u_{,yy}^L)^2 dx dy. \end{aligned} \quad (4.3)$$

Thus, from (3.3) it follows

$$\begin{aligned} \frac{C_0}{L^2} &\geq \int_0^1 \left(\int_{-L}^L \frac{(u_{,y}^L)^2}{2} dy - x \right)^2 dx = \int_0^1 \left(\frac{1}{2} \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) - x \right)^2 dx \\ &\geq C \left(\int_0^1 \left(\frac{1}{2} \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) \right)^2 dx - \frac{1}{3} \right), \end{aligned} \quad (4.4)$$

and

$$\frac{C_0}{L^2} \geq \int_{-1}^0 \left(\int_{-L}^L \frac{(u_{,y}^L)^2}{2} dy \right)^2 dx = \int_{-1}^0 \left(\frac{1}{2} \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) \right)^2 dx. \quad (4.5)$$

Hence, we obtain

$$|\mu^L|(((-1, 1) \times \mathbb{R})) = \mu^L(((-1, 1) \times \mathbb{R})) = \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(k, x) dx \leq C,$$

from which we deduce the existence of a (not relabeled) subsequence and $\mu \in \mathcal{M}_b^+(((-1, 1) \times \mathbb{R}))$ such that $\mu^L \xrightarrow{*} \mu$. In addition, (4.3) together with (3.3) and (3.4) yield

$$\begin{aligned} C &\geq \int_{-1}^1 \int_{-L}^L (u_{,x}^L)^2 + (u_{,yy}^L)^2 dy dx \\ &\geq \int_{-1}^1 \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) k^2 + \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \frac{1}{4k^2} \frac{(a_{,x}^L(x, k))^2}{a^L(x, k)} \right) dx \\ &\geq \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L}} |a_{,x}^L(x, k)| dx = |\mu_{,x}^L|(((-1, 1) \times \mathbb{R})), \end{aligned} \quad (4.6)$$

where the last inequality follows by Young's inequality. Hence, up to subsequence, we may deduce that there exists $\tilde{\mu} \in \mathcal{M}_b(\times(-1, 1) \times \mathbb{R})$ such that $\mu_{,x}^L \xrightarrow{*} \tilde{\mu}$. Moreover, given any $\varphi \in C_c^\infty((-1, 1) \times \mathbb{R})$, it holds

$$\begin{aligned} \int_{(-1,1) \times \mathbb{R}} \varphi \, d\tilde{\mu} &= \lim_{L \rightarrow +\infty} \int_{(-1,1) \times \mathbb{R}} \varphi \, d\mu_{,x}^L = - \lim_{L \rightarrow +\infty} \int_{(-1,1) \times \mathbb{R}} \varphi_{,x} \, d\mu^L \\ &= - \int_{(-1,1) \times \mathbb{R}} \varphi_{,x} \, d\mu, \end{aligned}$$

which in turn implies $\tilde{\mu} = \mu_{,x}$.

Step 2: we show that $\mu_{,x} \ll \mu$. By Remark 2.2 (ii) we have that $\mu_{,x}^L \ll \mu^L$. Now let $N \in \mathbb{N}$ be fixed and let $\mu_N^L := \mu^L \llcorner (-1, 1) \times (-N, N)$ and $\mu_N := \mu \llcorner (-1, 1) \times (-N, N)$. Then the following properties hold:

$$\begin{aligned} \mu_{N,x}^L &:= \mu_{,x}^L \llcorner (-1, 1) \times (-N, N), \quad \mu_{N,x} := \mu_{,x} \llcorner (-1, 1) \times (-N, N), \\ \mu_{N,x}^L &\ll \mu_N^L, \quad (\mu_N^L, \mu_{N,x}^L) \xrightarrow{*} (\mu_N, \mu_{N,x}), \end{aligned} \quad (4.7)$$

and

$$\frac{d\mu_{N,x}^L}{d\mu_N^L}(x, k) = \frac{d\mu_{,x}^L}{d\mu^L}(x, k) \llcorner (-1, 1) \times (-N, N).$$

Moreover, recalling the definition of μ^L and (4.6) we have

$$\begin{aligned} &\int_{(-1,1) \times (-N,N)} \frac{1}{4N^2} \left(\frac{d\mu_{N,x}^L}{d\mu_N^L}(x, k) \right)^2 d\mu_N^L \\ &\leq \int_{(-1,1) \times (-N,N)} \frac{1}{4k^2} \left(\frac{d\mu_{,x}^L}{d\mu^L}(x, k) \right)^2 d\mu^L \\ &\leq \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \frac{1}{4k^2} \frac{(a_{,x}^L(x, k))^2}{a^L(x, k)} dx \leq C. \end{aligned} \quad (4.8)$$

From (4.7), (4.8) and (Ambrosio et al. 2000), Example 2.36 pg. 67, and discussion at pg. 66) we deduce that $\mu_{N,x} \ll \mu_N$ for every $N \in \mathbb{N}$ and hence $\mu_{,x} \ll \mu$.

Step 3: we show that $\mu \in \mathcal{M}_b^+(\mathbb{R})$, that is, $\mu((-\infty, 0] \times \mathbb{R}) = 0$, and that

$$\int_{(0,1) \times \mathbb{R}} \phi(x) \, d\mu = \int_0^1 2x\phi(x) \, dx, \quad (4.9)$$

for all $\phi \in C_c^\infty((0, 1))$. To this purpose for fixed $\delta \in (0, 1)$ by (4.3), we have

$$\begin{aligned} \mu^L((-1, \delta) \times \mathbb{R}) &= \int_{-1}^{\delta} \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) \, dx \\ &\leq C \int_{-1}^0 \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) \right)^2 \, dx \\ &\quad + C \int_0^{\delta} \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) - x \right)^2 \, dx + C \int_0^{\delta} x^2 \, dx \\ &\leq \frac{C}{L^2} + C\delta^3. \end{aligned}$$

This together with the lower semi-continuity with respect to the weak* convergence gives

$$\mu((-1, 0] \times \mathbb{R}) \leq \mu((-1, \delta) \times \mathbb{R}) \leq \liminf_{L \rightarrow \infty} \mu^L((-1, \delta) \times \mathbb{R}) \leq C\delta^3.$$

By sending $\delta \rightarrow 0$ we deduce $\mu((-1, 0] \times \mathbb{R}) = 0$. It remains to show (4.9). Given $\phi \in C_c^\infty(0, 1)$ it holds

$$\int_{(0,1) \times \mathbb{R}} \phi(x) \, d\mu^L = \int_0^1 \phi(x) \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) - 2x \right) \, dx + \int_0^1 2x\phi(x) \, dx.$$

From (4.4) it follows that

$$\begin{aligned} \int_0^1 |\phi(x)| \left| \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) - 2x \right| \, dx &\leq C \|\phi\|_\infty \left(\int_0^1 \left(\frac{1}{2} \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) - x \right)^2 \, dx \right)^{1/2} \\ &\leq \frac{C}{L} \rightarrow 0, \end{aligned}$$

as $L \rightarrow +\infty$, so that

$$\lim_{L \rightarrow +\infty} \int_{(0,1) \times \mathbb{R}} \phi(x) \, d\mu^L = \int_0^1 2x\phi(x) \, dx. \quad (4.10)$$

Next we fix $R \geq 1$ and take $\psi_R \in C_c^\infty(\mathbb{R})$ such that $0 \leq \psi_R \leq 1$, $\psi_R(k) \equiv 1$ if $|k| < R$ and $\psi_R(k) \equiv 0$ if $|k| > R + 1$. We have

$$\int_{(0,1) \times \mathbb{R}} \phi(x) \, d\mu^L = \int_{(0,1) \times \mathbb{R}} \phi(x) \psi_R(k) \, d\mu^L + \int_{(0,1) \times \mathbb{R}} \phi(x) (1 - \psi_R(k)) \, d\mu^L. \quad (4.11)$$

The weak* convergence yields

$$\lim_{L \rightarrow +\infty} \int_{(0,1) \times \mathbb{R}} \phi(x) \psi_R(k) \, d\mu^L = \int_{(0,1) \times \mathbb{R}} \phi(x) \psi_R(k) \, d\mu,$$

whereas for the second term on the right-hand side of (4.11) we get

$$\begin{aligned} \int_{(0,1) \times \mathbb{R}} |\phi(x)(1 - \psi_R(k))| \, d\mu^L &\leq \int_0^1 |\phi(x)| \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L}, |k| > R} a^L(x, k) \right) dx \\ &\leq \frac{\|\phi\|_\infty}{R^2} \int_0^1 \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L}, |k| > R} a^L(x, k) k^2 \right) dx \\ &\leq \frac{\|\phi\|_\infty}{R^2} \int_0^1 \int_{-L}^L (u_{,yy})^2 \, dy \, dx \leq \frac{C}{R^2}, \end{aligned}$$

where the last two inequalities follow from (3.3) and (4.3). Thus, passing to the limit as $L \rightarrow +\infty$ in (4.11), we obtain

$$\begin{aligned} \int_{(0,1) \times \mathbb{R}} \phi(x) \psi_R(k) \, d\mu - \frac{C}{R^2} &\leq \lim_{L \rightarrow +\infty} \int_{(0,1) \times \mathbb{R}} \phi(x) \, d\mu^L \\ &\leq \int_{(0,1) \times \mathbb{R}} \phi(x) \psi_R(k) \, d\mu + \frac{C}{R^2}. \end{aligned}$$

Eventually by letting $R \rightarrow +\infty$ we deduce

$$\lim_{L \rightarrow +\infty} \int_{(0,1) \times \mathbb{R}} \phi(x) \, d\mu^L = \int_{(0,1) \times \mathbb{R}} \phi(x) \, d\mu,$$

which together with (4.10) yield (4.9). $\square$

Proposition 4.2 (Lower bound) *Let $\mathcal{F}_L$ and $\mathcal{F}_\infty$ be as in (2.6) and (2.9) respectively. Let for $L > 0$ be $(w^L, u^L) \subset \mathcal{A}_L^{\text{in}} \times \mathcal{A}_L^{\text{out}}$ a sequence converging to $\mu \in \mathcal{M}_\infty$ in the sense of Definition 2.3. Then there holds*

$$\liminf_{L \rightarrow \infty} \mathcal{F}_L(w^L, u^L) \geq \mathcal{F}_\infty(\mu). \quad (4.12)$$

Proof Let (w^L, u^L) be as in the statement and let $\mu^L := \mu^L(u^L)$ and $\mu_{,x}^L := \mu_{,x}^L(u^L)$ be defined accordingly to Definition 2.1, that is,

$$\mu^L = \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) \mathbb{L}^1 \llcorner (-1, 1) \times \delta_k, \quad \mu_{,x}^L = \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a_{,x}^L(x, k) \mathbb{L}^1 \llcorner (-1, 1) \times \delta_k.$$

Recalling (4.3), (3.3) and (3.4), we have that

$$\begin{aligned}
 \mathcal{F}_L(w^L, u^L) &\geq \int_{-1}^1 \int_{-L}^L (u_{,x}^L)^2 + (u_{,yy}^L)^2 \, dy \, dx \\
 &\geq \int_0^1 \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \frac{1}{4k^2} \frac{(a_{,x}^L(x, k))^2}{a^L(x, k)} + \sum_{k \in \frac{\pi\mathbb{Z}}{L}} a^L(x, k) k^2 \right) dx \\
 &= \int_{(0,1) \times \mathbb{R}} \left(k^2 + \frac{1}{4k^2} \left(\frac{d\mu_{,x}^L}{d\mu^L}(x, k) \right)^2 \right) d\mu^L \\
 &= \int_{(0,1) \times \mathbb{R}} k^2 d\mu^L + \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu^L}{d|\tilde{\mu}^L|}(x, k) \right)^{-1} \\
 &\quad \left(\frac{d\mu_{,x}^L}{d|\tilde{\mu}^L|}(x, k) \right)^2 d|\tilde{\mu}^L|,
 \end{aligned} \tag{4.13}$$

where $\tilde{\mu}^L := (\mu^L, \mu_{,x}^L)$, and the last equality follows from Remark 2.4 (iii). By Reshetnyak Theorem (cf. Ambrosio et al. 2000, Theorem 2.38) there hold

$$\liminf_{L \rightarrow +\infty} \int_{(0,1) \times \mathbb{R}} k^2 d\mu^L \geq \int_{(0,1) \times \mathbb{R}} k^2 d\mu, \tag{4.14}$$

and

$$\begin{aligned}
 &\liminf_{L \rightarrow +\infty} \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu^L}{d|\tilde{\mu}^L|}(x, k) \right)^{-1} \left(\frac{d\mu_{,x}^L}{d|\tilde{\mu}^L|}(x, k) \right)^2 d|\tilde{\mu}^L| \\
 &\geq \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu}{d|\tilde{\mu}|}(x, k) \right)^{-1} \left(\frac{d\mu_{,x}}{d|\tilde{\mu}|}(x, k) \right)^2 d|\tilde{\mu}|,
 \end{aligned} \tag{4.15}$$

with $\tilde{\mu} := (\mu, \mu_{,x})$. Gathering together (4.13), (4.14) and (4.15), we find

$$\begin{aligned}
 \liminf_{L \rightarrow \infty} \mathcal{F}_L(w^L, u^L) &\geq \int_{(0,1) \times \mathbb{R}} k^2 d\mu \\
 &\quad + \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu}{d|\tilde{\mu}|}(x, k) \right)^{-1} \left(\frac{d\mu_{,x}}{d|\tilde{\mu}|}(x, k) \right)^2 d|\tilde{\mu}| \\
 &= \mathcal{F}_\infty(\mu).
 \end{aligned}$$

□

5 Upper Bound

In this section we prove the $\Gamma - \lim \sup$ inequality.

Proposition 5.1 (Upper bound) *Let $\mu \in \mathcal{M}_\infty$. Then for $L > 0$ there exists a sequence $(w^L, u^L) \in \mathcal{A}_L^{\text{in}} \times \mathcal{A}_L^{\text{out}}$ that converges to $\mu \in \mathcal{M}_\infty$ in the sense of Definition 2.3 and such that*

$$\limsup_{L \rightarrow \infty} \mathcal{F}_L(w^L, u^L) \leq \mathcal{F}_\infty(\mu),$$

with $\mathcal{F}_L$ and $\mathcal{F}_\infty$ defined as in (2.6) and (2.9) respectively.

The proof of Proposition 5.1 is quite long and technical; for this reason we divide it into a number of intermediate steps. Let us now summarize the main steps of the upper bound construction. Let $\mu \in \mathcal{M}_\infty$ be given. We first construct a sequence of out-of-plane displacements $(u^L)_{L>0}$ by defining the corresponding Fourier coefficients a_k^L , with $k \in \frac{\pi\mathbb{Z}}{L}$. The latter have to satisfy at the same time:

- be regular enough in order to ensure $u^L \in \mathcal{A}_L^{\text{out}}$;
- the constraint should be (at least approximately) satisfied

$$2A^L(x) := \int_{-L}^L (u_{\cdot,y}^L(x, \cdot))^2 dy = \sum_{k \in \frac{\pi\mathbb{Z}}{L}} (a_k^L(x))^2 k^2 \simeq 2x \text{ for a.e. } x \in (0, 1);$$

- be such that we have good control of partial derivatives of u^L needed to estimate the energy functional $\mathcal{F}_L$.

In order to do that, we proceed as follows. We first dilate the measure μ in the x -variable by a factor of $\lambda > 1$ and get a new measure μ_λ , defined on a larger interval $[-1, \lambda]$ instead of $[-1, 1]$ while keeping the constraint intact. This operation is useful to have enough space near $x = 1$ to mollify in the sequel (Lemma 5.2). As a second step, we discretize μ_λ in the k -variable to obtain a sequence of measures $(\mu^L)_{L>0}$ of the form $\sum_{k \in \frac{\pi\mathbb{Z}}{L}} \bar{b}^L(x, k) \mathcal{L}^1 \llcorner (0, \lambda) \times \delta_k$ (Lemma 5.3). In the third step we regularize each $\bar{b}^L(x, k)$ by convolution with a mollification kernel $\rho_\varepsilon(x)$ at scale $\varepsilon = \varepsilon(L) \rightarrow 0$. Then we set $a_k^L(x) := \frac{1}{k} \sqrt{(\bar{b}^L(\cdot, k) * \rho_\varepsilon)(x)}$ and define u^L accordingly (cf. Lemma 5.4). The mollification procedure possibly produces small error in the constraint, i.e., $A^L(x) = x + o(1)$; hence, in the construction of the recovery sequence we rescale u^L by a factor $f^L(x) := \sqrt{x/A^L(x)}$ to recover back the constraint. Once we construct the out-of-plane displacement, we follow the ideas from Bella (2015) to construct the in-plane displacement (see proof of Proposition 5.1).

For any $\lambda \geq 1$ we define the following class of measures

$$\mathcal{M}_\infty^\lambda := \left\{ \mu \in \mathcal{M}_b^+((0, \lambda) \times \mathbb{R}) : \mu_{\cdot,x} \in \mathcal{M}_b((0, \lambda) \times \mathbb{R}), \quad \mu_{\cdot,x} \ll \mu, \right. \\ \left. \int_0^\lambda 2x\phi(x)dx = \int_{(0,\lambda) \times \mathbb{R}} \phi(x) d\mu(k, x) \quad \forall \phi \in C_c^\infty(0, \lambda) \right\}, \quad (5.1)$$

and the functional $\mathcal{F}_\infty^\lambda: \mathcal{M}_\infty^\lambda \rightarrow [0, +\infty]$

$$\mathcal{F}_\infty^\lambda(\mu) = \int_{(0,\lambda) \times \mathbb{R}} \left[k^2 + \frac{1}{4k^2} \left(\frac{d\mu_{\cdot,x}}{d\mu} \right)^2 \right] d\mu. \quad (5.2)$$

Then we have $\mathcal{M}_\infty = \mathcal{M}_\infty^1$ and $\mathcal{F}_\infty = \mathcal{F}_\infty^1$.

Lemma 5.2 (Dilation of μ) *Let $\mu \in \mathcal{M}_\infty$. Then for each $\lambda \in (1, 2)$ there exists $\mu_\lambda \in \mathcal{M}_\infty^\lambda$ such that*

$$(\mu_\lambda, \mu_{\lambda,\cdot,x}) \llcorner ((0, 1) \times \mathbb{R}) \xrightarrow{*} (\mu, \mu_{\cdot,x}) \text{ as } \lambda \searrow 1, \quad (5.3)$$

and

$$\begin{aligned} \int_{(0,\lambda) \times \mathbb{R}} k^2 d\mu_\lambda &= \lambda^2 \int_{(0,1) \times \mathbb{R}} k^2 d\mu, \\ \int_{(0,\lambda) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu_{\lambda,\cdot,x}}{d\mu_\lambda} \right)^2 d\mu_\lambda &= \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu_{\cdot,x}}{d\mu} \right)^2 d\mu \end{aligned} \quad (5.4)$$

so that, in particular

$$\lim_{\lambda \searrow 1} \mathcal{F}_\infty^\lambda(\mu_\lambda) = \mathcal{F}_\infty(\mu). \quad (5.5)$$

Moreover, $(v_{\frac{x}{\lambda}}, 2x)_{x \in (0,\lambda)}$ is a disintegration of μ_λ where $v_x \in \mathcal{M}_b^+(\mathbb{R})$ for $x \in (0, 1)$ is the measure given by Corollary 3.3.

Proof Let $\mu_\lambda \in \mathcal{M}_b^+((0, \lambda) \times \mathbb{R})$ be defined via duality as follows

$$\int_{(0,\lambda) \times \mathbb{R}} \psi(x, k) d\mu_\lambda = \lambda^2 \int_{(0,1) \times \mathbb{R}} \psi(\lambda x, k) d\mu, \quad (5.6)$$

for every $\psi \in C((0, \lambda) \times \mathbb{R})$. Notice that $\mu_{\lambda,\cdot,x} \in \mathcal{M}_b((0, \lambda) \times \mathbb{R})$ and is given by

$$\int_{(0,\lambda) \times \mathbb{R}} \psi(x, k) d\mu_{\lambda,\cdot,x} = \lambda \int_{(0,1) \times \mathbb{R}} \psi(\lambda x, k) d\mu_{\cdot,x}. \quad (5.7)$$

Hence, $\mu_{\lambda,\cdot,x} \ll \mu_\lambda$ and

$$\frac{d\mu_{\lambda,\cdot,x}}{d\mu_\lambda}(x, k) = \frac{1}{\lambda} \frac{d\mu_{\cdot,x}}{d\mu} \left(\frac{x}{\lambda}, k \right).$$

Moreover, (5.6) together with the change of variable $x = \lambda \hat{x}$ implies

$$\int_0^\lambda 2x \phi(x) dx = \int_{(0,\lambda) \times \mathbb{R}} \phi(x) d\mu_\lambda \quad \forall \phi \in C_c^\infty(0, \lambda).$$

It follows that $\mu_\lambda \in \mathcal{M}_\infty^\lambda$. Moreover, from (5.6) and (5.7) we deduce that

$$(\mu_\lambda, \mu_{\lambda,x}) \llcorner (\mathbb{R} \times (0, 1)) \xrightarrow{*} (\mu, \mu_{,x}) \quad \text{as } \lambda \rightarrow 1,$$

and (5.4) which implies (5.5). The fact that $(\nu_{\frac{x}{\lambda}}, 2x)_{x \in (0, \lambda)}$ is a disintegration of μ_λ follows again by a change of variable. $\square$

Lemma 5.3 (Discretisation of μ) *Let $\mu \in \mathcal{M}_\infty$ with $\mathcal{F}_\infty(\mu) < +\infty$ and let $\lambda = \lambda(L) \searrow 1$ as $L \rightarrow \infty$. Then there exists $(\mu^L) \subset \mathcal{M}_\infty^\lambda$ with the following properties:*

(i) $\mu^L = \sum_{k \in \frac{\pi\mathbb{Z}}{L}} \bar{b}^L(x, k) \mathbb{L}^1 \llcorner (0, \lambda) \times \delta_k$ with

$$\bar{b}^L(\cdot, k) \in W^{1,1}(0, \lambda) \quad \text{and} \quad \sum_{k \in \frac{\pi\mathbb{Z}}{L}} \bar{b}^L(x, k) = 2x, \quad \forall x \in (0, \lambda), \quad (5.8)$$

$$\begin{aligned} \mathcal{F}_\infty^\lambda(\mu^L) &= \int_0^\lambda \sum_{k \in \frac{\pi\mathbb{Z}}{L}} k^2 \bar{b}^L(x, k) \, dx \\ &\quad + \int_0^\lambda \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \frac{1}{4k^2} \frac{(\bar{b}_{,x}^L(x, k))^2}{\bar{b}^L(x, k)} \, dx; \end{aligned} \quad (5.9)$$

(ii) $(\mu^L, \mu_{,x}^L) \llcorner ((0, 1) \times \mathbb{R}) \xrightarrow{*} (\mu, \mu_{,x})$;

(iii) $\limsup_{L \rightarrow \infty} \mathcal{F}_\infty^\lambda(\mu^L) \leq \mathcal{F}_\infty(\mu)$.

Proof For each $L > 1$ let $\mu_\lambda \in \mathcal{M}_\infty^\lambda$ be the measure given by Lemma 5.2 and let $(\nu_{\frac{x}{\lambda}}, 2x)_{x \in (0, \lambda)}$ be the corresponding disintegration. We then define $\mu^L \in \mathcal{M}_b^+((0, \lambda) \times \mathbb{R})$ as

$$\mu^L := \sum_{k \in \frac{\pi\mathbb{Z}}{L}} \bar{b}^L(x, k) \mathbb{L}^1 \llcorner (0, \lambda) \times \delta_k, \quad (5.10)$$

where for $(x, k) \in (0, \lambda) \times \frac{\pi\mathbb{Z}}{L}$ we set

$$\bar{b}^L(x, k) := \begin{cases} 0 & \text{if } k = 0, \\ 2x \nu_{\frac{x}{\lambda}}(I_k^L) & \text{if } k \neq 0, \end{cases} \quad \text{and} \quad I_k^L := \begin{cases} (k - \frac{\pi}{L}, k] & \text{if } k > 0, \\ [k, k + \frac{\pi}{L}) & \text{if } k < 0. \end{cases} \quad (5.11)$$

Now, for each $k \in \frac{\pi\mathbb{Z}}{L}$, $\bar{b}^L(\cdot, k) \in W^{1,1}(0, \lambda)$ with

$$\bar{b}_{,x}^L(x, k) = \begin{cases} 0 & \text{if } k = 0, \\ 2x \int_{I_k^L} \frac{d\mu_{\lambda,x}}{d\mu_\lambda}(x, \hat{k}) \, d\nu_{\frac{x}{\lambda}}(\hat{k}) & \text{if } k \neq 0. \end{cases}$$

Indeed, if $k = 0$ there is nothing to prove. If instead $k \neq 0$, for $\phi \in C_c^\infty(0, \lambda)$ from the definition of $\nu_{\frac{x}{\lambda}}$ and recalling Remark 2.2 we get

$$\begin{aligned} \int_0^\lambda \bar{b}^L(x, k) \dot{\phi}(x) dx &= \int_0^\lambda \int_{\mathbb{R}} \chi_{I_k^L}(\hat{k}) \dot{\phi}(x) d\nu_{\frac{x}{\lambda}}(\hat{k}) 2x dx \\ &= \int_{(0, \lambda) \times \mathbb{R}} \chi_{I_k^L}(\hat{k}) \dot{\phi}(x) d\mu_\lambda \\ &= - \int_{(0, \lambda) \times \mathbb{R}} \chi_{I_k^L}(\hat{k}) \phi(x) d\mu_{\lambda, x} \\ &= - \int_{(0, \lambda) \times \mathbb{R}} \chi_{I_k^L}(\hat{k}) \phi(x) \frac{d\mu_{\lambda, x}}{d\mu_\lambda}(x, \hat{k}) d\mu_\lambda \\ &= - \int_0^\lambda \int_{I_k^L} \frac{d\mu_{\lambda, x}}{d\mu_\lambda}(x, \hat{k}) d\nu_{\frac{x}{\lambda}}(\hat{k}) 2x \phi(x) dx; \end{aligned}$$

furthermore by Young's inequality and (5.5)

$$\begin{aligned} \int_0^\lambda |\bar{b}_{,x}^L(x, k)| dx &\leq \int_{I_k^L \times (0, \lambda)} \left| \frac{d\mu_{\lambda, x}}{d\mu_\lambda} \right| d\mu_\lambda \\ &\leq \frac{1}{2} \int_{I_k^L \times (0, \lambda)} \left(k^2 + \frac{1}{k^2} \left(\frac{d\mu_{\lambda, x}}{d\mu_\lambda} \right)^2 \right) d\mu_\lambda \leq C. \end{aligned}$$

Thus, in particular

$$\int_0^\lambda \sum_{k \in \frac{\pi\mathbb{Z}}{L}} |\bar{b}_{,x}^L(x, k)| dx \leq \mathcal{F}_\infty^\lambda(\mu_\lambda) \leq C.$$

As a consequence we have that $\mu_{,x}^L \in \mathcal{M}_b((0, \lambda) \times \mathbb{R})$, and

$$\mu_{,x}^L = \sum_{k \in \frac{\pi\mathbb{Z}}{L}} \bar{b}_{,x}^L(x, k) \mathbb{1}_{\square(0, \lambda) \times \delta_k},$$

and by Remark 2.2 (ii) $\tilde{\mu}_{,x}^L \ll \tilde{\mu}^L$. Moreover, as $\nu_{\frac{x}{\lambda}}$ is a probability measure, there holds

$$\sum_{k \in \frac{\pi\mathbb{Z}}{L}} \bar{b}^L(x, k) = 2x \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \nu_{\frac{x}{\lambda}}(I_k^L) \right) = 2x, \quad \forall x \in (0, \lambda).$$

Note in particular that $\mu^L \in \mathcal{M}_\infty^\lambda$. Moreover, (5.9) readily follows and (i) is proved. We next show (ii). Take $\varphi \in C_c^\infty((0, 1) \times \mathbb{R})$, so that from (5.11) we obtain

$$\begin{aligned} \int_{(0,1) \times \mathbb{R}} \varphi \, d\mu^L &= \int_0^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \varphi(x, k) v_{\frac{x}{\lambda}}(I_k^L) 2x \, dx \\ &= \int_0^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \int_{I_k^L} (\varphi(x, k) - \varphi(x, \hat{k})) \, dv_{\frac{x}{\lambda}}(\hat{k}) 2x \, dx \quad (5.12) \\ &\quad + \int_{(0,1) \times \mathbb{R}} \varphi \, d\mu_\lambda. \end{aligned}$$

Since φ is uniformly continuous for every $\varepsilon > 0$, there is $L_0 > 1$ such that for all $L \geq L_0$

$$|\varphi(x, k) - \varphi(x, \hat{k})| < \varepsilon \quad \forall x \in (0, \lambda), \quad \forall k \in \frac{\pi\mathbb{Z}}{L}, \quad \forall \hat{k} \in I_k^L,$$

from which we readily deduce that

$$\begin{aligned} \int_0^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \int_{I_k^L} |\varphi(x, k) - \varphi(x, \hat{k})| \, dv_{\frac{x}{\lambda}}(\hat{k}) 2x \, dx \\ \leq \mu_\lambda((0, \lambda) \times \mathbb{R}) \varepsilon = \lambda^2 \mu((0, 1) \times \mathbb{R}) \varepsilon. \end{aligned} \quad (5.13)$$

From (5.12), (5.13), (5.3) and the arbitrariness of ε , we infer $\mu^L \llcorner ((0, 1) \times \mathbb{R}) \xrightarrow{*} \mu$ as $L \rightarrow +\infty$. By analogous arguments we get $\mu_{\lambda,x}^L \llcorner ((0, 1) \times \mathbb{R}) \xrightarrow{*} \mu_{\lambda,x}$ as $L \rightarrow +\infty$. It remains to prove (iii). We start by observing that for all $\delta > 0$ and all $\hat{k} \in I_k^L$ we have

$$k^2 \leq \left(\hat{k} + \frac{\pi}{L} \right)^2 \leq (1 + \delta) \hat{k}^2 + (1 + \delta^{-1}) \frac{\pi^2}{L^2},$$

so that

$$\begin{aligned} \int_0^\lambda \sum_{k \in \frac{\pi\mathbb{Z}}{L}} k^2 \bar{b}^L(x, k) \, dx &= \int_0^\lambda \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \int_{I_k^L} k^2 \, dv_{\frac{x}{\lambda}}(\hat{k}) 2x \, dx \\ &= \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \int_{(0,\lambda) \times I_k^L} k^2 \, d\mu_\lambda(x, \hat{k}) \\ &\leq \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \int_{(0,\lambda) \times I_k^L} \left((1 + \delta) \hat{k}^2 + (1 + \delta^{-1}) \frac{\pi^2}{L^2} \right) d\mu_\lambda(x, \hat{k}) \\ &= (1 + \delta) \int_{(0,\lambda) \times \mathbb{R}} \hat{k}^2 \, d\mu_\lambda + \mu_\lambda((0, \lambda) \times \mathbb{R}) (1 + \delta^{-1}) \frac{\pi^2}{L^2}. \end{aligned} \quad (5.14)$$

Moreover, there holds

$$\int_0^\lambda \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \frac{1}{4k^2} \frac{(\bar{b}_{,x}^L(x, k))^2}{\bar{b}^L(x, k)} dx = \int_0^\lambda \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \frac{1}{4k^2} \left(\frac{\bar{b}_{,x}^L(x, k)}{\bar{b}^L(x, k)} \right)^2 \bar{b}^L(x, k) dx. \quad (5.15)$$

Since $1/|k| \leq 1/|\hat{k}|$ for $\hat{k} \in I_k^L$ the following inequality follows

$$\frac{1}{2|k|} \frac{\bar{b}_{,x}^L(x, k)}{\bar{b}^L(x, k)} = \frac{1}{2|k|} \int_{I_k^L} \frac{d\mu_{\lambda,x}}{d\mu_\lambda}(x, \hat{k}) dv_{\frac{x}{\lambda}}(\hat{k}) \leq \int_{I_k^L} \frac{1}{2|\hat{k}|} \frac{d\mu_{\lambda,x}}{d\mu_\lambda}(x, \hat{k}) dv_{\frac{x}{\lambda}}(\hat{k}). \quad (5.16)$$

Now combining (5.15) with (5.16), we obtain

$$\begin{aligned} & \int_0^\lambda \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \frac{1}{4k^2} \frac{(\bar{b}_{,x}^L(x, k))^2}{\bar{b}^L(x, k)} dx \\ &= \int_0^\lambda \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \left(\int_{I_k^L} \frac{1}{2|\hat{k}|} \frac{d\mu_{\lambda,x}}{d\mu_\lambda}(x, \hat{k}) dv_{\frac{x}{\lambda}}(\hat{k}) \right)^2 v_{\frac{x}{\lambda}}(I_k^L) 2x dx \\ &\leq \int_0^\lambda \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \int_{I_k^L} \frac{1}{4\hat{k}^2} \left(\frac{d\mu_{\lambda,x}}{d\mu_\lambda}(x, \hat{k}) \right)^2 dv_{\frac{x}{\lambda}}(\hat{k}) 2x dx \\ &= \int_{(0,\lambda) \times \mathbb{R}} \frac{1}{4\hat{k}^2} \left(\frac{d\mu_{\lambda,x}}{d\mu_\lambda}(x, \hat{k}) \right)^2 d\mu_\lambda \\ &= \int_{(0,1) \times \mathbb{R}} \frac{1}{4\hat{k}^2} \left(\frac{d\mu_{,x}}{d\mu}(x, \hat{k}) \right)^2 d\mu, \end{aligned} \quad (5.17)$$

where the inequality follows by Jensen's inequality. Finally, gathering together (5.14) and (5.17) and recalling (5.5), we infer

$$\limsup_{L \rightarrow \infty} \mathcal{F}_\infty^\lambda(\mu^L) \leq (1 + \delta) \limsup_{L \rightarrow \infty} \mathcal{F}_\infty^\lambda(\mu_\lambda) \leq (1 + \delta) \mathcal{F}_\infty(\mu).$$

By arbitrariness of δ , (iii) follows by letting $\delta \rightarrow 0$. $\square$

In the next lemma we continue our quest toward definition of the in-plane displacement. Picking up the construction of Fourier coefficients from Lemma 5.3, to conclude we require its higher regularity (in particular in the x -variable). A natural idea would be to mollify the coefficients, but since we want to preserve the constraint (or at least not to change it much), we will mollify squares of the coefficients—in the end the constraint is sum (integral) of these squares, so it behaves well w.r.t to their mollification. Unfortunately, we could not use simple compactly supported smoothing kernel, since

for our kernel we require that its derivative can be controlled by the kernel itself—this satisfies, e.g., an exponentially decaying kernel, but not compactly supported kernel. Using $e^{-|x|}$ as kernel works well provided one can reasonably extend the coefficients (precisely their squares) to the whole $\mathbb{R}$. This is trivial for $x < -1$ (zero extension) and also for $x > \lambda$ (extension by a suitable constant). Even though obviously anywhere in $x < 0$ and $x > \lambda$ the constraint is not equal x , which would be enough to show using symmetry of the kernel that the constraint after mollification stays intact, as long as x stays away from these two sets the violation from the constraint being x , denoted by f^L , is exponentially small. Since we need to focus on the interval $[0, 1]$, we definitely stay away from the set $x > \lambda$ (at least measured at the ε -scale). Near $x = 0$ the situation is completely different (the constraint violation f^L is far from 1 there), but this critical region is relatively small (which helps to get the estimates). We therefore use multiplication by f^L to completely recover the constraint, but also cut-off near $x = 0$ on scale even smaller than the convolution scale (see proof of Proposition 5.1 below).

Though it might look strange to first mollify the coefficients which destroys the constraint, and then multiply by the error to recover the constraint back, it is very natural: for the error we have basically explicit formula, since it is expressed in terms of mollification of the constraint and not of single coefficients, and we can therefore control it very well—including its higher derivatives, so that we can still control the regularity.

Lemma 5.4 (Construction of u) *Let $\mu \in \mathcal{M}_\infty$ be such that $\mathcal{F}_\infty(\mu) < +\infty$. Let $\varepsilon = \varepsilon(L) > 0$ and $n = n(L) \in \mathbb{N}$ be such that*

$$\lim_{L \rightarrow +\infty} \varepsilon(L) = 0, \quad \lim_{L \rightarrow +\infty} n(L) = \lim_{L \rightarrow +\infty} \frac{L}{n(L)} = +\infty.$$

Then there exists $\hat{u}^L \in \mathcal{A}_L^{\text{out}} \cap \mathcal{A}_{L_0}^{\text{out}}$ with $L_0 := L/n(L)$ that satisfies the following properties: let

$$A^L(x) := \frac{1}{2} \int_{-L}^L (\hat{u}_{\cdot,y}^L(x, \cdot))^2 dy \quad \text{and} \quad f^L(x) := \sqrt{\frac{x}{A^L(x)}}.$$

Then for all $x \in (0, 1)$ and $N \in \mathbb{N}$ there holds

$$\max\{x, \varepsilon\} \lesssim A^L(x) \lesssim \max\{x, \varepsilon\}; \quad (5.18)$$

$$(f^L(x))^2 \lesssim \frac{x}{\max\{x, \varepsilon\}}, \quad (f^L(x))^2 \leq 1 + o_L(1); \quad (5.19)$$

$$(f^L(x))^2 \geq 1 + o_N(1) \quad \text{if } x \in (N\varepsilon, 1); \quad (5.20)$$

$$(f^L(x))^2 \lesssim \frac{\max\{e^{-\frac{x}{\varepsilon}}, e^{-\frac{1}{\sqrt{\varepsilon}}}\}}{x\varepsilon}, \quad (\ddot{f}^L(x))^2 \lesssim \frac{1}{x^3\varepsilon}; \quad (5.21)$$

there exists a continuous increasing function $\omega: [0, +\infty) \rightarrow [0, +\infty)$ with $\omega(0) = 0$ such that

$$\int_{-L}^L (\hat{u}^L(x, \cdot))^2 dy \lesssim \begin{cases} \max\{x, \varepsilon\}(\omega(2N\varepsilon) + Ne^{-N}) & \text{if } x \in (0, N\varepsilon) \\ x & \text{if } x \in (N\varepsilon, 1) \end{cases} \quad (5.22)$$

$$\int_{-L}^L (\hat{u}^L(x, \cdot))^4 dy \lesssim L_0^2 (\max\{\varepsilon, x\})^2; \quad (5.23)$$

$$\int_{-L}^L \left((\hat{u}_{,yy}^L(x, \cdot))^2 + (\hat{u}_{,x}^L(x, \cdot))^2 \right) dy \lesssim \frac{1}{\varepsilon}. \quad (5.24)$$

Moreover,

$$(\mu^L(\hat{u}^L), \mu_{,x}^L(\hat{u}^L)) \overset{*}{\rightharpoonup} (\mu, \mu_{,x}) \text{ in } \mathcal{M}_b((-1, 1) \times \mathbb{R})^2; \quad (5.25)$$

$$\limsup_{L \rightarrow \infty} \int_{-L}^L \int_{-1}^1 ((\hat{u}_{,x}^L)^2 + (\hat{u}_{,yy}^L)^2) dx dy \leq \mathcal{F}_\infty(\mu); \quad (5.26)$$

$$\int_{-L}^L \int_{-1}^1 \left((\hat{u}_{,xy}^L)^2 + (\hat{u}_{,xx}^L)^2 + (\hat{u}_{,yyx}^L)^2 \right) dx dy \lesssim \frac{1}{\varepsilon^2}; \quad (5.27)$$

$$\int_{-L}^L \int_{-1}^1 (\hat{u}_{,x}^L)^4 dx dy \lesssim \frac{L_0^2}{\varepsilon^2}. \quad (5.28)$$

Finally, since $\hat{u}^L$ and all its derivatives are $2L_0$ -periodic in the y -variable, the above estimates still hold true if we replace the average integral on $[-L, L]$ with the average integral on $[-L_0, L_0]$.

Proof Let $\mu \in \mathcal{M}_\infty$, $\varepsilon = \varepsilon(L)$, $n = n(L)$ and L_0 be as in the statement. We set

$$\lambda = \lambda(L) := (1 + \sqrt{\varepsilon}) \searrow 1 \text{ as } L \rightarrow +\infty,$$

hence in particular $\lambda \searrow 1$ as $L_0 \rightarrow +\infty$.

We construct $\hat{u}^L \in \mathcal{A}_{L_0}^{\text{out}}$ and then we extend it periodically in $[-1, 1] \times [-L, L]$, without relabeling it. In this way, since $L = n(L)L_0$ with $n(L) \in \mathbb{N}$, we have $\hat{u}^L \in \mathcal{A}_L^{\text{out}}$. The main idea is that of discretizing the measure μ in the variable k to get a measure concentrated on lines $\mathbb{R} \times \{k\}$ with $k \in \frac{\pi\mathbb{Z}}{L_0}$ where the weight on each line is a coefficient $b^{L_0}(x, k)$. Afterward, we define $\hat{u}^L$ as in such a way that its Fourier coefficients $a_k^L(x)$ are as close as possible to $\sqrt{b^{L_0}(x, k)/k}$ but at the same time have better regularity to ensure $\hat{u}^L \in \mathcal{A}_L^{\text{out}}$. In order to do that, we first dilate μ with a factor λ in the x -variable as in Lemma 5.2, then we discretize using Lemma 5.3, and finally we mollify $b^{L_0}(\cdot, k)$ at scale ε after a suitable extension in $\mathbb{R}$.

To this purpose we let $(\mu^{L_0}) \subset \mathcal{M}_\infty^\lambda$ be the sequence of Lemma 5.3 for the parameter L_0 , which is of the form

$$\mu^{L_0} = \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(x, k) \mathbb{L}^1 \llcorner (0, \lambda) \times \delta_k.$$

By the mean value theorem, for each λ , we can find $\bar{\lambda} \in (\frac{\lambda+1}{2}, \lambda)$ such that

$$\sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(\bar{\lambda}, k) k^2 \leq \int_{\frac{\lambda+1}{2}}^{\lambda} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(x, k) k^2 dx. \quad (5.29)$$

By truncating $\bar{b}^{L_0}$ at $x = \bar{\lambda}$ we can define $b^{L_0}: \mathbb{R} \times \frac{\pi\mathbb{Z}}{L_0} \rightarrow \mathbb{R}$ as

$$b^{L_0}(x, k) := \begin{cases} 0 & \text{if } x \leq 0, \\ \bar{b}^{L_0}(x, k) & \text{if } 0 < x < \bar{\lambda}, \\ \bar{b}^{L_0}(\bar{\lambda}, k) & \text{if } x \geq \bar{\lambda}. \end{cases} \quad (5.30)$$

Let $\rho_\varepsilon(x) := \frac{1}{2\varepsilon} e^{\frac{-|x|}{\varepsilon}}$ and note that in particular

$$|\dot{\rho}_\varepsilon(x)| = \frac{1}{\varepsilon} \rho_\varepsilon(x). \quad (5.31)$$

Finally we let $a^L: \mathbb{R} \times \frac{\pi\mathbb{Z}}{L_0} \rightarrow \mathbb{R}$ be defined as

$$a^L(x, k) := (b^{L_0}(\cdot, k) * \rho_\varepsilon)(x),$$

and $\hat{u}^L \in \mathcal{A}_{L_0}^{\text{out}}$ be the function

$$\hat{u}^L(x, y) := \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k > 0} \frac{\sqrt{a^L(x, k)}}{k} \sqrt{2} \sin(ky) + \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k < 0} \frac{\sqrt{a^L(x, k)}}{k} \sqrt{2} \cos(ky).$$

Eventually we extend $\hat{u}^L$, without relabeling it, periodically in $[-1, 1] \times [-L, L]$.

Step 1: in this step we show (5.18)–(5.21). By (3.3) and (5.8) we have that

$$\begin{aligned} 2A^L(x) &= \int_{-L}^L (\hat{u}_{,y})^2 dy = \int_{-L_0}^{L_0} (\hat{u}_{,y})^2 dy \\ &= \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} a^L(x, k) = \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} b^{L_0}(\cdot, k) \right) * \rho_\varepsilon(x) \\ &= 2(x\chi_{(0, \bar{\lambda})} + \bar{\lambda}\chi_{(\bar{\lambda}, +\infty)}) * \rho_\varepsilon(x) = 2x\chi_{\{x \geq 0\}} + \varepsilon \left(e^{\frac{-|x|}{\varepsilon}} - e^{\frac{x-\bar{\lambda}}{\varepsilon}} \right), \end{aligned} \quad (5.32)$$

for all $x \in [-1, 1]$. Then for ε small enough we have

$$\begin{aligned} \frac{\varepsilon}{2e} &\leq 2A^L(x) \leq 3\varepsilon \quad \text{if } x \in (0, \varepsilon), \\ x &\leq 2x + \varepsilon \left(e^{\frac{-1}{\varepsilon}} - e^{\frac{1-\bar{\lambda}}{\varepsilon}} \right) \leq 2A^L(x) \leq 3x \quad \text{if } x \in (\varepsilon, 1), \end{aligned}$$

so that

$$\frac{1}{2e} \max\{x, \varepsilon\} \leq 2A^L(x) = \int_{-L}^L (\hat{u}_y)^2 dy \leq 3 \max\{x, \varepsilon\}.$$

We have that

$$(f^L(x))^2 = \frac{x}{A^L(x)},$$

and by estimates above

$$(f^L(x))^2 \lesssim \frac{x}{\max\{x, \varepsilon\}} \lesssim \frac{x}{\varepsilon} \quad \text{if } x \in (0, \varepsilon).$$

Observing that $(e^{\frac{-x}{\varepsilon}} - e^{\frac{x-\bar{\lambda}}{\varepsilon}}) \geq 0$ if and only if $x \in (0, \bar{\lambda}/2)$ we have

$$(f^L(x))^2 = \frac{x}{x + \frac{\varepsilon}{2}(e^{\frac{-x}{\varepsilon}} - e^{\frac{x-\bar{\lambda}}{\varepsilon}})} \leq 1 \quad \text{in } (0, \bar{\lambda}/2),$$

and

$$\begin{aligned} (f^L(x))^2 &\leq \frac{x}{x + \frac{\varepsilon}{2}(e^{\frac{-1}{\varepsilon}} - e^{\frac{1-\bar{\lambda}}{\varepsilon}})} \\ &\leq 1 + \frac{\frac{\varepsilon}{2}|e^{\frac{-1}{\varepsilon}} - e^{\frac{1-\bar{\lambda}}{\varepsilon}}|}{\frac{1}{2} + \frac{\varepsilon}{2}(e^{\frac{-1}{\varepsilon}} - e^{\frac{1-\bar{\lambda}}{\varepsilon}})} \\ &= 1 + \frac{\varepsilon}{8} \quad \text{for } x \in (\bar{\lambda}/2, 1). \end{aligned}$$

Moreover, we have that

$$\begin{aligned} (f^L(x))^2 &\geq \frac{x}{x + \frac{\varepsilon}{2}e^{-N}} = 1 - \frac{\frac{\varepsilon}{2}e^{-N}}{x + \frac{\varepsilon}{2}e^{-N}} \geq 1 - \frac{\frac{\varepsilon}{2}e^{-N}}{N\varepsilon} \\ &= 1 + o_N(1) \quad \text{for } x \in [N\varepsilon, 1). \end{aligned}$$

A direct computation shows that

$$\begin{aligned} (\dot{f}^L(x))^2 &= \frac{(A^L(x) - x\dot{A}^L(x))^2}{4x(A^L(x))^3} \\ &\lesssim \frac{(\max\{x, \varepsilon\})^2 \max\{e^{-\frac{x}{\varepsilon}}, e^{\frac{1-\bar{\lambda}}{\varepsilon}}\}}{x(\max\{x, \varepsilon\})^3} \\ &\lesssim \frac{\max\{e^{-\frac{x}{\varepsilon}}, e^{\frac{1}{\sqrt{\varepsilon}}}\}}{x \max\{x, \varepsilon\}} \lesssim \frac{\max\{e^{-\frac{x}{\varepsilon}}, e^{\frac{1}{\sqrt{\varepsilon}}}\}}{x\varepsilon}, \end{aligned}$$

where the second inequality follows from $\frac{1-\bar{\lambda}}{\varepsilon} \leq \frac{1-\frac{\lambda+1}{2}}{\varepsilon} = \frac{1}{2\sqrt{\varepsilon}}$. Furthermore we have

$$\begin{aligned} (\ddot{f}^L(x))^2 &\lesssim \frac{x(\ddot{A}^L(x))^2}{(A^L(x))^3} + \frac{1}{x^3 A^L(x)} \\ &\quad + \frac{(\dot{A}^L(x))^2}{x(A^L(x))^3} + \frac{x(\dot{A}^L(x))^4}{(A^L(x))^5} \\ &\lesssim \frac{x\varepsilon^{-2}e^{-\frac{x}{\varepsilon}}}{(\max\{x, \varepsilon\})^3} + \frac{1}{x^3 \max\{x, \varepsilon\}} \\ &\quad + \frac{1}{x(\max\{x, \varepsilon\})^3} + \frac{x}{(\max\{x, \varepsilon\})^5}. \end{aligned} \quad (5.33)$$

Hence,

$$(\ddot{f}^L(x))^2 \lesssim \frac{1}{\varepsilon^4} + \frac{1}{x^3 \varepsilon} + \frac{1}{x \varepsilon^3} \lesssim \frac{1}{x^3 \varepsilon} \quad \text{if } x \in (0, \varepsilon),$$

and

$$(\ddot{f}^L(x))^2 \lesssim \frac{e^{-\frac{x}{\varepsilon}}}{x^2 \varepsilon^2} + \frac{1}{x^4} \lesssim \frac{1}{x^3 \varepsilon} \quad \text{if } x \in (\varepsilon, 1).$$

Step 2: in this step we show (5.22). By (3.2) it holds

$$\begin{aligned} \int_{-L}^L (\hat{u}^L(x, \cdot))^2 dy &= \int_{-L_0}^{L_0} (\hat{u}^L(x, \cdot))^2 dy = \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{a^L(x, k)}{k^2} \\ &= \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{b^{L_0}(\cdot, k)}{k^2} \right) * \rho_\varepsilon(x) \\ &= \int_0^{+\infty} \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{b^{L_0}(z, k)}{k^2} \right) \rho_\varepsilon(x - z) dz. \end{aligned} \quad (5.34)$$

Moreover, by the fundamental theorem of calculus and Hölder's inequality we have

$$b^{L_0}(z, k) = \left(\sqrt{b^{L_0}(z, k)} \right)^2 = \left(\int_0^z \frac{b_{,x}^{L_0}(\hat{z}, k)}{2\sqrt{b^{L_0}(\hat{z}, k)}} d\hat{z} \right)^2 \leq z \int_0^z \frac{(b_{,x}^{L_0}(\hat{z}, k))^2}{4b^{L_0}(\hat{z}, k)} d\hat{z}. \quad (5.35)$$

Combining (5.34) with (5.35) we find

$$\int_{-L}^L (\hat{u}^L(x, \cdot))^2 dy \leq \int_0^{+\infty} \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \int_0^z \frac{(b_{,x}^{L_0}(\hat{z}, k))^2}{4k^2 b^{L_0}(\hat{z}, k)} d\hat{z} \right) z \rho_\varepsilon(x - z) dz$$

By definition of b^{L_0} it follows that

$$\begin{aligned} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \int_0^z \frac{(b_{,x}^{L_0}(\hat{z}, k))^2}{4k^2 b^{L_0}(\hat{z}, k)} d\hat{z} &= \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \int_0^{z \wedge \bar{\lambda}} \frac{(\bar{b}_{,x}^{L_0}(\hat{z}, k))^2}{4k^2 \bar{b}^{L_0}(\hat{z}, k)} d\hat{z} \\ &\leq \int_{(0, z \wedge \bar{\lambda}) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu_{,x}^{L_0}}{d\mu^{L_0}} \right)^2 d\mu^{L_0} =: \omega(z), \end{aligned} \quad (5.36)$$

where the last inequality can be obtained by arguing exactly as in (5.17). Therefore, we deduce that

$$\oint_{-L_0}^{L_0} (\hat{u}^L(x, \cdot))^2 dy \leq \int_0^{+\infty} \omega(z) z \rho_\varepsilon(x - z) dz.$$

Note that $\omega(z) \rightarrow 0$ as $z \rightarrow 0$ and $\omega(z) \leq \omega(\bar{\lambda}) \leq (\lambda^2) \mathcal{F}_\infty(\mu) \leq C$. Let $N \geq 2$ be a natural number. Assume $x \in (0, N\varepsilon]$. Since ω is increasing, we have

$$\begin{aligned} \int_0^{+\infty} \omega(z) z \rho_\varepsilon(x - z) dz &\leq \omega(2N\varepsilon) \int_0^{2N\varepsilon} z \rho_\varepsilon(x - z) dz \\ &\quad + \omega(\bar{\lambda}) \int_{2N\varepsilon}^{+\infty} z \rho_\varepsilon(x - z) dz \\ &\leq \omega(2N\varepsilon) \max\{x, \varepsilon\} + CNe^{-N}\varepsilon \\ &\lesssim \max\{x, \varepsilon\}(\omega(2N\varepsilon) + Ne^{-N}). \end{aligned} \quad (5.37)$$

If instead $x \in (N\varepsilon, 1)$, we get

$$\int_0^{+\infty} \omega(z) z \rho_\varepsilon(x - z) dz \leq \omega(\bar{\lambda}) \int_0^{+\infty} z \rho_\varepsilon(x - z) dz \lesssim x. \quad (5.38)$$

Step 3: in this step we show (5.23). By the mean value theorem and the fact that $u^L(x, \cdot)$ is $2L_0$ -periodic, for fixed x , we can find $y_0 = y_0(x) \in [-L_0, L_0]$ such that

$$\hat{u}^L(x, y_0) = \oint_{-L_0}^{L_0} \hat{u}^L(x, \hat{y}) d\hat{y} = 0,$$

where the second equality follows by the definition of $\hat{u}^L$ and the fact that

$$\begin{aligned} \oint_{-L_0}^{L_0} \sin(k\hat{y}) d\hat{y} &= \frac{k}{2L_0} (-\cos(kL_0) + \cos(kL_0)) = 0, \\ \oint_{-L_0}^{L_0} \cos(k\hat{y}) d\hat{y} &= \frac{k}{2L_0} (\sin(kL_0) - \sin(kL_0)) = 0, \end{aligned}$$

for all $k \in \frac{\pi\mathbb{Z}}{L_0}$. Thus, by the fundamental theorem of calculus, Hölder's inequality and Plancherel it holds

$$\begin{aligned} |\hat{u}^L(x, y)| &= \left| \int_{y_0}^y \hat{u}_{,y}^L(x, y') \, dy' \right| \leq \sqrt{2L_0} \left(\int_{-L_0}^{L_0} (\hat{u}_{,y}^L)^2 \, dy' \right)^{\frac{1}{2}} \\ &= 2\sqrt{2}L_0\sqrt{A^L(x)}. \end{aligned}$$

This together with steps 1 and 2 yields

$$\begin{aligned} \oint_{-L}^L (\hat{u}^L)^4 \, dy &= \oint_{-L_0}^{L_0} \int_{-1}^1 (\hat{u}^L)^4 \, dy \\ &\lesssim L_0^2 A^L(x) \oint_{-L_0}^{L_0} (\hat{u}^L)^2 \, dy \lesssim L_0^2 (\max\{x, \varepsilon\})^2. \end{aligned}$$

Step 4: in this step we show (5.24). By (3.3) the definition of a^L and (5.29)

$$\begin{aligned} \oint_{-L}^L (\hat{u}_{,yy}^L)^2 \, dy &= \oint_{-L_0}^{L_0} (\hat{u}_{,yy}^L)^2 \, dy = \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} k^2 a^L(x, k) = \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} k^2 b^{L_0}(x, \cdot) * \rho_\varepsilon(x) \\ &= \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \left(k^2 \int_0^{\bar{\lambda}} \bar{b}^{L_0}(z, k) \rho_\varepsilon(x - z) \, dz \right) \\ &\quad + \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(\bar{\lambda}, k) k^2 \int_{\bar{\lambda}}^{+\infty} \rho_\varepsilon(x - z) \, dz \\ &\leq \|\rho_\varepsilon\|_\infty \int_0^{\bar{\lambda}} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(x, k) k^2 \, dx \\ &\quad + e^{\frac{x-\bar{\lambda}}{\varepsilon}} \oint_{\frac{\lambda+1}{2}}^{\bar{\lambda}} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(x, k) k^2 \, dx \\ &\lesssim \left(\frac{1}{\varepsilon} + \frac{1}{\lambda-1} e^{\frac{x-\bar{\lambda}}{\varepsilon}} \right) \int_0^{\bar{\lambda}} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(x, k) k^2 \, dx \\ &\lesssim \left(\frac{1}{\varepsilon} + \frac{1}{\lambda-1} e^{\frac{x-\bar{\lambda}}{\varepsilon}} \right) \mathcal{F}_\infty^\lambda(\mu^{L_0}) \lesssim \frac{1}{\varepsilon}. \end{aligned} \tag{5.39}$$

Since the function $(z_1, z_2) \mapsto z_1^2/z_2$ is convex, by Jensen's inequality we have

$$\frac{(a_{,x}^L(x, k))^2}{a^L(x, k)} = \frac{(b_{,x}^{L_0}(\cdot, k) * \rho_\varepsilon(x))^2}{b^{L_0}(\cdot, k) * \rho_\varepsilon(x)} \leq \frac{(b_{,x}^{L_0}(\cdot, k))^2}{b^{L_0}(\cdot, k)} * \rho_\varepsilon(x).$$

This together with (3.3) and (3.4) implies

$$\begin{aligned}
 \int_{-L}^L (u_{,x}^L)^2 dy &= \int_{-L_0}^{L_0} (u_{,x}^L)^2 dy = \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{4k^2} \frac{(a_{,x}^L(x, k))^2}{a^L(x, k)} \\
 &\leq \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{4k^2} \frac{(b_{,x}^{L_0}(\cdot, k))^2}{b^{L_0}(\cdot, k)} * \rho_\varepsilon(x) \\
 &= \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \left(\frac{1}{4k^2} \int_0^{\bar{\lambda}} \frac{(b_{,x}^{L_0}(z, k))^2}{b^{L_0}(z, k)} \rho_\varepsilon(x - z) dz \right) \\
 &\leq \|\rho_\varepsilon\|_\infty \int_0^{\bar{\lambda}} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{4k^2} \frac{(b_{,x}^{L_0}(x, k))^2}{b^{L_0}(x, k)} dx \leq \frac{1}{\varepsilon} \mathcal{F}_\infty^\lambda(\mu^{L_0}) \lesssim \frac{1}{\varepsilon}.
 \end{aligned} \tag{5.40}$$

Thus, combining (5.39) with (5.40) we infer (5.24).

Step 5: in this step we show (5.25). By recalling Definition 2.1 we have

$$\hat{\mu}^L := \mu^L(\hat{u}^L) = \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} a^L(x, k) \mathcal{L}^1 \llcorner (-1, 1) \times \delta_k,$$

and

$$\hat{\mu}_{,x}^L = \mu_{,x}^L(\hat{u}^L) = \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} a_{,x}^L(x, k) \mathcal{L}^1 \llcorner (-1, 1) \times \delta_k,$$

Let $\varphi \in C_c^\infty((-1, 1) \times \mathbb{R})$. Then

$$\int_{(-1, 1) \times \mathbb{R}} \varphi d\hat{\mu}^L = \int_{(-1, 1) \times \mathbb{R}} \varphi d\hat{\mu}^L - \int_{(0, 1) \times \mathbb{R}} \varphi d\mu^{L_0} + \int_{(0, 1) \times \mathbb{R}} \varphi d\mu^{L_0}.$$

By Lemma 5.3 we have that

$$\lim_{L \rightarrow \infty} \int_{(0, 1) \times \mathbb{R}} \varphi d\mu^{L_0} = \int_{(0, 1) \times \mathbb{R}} \varphi d\mu;$$

hence, it suffices to show that

$$\lim_{L \rightarrow +\infty} \left(\int_{(-1, 1) \times \mathbb{R}} \varphi d\hat{\mu}^L - \int_{(0, 1) \times \mathbb{R}} \varphi d\mu^{L_0} \right) = 0.$$

Indeed, by (5.32) and (5.8) we have

$$\begin{aligned} \left| \int_{(-1,1) \times \mathbb{R}} \varphi \, d\hat{\mu}^L - \int_{(0,1) \times \mathbb{R}} \varphi \, d\mu^{L_0} \right| &\leq \|\varphi\|_\infty \left| \int_{(-1,1) \times \mathbb{R}} d\hat{\mu}^L - \int_{(0,1) \times \mathbb{R}} d\mu^{L_0} \right| \\ &= \|\varphi\|_\infty \left| \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} a^L(x, k) \, dx - \int_0^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^L(x, k) \, dx \right| \\ &= \|\varphi\|_\infty \left| \int_{-1}^1 \varepsilon \left(e^{-\frac{|x|}{\varepsilon}} - e^{-\frac{x-\bar{\lambda}}{\varepsilon}} \right) dx \right| \leq C\varepsilon^2 \rightarrow 0 \quad \text{as } L \rightarrow +\infty. \end{aligned}$$

Moreover, we have

$$\int_{(-1,1) \times \mathbb{R}} \varphi \, d\hat{\mu}_{,x}^L = - \int_{(-1,1) \times \mathbb{R}} \varphi_{,x} \, d\hat{\mu}^L \rightarrow - \int_{(-1,1) \times \mathbb{R}} \varphi_{,x} \, d\mu = \int_{(-1,1) \times \mathbb{R}} \varphi \, d\mu_{,x}.$$

Step 6: in this step we show (5.26). By (3.3), (5.30) we have

$$\begin{aligned} \int_{-L_0}^{L_0} \int_{-1}^1 (\hat{a}_{,yy}^L)^2 \, dx \, dy &= \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} a^L(x, k) k^2 \, dx \\ &= \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} b^{L_0}(\cdot, k) * \rho_\varepsilon(x) k^2 \, dx \\ &= \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \left(k^2 \int_0^{\bar{\lambda}} \bar{b}^{L_0}(z, k) \rho_\varepsilon(x-z) \, dz \right) dx \\ &\quad + \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(\bar{\lambda}, k) k^2 \int_{-1}^1 \int_{\bar{\lambda}}^{+\infty} \rho_\varepsilon(x-z) \, dz. \end{aligned} \quad (5.41)$$

Fubini's theorem yields

$$\int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \left(k^2 \int_0^{\bar{\lambda}} \bar{b}^{L_0}(z, k) \rho_\varepsilon(x-z) \, dz \right) dx \leq \int_0^{\bar{\lambda}} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(x, k) k^2 \, dx. \quad (5.42)$$

while from (5.29) we deduce

$$\begin{aligned} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(\bar{\lambda}, k) k^2 \int_{-1}^1 \int_{\bar{\lambda}}^{+\infty} \rho_\varepsilon(x-z) \, dz \, dx \\ \leq \frac{\varepsilon}{2} \left(e^{-\frac{1-\bar{\lambda}}{\varepsilon}} - e^{-\frac{-1-\bar{\lambda}}{\varepsilon}} \right) \int_{\frac{\bar{\lambda}+1}{2}}^{\bar{\lambda}} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \bar{b}^{L_0}(x, k) k^2 \, dx. \end{aligned} \quad (5.43)$$

Analogously from (3.3), (3.4) and Jensen's inequality, it holds

$$\begin{aligned} \int_{-L_0}^{L_0} \int_{-1}^1 (\hat{u}_{,x}^L)^2 dx dy &= \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{4k^2} \frac{(a_{,x}^L(x, k))^2}{a^L(x, k)} dx \\ &\leq \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{4k^2} \frac{(b_{,x}^{L_0}(\cdot, k))^2}{b^{L_0}(\cdot, k)} * \rho_\varepsilon(x) dx \quad (5.44) \\ &\leq \int_0^{\bar{\lambda}} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{4k^2} \frac{(\bar{b}_{,x}^{L_0}(x, k))^2}{\bar{b}^{L_0}(x, k)} dx. \end{aligned}$$

Gathering together (5.41)–(5.44), we obtain

$$\int_{-L}^L \int_{-1}^1 ((\hat{u}_{,x}^L)^2 + (\hat{u}_{,yy}^L)^2) dx dy \leq \left(1 + \frac{C\varepsilon(e^{\frac{1-\bar{\lambda}}{\varepsilon}} - e^{\frac{-1-\bar{\lambda}}{\varepsilon}})}{1-\lambda}\right) \mathcal{F}_\infty^\lambda(\mu^{L_0}), \quad (5.45)$$

and hence, by letting $L \rightarrow +\infty$ and recalling Lemma 5.3 (iii) we deduce (5.26).

Step 7: in this step we show (5.27). By (3.3), (3.4)

$$\begin{aligned} \int_{-L}^L \int_{-1}^1 (\hat{u}_{,xy}^L)^2 dx dy &= \int_{-L_0}^{L_0} \int_{-1}^1 (\hat{u}_{,xy}^L)^2 dx dy \\ &= \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{4} \frac{(a_{,x}^L(x, k))^2}{a^L(x, k)} dx. \quad (5.46) \end{aligned}$$

Next we observe that (5.31) yields

$$\begin{aligned} (a_{,x}^L(x, k))^2 &= (b^L(\cdot, k) * \dot{\rho}_\varepsilon(x))^2 \leq (b^L(\cdot, k) * |\dot{\rho}_\varepsilon|(x))^2 \\ &= \frac{1}{\varepsilon^2} (b^L(\cdot, k) * \rho_\varepsilon(x))^2 \leq \frac{1}{\varepsilon^2} (a^L(x, k))^2, \quad (5.47) \end{aligned}$$

so that combining (5.46) with (5.47) and recalling (5.32) we obtain

$$\int_{-L}^L \int_{-1}^1 (\hat{u}_{,xy}^L)^2 dx dy \leq \frac{1}{4\varepsilon^2} \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} (a^L(x, k))^2 dx \lesssim \frac{1}{\varepsilon^2}. \quad (5.48)$$

In a similar way (3.3) and (3.4) give

$$\begin{aligned}
 \oint_{-L}^L \int_{-1}^1 (\hat{u}_{,xx}^L)^2 dx dy &= \oint_{-L_0}^{L_0} \int_{-1}^1 (\hat{u}_{,xx}^L)^2 dx dy \\
 &= \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{k^2} \left[\left(\sqrt{a^L(x, k)} \right)_{,xx} \right]^2 dx \\
 &\lesssim \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{k^2} \frac{(a_{,xx}^L(x, k))^2}{a^L(x, k)} dx \\
 &\quad + \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{k^2} \frac{(a_{,x}^L(x, k))^4}{(a^L(x, k))^3} dx \quad (5.49) \\
 &\lesssim \frac{1}{\varepsilon^2} \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{1}{4k^2} \frac{(a_{,x}^L(x, k))^2}{a^L(x, k)} dx \\
 &\lesssim \frac{1}{\varepsilon^2} \int_{-1}^1 \oint_{-L_0}^{L_0} (\hat{u}_{,x}^L)^2 dy dx \\
 &\lesssim \frac{1}{\varepsilon^2} \mathcal{F}_\infty^\lambda(\mu^{L_0}) \lesssim \frac{1}{\varepsilon^2},
 \end{aligned}$$

where the second inequality follows from

$$(a_{,x}^L(x, k))^4 = (a_{,x}^L(x, k))^2 (b^L(\cdot, k) * \dot{\rho}_\varepsilon(x))^2 \leq \frac{1}{\varepsilon^2} (a_{,x}^L(x, k))^2 (a^L(x, k))^2,$$

and

$$(a_{,xx}^L(x, k))^2 = (b_{,x}^L(\cdot, k) * \dot{\rho}_\varepsilon(x))^2 \leq \frac{1}{\varepsilon^2} (a_{,x}^L(x, k))^2.$$

Moreover, appealing again to (5.47) we find

$$\begin{aligned}
 \oint_{-L}^L \int_{-1}^1 (\hat{u}_{,yyx}^L)^2 dx dy &= \oint_{-L_0}^{L_0} \int_{-1}^1 (\hat{u}_{,yyx}^L)^2 dx dy \\
 &= \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} k^2 \left[\left(\sqrt{a^L(x, k)} \right)_{,yx} \right]^2 dx \\
 &= \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} \frac{k^2}{4} \frac{(a_{,x}^L(x, k))^2}{a^L(x, k)} dx \\
 &\lesssim \frac{1}{\varepsilon^2} \int_{-1}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k \neq 0} k^2 a^L(x, k) dx
 \end{aligned}$$

$$\lesssim \frac{1}{\varepsilon^2} \int_{-1}^1 \int_{-L_0}^{L_0} (\hat{u}_{,yy}^L)^2 dy dx \lesssim \frac{1}{\varepsilon^2} \mathcal{F}_\infty^\lambda(\mu^{L_0}) \lesssim \frac{1}{\varepsilon^2}. \quad (5.50)$$

Eventually gathering together (5.48)–(5.50), we deduce (5.27).

Step 8: in this step we show (5.28). Analogously to step 3 we can find $y_0 \in [-L_0, L_0]$ such that

$$\hat{u}_{,x}^L(x, y_0) = \int_{-L_0}^{L_0} \hat{u}_{,x}^L(x, \hat{y}) d\hat{y} = 0,$$

so that by the fundamental theorem of calculus, Hölder's inequality it holds

$$\begin{aligned} |\hat{u}_{,x}^L(x, y)| &= \left| \int_{y_0}^y \hat{u}_{,xy}^L(x, y') dy' \right| \leq \sqrt{L_0} \left(\int_{-L_0}^{L_0} (\hat{u}_{,xy}^L)^2 dy' \right)^{\frac{1}{2}} \\ &= \sqrt{2} L_0 \left(\sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} \frac{1}{4} \frac{(a_{,x}^L(x, k))^2}{a^L(x, k)} \right)^{\frac{1}{2}} \quad (5.51) \\ &\lesssim L_0 \left(\frac{1}{4\varepsilon^2} \sum_{k \in \frac{\pi\mathbb{Z}}{L}, k \neq 0} a^L(x, k) \right)^{\frac{1}{2}} \lesssim \frac{L_0}{\varepsilon}, \end{aligned}$$

where the last two inequalities follow from (5.47) and (5.32). Therefore, (5.51) gives

$$\int_{-L_0}^{L_0} \int_{-1}^1 (\hat{u}_{,x}^L)^4 dx dy \lesssim \frac{L_0^2}{\varepsilon^2} \int_{-L_0}^{L_0} \int_{-1}^1 (\hat{u}_{,x}^L)^2 dx dy \lesssim \frac{L_0^2}{\varepsilon^2},$$

and the proof is concluded. $\square$

We are now in a position to prove Proposition 5.1.

Proof of Proposition 5.1 Let $\mu \in \mathcal{M}_\infty$ be as in the statement. Let $\varepsilon = \varepsilon(L) > 0$ and $n = n(L) \in \mathbb{N}$ be such that

$$\lim_{L \rightarrow +\infty} \varepsilon(L) = 0, \quad \lim_{L \rightarrow +\infty} n(L) = \lim_{L \rightarrow +\infty} \frac{L}{n(L)} = +\infty, \quad (5.52)$$

to be chosen later. Let $\hat{u}^L \in \mathcal{A}_L^{\text{out}} \cap \mathcal{A}_{L_0}^{\text{out}}$ with $L_0 := L/n(L)$ be the function given by Lemma 5.4. Recall that

$$A^L(x) = \frac{1}{2} \int_{-L}^L (\hat{u}_{,y}^L)^2 dy \quad \text{and} \quad f^L(x) = \sqrt{\frac{x}{A^L(x)}} \quad \text{for } x \geq 0.$$

Furthermore we let $M = M(L) \in \mathbb{N}$, $M \geq 2$ to be chosen later such that setting $\delta = \delta(L) := \frac{\varepsilon(L)}{M(L)} < \varepsilon(L)$ we have

$$\lim_{L \rightarrow +\infty} M(L) = +\infty, \quad \text{and} \quad \lim_{L \rightarrow +\infty} \delta(L) = \lim_{L \rightarrow +\infty} M^2(L) \delta(L) = 0. \quad (5.53)$$

We consider $\psi_\delta \in C^\infty(\mathbb{R})$ such that

$$\psi_\delta \equiv 0 \text{ in } (-\infty, \delta], \quad \psi_\delta \equiv 1 \text{ in } [2\delta, +\infty), \quad |\dot{\psi}_\delta(x)| \leq C\delta^{-1}, \quad |\ddot{\psi}_\delta(x)| \leq C\delta^{-2}.$$

Note that $\dot{\psi}_\delta = \ddot{\psi}_\delta = 0$ in $(\delta, 2\delta)^c$. We next define $(w^L, u^L) = ((w_1^L, w_2^L), u^L)$ as follows:

$$\begin{aligned} u^L(x, y) &:= \psi_\delta(x) f^L(x) \hat{u}^L(x, y), \\ w_2^L(x, y) &:= \psi_\delta^2(x) xy + B^L(x) - \frac{1}{2} \int_0^y (u_{,y}^L)^2 dy', \\ w_1^L(x, y) &:= x - \frac{1}{L^2} \int_0^y (w_{2,x}^L + u_{,x}^L u_{,y}^L) dy', \end{aligned}$$

where

$$B^L(x) := \frac{1}{2} \int_{-L_0}^{L_0} \int_0^y (u_{,y}^L)^2 dy' dy - \int_{-L_0}^{L_0} \int_0^x u_{,x}^L u_{,y}^L dx' dy.$$

Clearly $u^L \in \mathcal{A}_L^{\text{out}} \cap \mathcal{A}_{L_0}^{\text{out}}$. We show that $w^L \in \mathcal{A}_L^{\text{in}} \cap \mathcal{A}_{L_0}^{\text{in}}$. Precisely to see that $w^L(x, \cdot)$ is $2L_0$ periodic we use the following fact:

A differentiable function h is T -periodic if h' is T -periodic and $h(t) = h(t + T)$ for some t .

The function $w_{2,y}^L(x, \cdot)$ is $2L_0$ -periodic, since $u_{,y}^L$ is, and from (3.3) satisfies

$$\begin{aligned} w_2^L(x, L_0) - w_2^L(x, -L_0) &= 2L_0 \psi_\delta^2(x) x - \int_{-L_0}^{L_0} (u_{,y}^L)^2 dy \\ &= 2L_0 \psi_\delta^2(x) (x - (f^L(x))^2 A^L(x)) = 0, \end{aligned}$$

from which we deduce $w_2^L(x, \cdot)$ is $2L_0$ -periodic. Using this periodicity, and in particular also of $w_{2,x}^L$, we see that $w_{1,y}^L(x, \cdot)$ is $2L_0$ -periodic. Moreover, we have

$$\begin{aligned} w_1^L(x, L_0) - w_1^L(x, -L_0) &= -\frac{1}{L^2} \int_{-L_0}^{L_0} (w_{2,x}^L + u_{,x}^L u_{,y}^L) dy \\ &= -\frac{1}{L^2} \left(2L_0 \dot{B}^L(x) - \int_{-L_0}^{L_0} \int_0^y u_{,y}^L u_{,xy}^L dy' dy + \int_{-L_0}^{L_0} u_{,x}^L u_{,y}^L dy \right) = 0, \end{aligned}$$

where the second and the third equalities follow from

$$w_{2,x}^L = (\psi_\delta^2(x)x)'y + \dot{B}^L(x) - \int_0^y u_{,y}^L u_{,xy}^L dy',$$

and

$$2L_0 \dot{B}^L(x) = 2L_0 \left(\int_{-L_0}^{L_0} u_{,y}^L u_{,xy}^L dy - \int_{-L_0}^{L_0} u_{,x}^L u_{,y}^L dy \right).$$

Thus, we deduce that $w_{1,y}^L$ is $2L_0$ -periodic. For the reader convenience we divide the rest of the proof into several steps. We will repeatedly use that the averaged integral over $(-L, L)$ of a $2L_0$ -periodic function is equal to the averaged integral over $(-L_0, L_0)$ of the same function.

Step 1: we show that (w^L, u^L) converges to μ in the sense of Definition 2.3. We have that

$$\hat{u}^L(x, y) = \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k > 0} a_k^L(x) \sin(ky) + \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k < 0} a_k^L(x) \cos(ky),$$

so that

$$\begin{aligned} u^L(x, y) &= \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k > 0} \psi_\delta(x) f^L(x) a_k^L(x) \sin(ky) \\ &\quad + \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}, k < 0} \psi_\delta(x) f^L(x) a_k^L(x) \cos(ky). \end{aligned}$$

Therefore, we get

$$\begin{aligned} \mu^L := \mu^L(u^L) &= \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \psi_\delta^2(x) (f^L(x))^2 (a_k^L(x))^2 k^2 \mathcal{L}^1 \lfloor (-1, 1) \times \delta_k \\ &= \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \psi_\delta^2(x) (f^L(x))^2 (a_k^L(x))^2 k^2 \mathcal{L}^1 \lfloor (\delta, 1) \times \delta_k. \end{aligned}$$

We show $\mu^L \xrightarrow{*} \mu$. We fix $\varphi \in C_c^\infty((-1, 1) \times \mathbb{R})$ and we write

$$\begin{aligned} \int_{(-1,1) \times \mathbb{R}} \varphi d\mu^L &= \left(\int_{(-1,1) \times \mathbb{R}} \varphi d\mu^L - \int_{(-1,1) \times \mathbb{R}} \varphi d\mu^L(\hat{u}^L) \right) \\ &\quad + \int_{(-1,1) \times \mathbb{R}} \varphi d\mu^L(\hat{u}^L). \end{aligned}$$

By Lemma 5.4 it holds

$$\lim_{L \rightarrow +\infty} \int_{(-1,1) \times \mathbb{R}} \varphi d\mu^L(\hat{u}^L) = \int_{(-1,1) \times \mathbb{R}} \varphi d\mu.$$

Therefore, it is sufficient to show that

$$\lim_{L \rightarrow +\infty} \left(\int_{(-1,1) \times \mathbb{R}} \varphi \, d\mu^L - \int_{(-1,1) \times \mathbb{R}} \varphi \, d\mu^L(\hat{u}^L) \right) = 0.$$

Recalling that $\psi_\delta = 0$ in $(-1, \delta)$, $\psi_\delta = 1$ in $(2\delta, 1)$ and $M\varepsilon = M\delta^2 \geq 2\delta$, we have

$$\begin{aligned} & \left| \int_{(-1,1) \times \mathbb{R}} \varphi \, d\mu^L - \int_{(0,1) \times \mathbb{R}} \varphi \, d\mu^L(\hat{u}^L) \right| \\ &= \left| \int_0^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} \varphi(x, k) (a_k^L(x))^2 k^2 \left(\psi_\delta^2(x) (f^L(x))^2 - 1 \right) dx \right| \\ &\leq \|\varphi\|_\infty \left| \int_0^{M\varepsilon} \left(\psi_\delta^2(x) (f^L(x))^2 - 1 \right) \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} (a_k^L(x))^2 k^2 dx \right| \\ &\quad + \|\varphi\|_\infty \left| \int_{M\varepsilon}^1 \left((f^L(x))^2 - 1 \right) \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} (a_k^L(x))^2 k^2 dx \right|. \end{aligned}$$

Since

$$\sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} (a_k^L(x))^2 k^2 = A^L(x) = \frac{1}{2} \int_{-L}^L (\hat{u}_{\cdot, y}(x, \cdot))^2 dy,$$

by (5.18) and (5.19) we have

$$\begin{aligned} & \left(\psi_\delta^2(x) (f^L(x))^2 - 1 \right) \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} (a_k^L(x))^2 k^2 \\ &\leq \left(\psi_\delta^2(x) (1 + o_L(1)) - 1 \right) \max\{x, \varepsilon\} \end{aligned}$$

which together with (5.53) implies

$$\begin{aligned} & \|\varphi\|_\infty \left| \int_0^{M\varepsilon} \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} (a_k^L(x))^2 k^2 (\psi_\delta^2(x) (f^L(x))^2 - 1) dx \right| \\ &\leq CM\varepsilon = CM^2\delta \rightarrow 0 \quad \text{as } L \rightarrow +\infty. \end{aligned}$$

Whereas (5.19), (5.20) with $N = M$ and the fact that $M = M(L) \rightarrow +\infty$ imply

$$(f^L(x))^2 = 1 + o_L(1) \quad \text{in } [M\varepsilon, 1),$$

so that

$$\|\varphi\|_\infty \left| \int_{M\varepsilon}^1 \sum_{k \in \frac{\pi\mathbb{Z}}{L_0}} a_k^L(x) k^2 (f^L(x) - 1) dx \right| \leq o_L(1) \rightarrow 0 \quad \text{as } L \rightarrow +\infty.$$

Eventually by duality we have

$$\int_{\mathbb{R} \times (-1,1)} \varphi d\mu_{,x}^L = - \int_{\mathbb{R} \times (-1,1)} \varphi_{,x} d\mu^L \rightarrow - \int_{\mathbb{R} \times (-1,1)} \varphi_{,x} d\mu = \int_{\mathbb{R} \times (-1,1)} \varphi d\mu_{,x},$$

which in turn implies $\mu_{,x}^L \xrightarrow{*} \mu_{,x}$.

Step 2: we show that

$$\limsup_{L \rightarrow +\infty} \int_{-L}^L \int_{-1}^1 ((u_{,x}^L)^2 + (u_{,yy}^L)^2) dx dy \leq \mathcal{F}_\infty(\mu) + C \lim_{L \rightarrow +\infty} \omega(2M^2\delta) \log M. \quad (5.54)$$

To this purpose we note that

$$u_{,x}^L(x, y) = \psi_\delta(x) f^L(x) \hat{u}_{,x}^L(x, y) + \dot{\psi}_\delta(x) f^L(x) \hat{u}^L(x, y) + \psi_\delta(x) \dot{f}^L(x) \hat{u}^L(x, y). \quad (5.55)$$

Therefore, by Young's inequality

$$\begin{aligned} \int_{-L}^L \int_{-1}^1 (u_{,x}^L)^2 dx dy &\leq (1 + \alpha) \int_{-L}^L \int_{\delta}^1 (f^L(x))^2 (\hat{u}_{,x}^L)^2 dx dy \\ &\quad + 2(1 + \alpha^{-1}) \frac{1}{\delta^2} \int_{-L}^L \int_{\delta}^{2\delta} (f^L(x))^2 (\hat{u}^L)^2 dx dy \\ &\quad + 2(1 + \alpha^{-1}) \int_{-L}^L \int_{\delta}^1 (\dot{f}^L(x))^2 (\hat{u}^L)^2 dx dy, \end{aligned} \quad (5.56)$$

for any $\alpha > 0$. In this way by recalling (5.19) (5.22) and (5.21) we have

$$\int_{-L}^L \int_{-1}^1 (f^L(x))^2 (\hat{u}_{,x}^L)^2 dx dy \leq (1 + o_L(1)) \int_{-L}^L \int_{-1}^1 (\hat{u}_{,x}^L)^2 dx dy, \quad (5.57)$$

$$\begin{aligned} \frac{1}{\delta^2} \int_{-L}^L \int_{\delta}^{2\delta} (f^L(x))^2 (\hat{u}^L)^2 dx dy &\lesssim \frac{1}{\delta^2} \delta(\omega(2M\varepsilon) + Me^{-M}) \int_{\delta}^{2\delta} dx \\ &\lesssim (\omega(2M\varepsilon) + Me^{-M}), \end{aligned} \quad (5.58)$$

and

$$\begin{aligned}
 \int_{-L}^L \int_{\delta}^1 (\dot{f}^L(x))^2 (\hat{u}^L)^2 \, dx \, dy &\lesssim \int_{\delta}^{\varepsilon} \frac{1}{x\varepsilon} \varepsilon (\omega(2M\varepsilon) + Me^{-M}) \, dx \\
 &\quad + \int_{\varepsilon}^{\sqrt{\varepsilon}} \frac{e^{-\frac{x}{\varepsilon}}}{x\varepsilon} x (\omega(2M\varepsilon) + Me^{-M}) \, dx \\
 &\quad + \int_{\sqrt{\varepsilon}}^1 \frac{e^{-\frac{1}{\sqrt{\varepsilon}}}}{x\varepsilon} x \, dx \\
 &\lesssim (\log \frac{\varepsilon}{\delta} + 1) (\omega(2M\varepsilon) + Me^{-M}) \\
 &\quad + \frac{e^{-\frac{1}{\sqrt{\varepsilon}}}}{\varepsilon}.
 \end{aligned} \tag{5.59}$$

Gathering together (5.56)–(5.59) and recalling that $\delta = \varepsilon/M$, we deduce

$$\begin{aligned}
 \int_{-L}^L \int_{-1}^1 (u_{,x}^L)^2 \, dx \, dy &\leq (1 + \bar{\alpha}) \int_{-L}^L \int_{-1}^1 (\hat{u}_{,x}^L)^2 \, dx \, dy \\
 &\quad + C(\log M + 1) (\omega(2M^2\delta) + Me^{-M}) + o_L(1),
 \end{aligned} \tag{5.60}$$

with $\bar{\alpha} := \alpha + \alpha o_L(1) + o_L(1)$. Moreover, as $u_{,yy}^L = \psi_{\delta}(x) f^L(x) \hat{u}_{,yy}^L$ we get

$$\int_{-L}^L \int_{-1}^1 (u_{,yy}^L)^2 \, dx \, dy \leq \int_{-L}^L \int_{-1}^1 (f^L(yy))^2 (\hat{u}_{,yy}^L)^2 \, dx \, dy \leq \int_{-L}^L \int_{-1}^1 (\hat{u}_{,x}^L)^2 \, dx \, dy. \tag{5.61}$$

By (5.60), (5.61), (5.26) and the fact that $M^2\delta \rightarrow 0$ we finally deduce

$$\begin{aligned}
 \limsup_{L \rightarrow +\infty} \int_{-L}^L \int_{-1}^1 ((u_{,x}^L)^2 + (u_{,yy}^L)^2) \, dx \, dy &\leq (1 + \alpha) \mathcal{F}_{\infty}(\mu) \\
 &\quad + C \lim_{L \rightarrow +\infty} \omega(2M^2\delta) \log M.
 \end{aligned}$$

Eventually by the arbitrariness of α we infer the desired estimate.

Step 3: we show that

$$L^2 \int_{-L}^L \int_{-1}^1 \left(w_{1,x}^L + \frac{(u_{,x}^L)^2}{2L^2} - 1 \right)^2 \, dx \, dy \lesssim \frac{L_0^4}{L^2} \frac{1}{\delta^2 \varepsilon} \lesssim \frac{L_0^4}{L^2} \frac{1}{\delta^3 M}. \tag{5.62}$$

By Young's inequality we have

$$\begin{aligned} L^2 \int_{-L}^L \int_{-1}^1 \left(w_{1,x}^L + \frac{(u_{1,x}^L)^2}{2L^2} - 1 \right)^2 dx dy &\lesssim \int_{-L}^L \int_{-1}^1 \frac{(u_{1,x}^L)^4}{L^2} dx dy \\ &+ L^2 \int_{-L}^L \int_{-1}^1 (w_{1,x}^L - 1)^2 dx dy. \end{aligned} \quad (5.63)$$

We estimate the first term on the right-hand side of (5.63). By (5.55) it follows

$$\begin{aligned} \int_{-L}^L \int_{-1}^1 (u_{1,x}^L)^4 dx dy &\lesssim \int_{-L}^L \int_{-1}^1 (f^L(x))^4 (\hat{u}_{1,x}^L)^4 dx dy \\ &+ \frac{1}{\delta^4} \int_{-L}^L \int_{\delta}^{2\delta} (f^L(x))^4 (\hat{u}^L)^4 dx dy + \int_{-L}^L \int_{\delta}^1 (f^L(x))^4 (\hat{u}^L)^4 dx dy. \end{aligned} \quad (5.64)$$

By (5.19) and (5.28) we have

$$\int_{-L}^L \int_{-1}^1 (f^L(x))^4 (\hat{u}_{1,x}^L)^4 dx dy \lesssim \frac{L_0^2}{\varepsilon^2}, \quad (5.65)$$

whereas from (5.19), (5.23), and the fact that $x \in (\delta, 2\delta)$ we get

$$\frac{1}{\delta^4} \int_{-L}^L \int_{\delta}^{2\delta} (f^L(x))^4 (\hat{u}^L)^4 dx dy \lesssim \frac{1}{\delta^4} \frac{\delta^2}{\varepsilon^2} L_0^2 \int_{\delta}^{2\delta} (\max\{x, \varepsilon\})^2 dx \lesssim \frac{L_0^2}{\delta}. \quad (5.66)$$

Finally, by (5.21) and (5.23)

$$\int_{-L}^L \int_{\delta}^1 (f^L(x))^4 (\hat{u}^L)^4 dx dy \lesssim L_0^2 \int_{\delta}^1 \frac{1}{x^2 \varepsilon^2} (\max\{x, \varepsilon\})^2 dx \lesssim L_0^2 \left(\frac{1}{\delta} + \frac{1}{\varepsilon^2} \right). \quad (5.67)$$

Thus, gathering together (5.64)–(5.67) we infer

$$\int_{-L}^L \int_{-1}^1 \frac{(u_{1,x}^L)^4}{L^2} dx dy \lesssim \frac{L_0^2}{L^2} \left(\frac{1}{\delta} + \frac{1}{\varepsilon^2} \right). \quad (5.68)$$

We now pass to estimate the second term on the right-hand side of (5.63). To this aim we observe that integrating by parts it holds

$$\begin{aligned} w_{2,x}^L + u_{1,x}^L u_{1,y}^L &= (x \psi_{\delta}^2(x))' y + \dot{B}^L(x) - \int_0^y u_{1,y}^L u_{1,x,y}^L dy' + u_{1,x}^L u_{1,y}^L(x, y) \\ &= (x \psi_{\delta}^2(x))' y + \dot{B}^L(x) + \int_0^y u_{1,yy} u_{1,x} dy' + u_{1,x}^L u_{1,y}^L|_{y=0} \quad (5.69) \\ &= (x \psi_{\delta}^2(x))' y + \int_0^y u_{1,yy} u_{1,x} dy' + C^L(x), \end{aligned}$$

where

$$C^L(x) := - \oint_{-L}^L \int_0^y u_{,yy} u_{,x} \, dy' \, dy = - \oint_{-L_0}^{L_0} \int_0^y u_{,yy} u_{,x} \, dy' \, dy,$$

and the last equality is a consequence of the following identity

$$\dot{B}^L(x) + u_{,x}^L u_{,y}^L|_{y=0} = \oint_{-L}^L \left(\int_0^y u_{,y}^L u_{,xy}^L \, dy' - u_{,x}^L u_{,y}^L + u_{,x}^L u_{,y}^L|_{y=0} \right) dy = C^L(x).$$

Using (5.69) and the definition of ω , we get

$$\begin{aligned} w_{1,x}^L - 1 &= -\frac{1}{L^2} \int_0^y \left((x\psi_\delta^2(x))'' y' + \int_0^{y'} (u_{,yy}^L u_{,xx}^L + u_{,yyx}^L u_{,x}^L) \, dy'' + \dot{C}^L(x) \right) dy' \\ &= -\frac{1}{L^2} \left((x\psi_\delta^2(x))'' \frac{y^2}{2} + \int_0^y \int_0^{y'} (u_{,yy}^L u_{,xx}^L + u_{,yyx}^L u_{,x}^L) \, dy'' \, dy' + \dot{C}^L(x)y \right). \end{aligned}$$

This together with Young's inequality gives

$$\begin{aligned} L^2 \oint_{-L}^L \int_{-1}^1 (w_{1,x}^L - 1)^2 \, dx \, dy &= L^2 \oint_{-L_0}^{L_0} \int_{-1}^1 (w_{1,x}^L - 1)^2 \, dx \, dy \\ &\lesssim \frac{1}{L^2} \oint_{-L_0}^{L_0} \int_{-1}^1 ((x\psi_\delta^2(x))'')^2 y^4 \, dx \, dy \\ &\quad + \frac{1}{L^2} \oint_{-L_0}^{L_0} \int_{-1}^1 \left(\int_0^y \int_0^{y'} (u_{,yy}^L u_{,xx}^L + u_{,yyx}^L u_{,x}^L) \, dy'' \, dy' \right)^2 \, dx \, dy \\ &\quad + \frac{1}{L^2} \oint_{-L_0}^{L_0} \int_{-1}^1 (\dot{C}^L(x))^2 y^2 \, dx \, dy. \end{aligned} \quad (5.70)$$

As $(x\psi_\delta^2(x))'' = 4\psi_\delta(x)\dot{\psi}_\delta(x) + 2x\psi_\delta(x)\ddot{\psi}_\delta(x) + 2x(\dot{\psi}_\delta(x))^2$ in $(\delta, 2\delta)$ and $(x\psi_\delta^2(x))'' = 0$; otherwise, we have

$$\frac{1}{L^2} \oint_{-L_0}^{L_0} \int_{-1}^1 ((x\psi_\delta^2(x))'')^2 y^4 \, dx \, dy \lesssim \frac{L_0^4}{L^2} \int_\delta^{2\delta} ((x\psi_\delta^2(x))'')^2 \, dx \lesssim \frac{L_0^4}{L^2} \frac{1}{\delta}. \quad (5.71)$$

We now estimate the second term on the right-hand side of (5.70). We first observe that if a, b, c, d are $2L_0$ -periodic then by applying in order Hölder, Young and Jensen

inequalities we have

$$\begin{aligned}
 & \left[\int_0^y \int_0^{y'} (ab + cd) \, dy'' \, dy' \right]^2 \\
 & \leq \left[\int_0^y \|a\|_{L^2(0,y')} \|b\|_{L^2(0,y')} + \|c\|_{L^2(0,y')} \|d\|_{L^2(0,y')} \, dy' \right]^2 \\
 & \lesssim \left[\int_0^y \|a\|_{L^2(0,y')} \|b\|_{L^2(0,y')} \, dy' \right]^2 + \left[\int_0^y \|c\|_{L^2(0,y')} \|d\|_{L^2(0,y')} \, dy' \right]^2 \\
 & \lesssim L_0^2 \int_{-L_0}^{L_0} \|a\|_{L^2(0,y')}^2 \|b\|_{L^2(0,y')}^2 \, dy' + L_0^2 \int_{-L_0}^{L_0} \|c\|_{L^2(0,y')}^2 \|d\|_{L^2(0,y')}^2 \, dy' \\
 & \lesssim L_0^4 \left[\left(\int_{-L_0}^{L_0} a^2 \, dy'' \right) \left(\int_{-L_0}^{L_0} b^2 \, dy'' \right) + \left(\int_{-L_0}^{L_0} c^2 \, dy'' \right) \left(\int_{-L_0}^{L_0} d^2 \, dy'' \right) \right].
 \end{aligned}$$

Therefore, it follows that

$$\begin{aligned}
 & \frac{1}{L^2} \int_{-L_0}^{L_0} \int_{-1}^1 \left(\int_0^y \int_0^{y'} (u_{,yy}^L u_{,xx}^L + u_{,yyx}^L u_{,x}^L) \, dy'' \, dy' \right)^2 \, dx \, dy \\
 & = \frac{1}{L^2} \int_{-L_0}^{L_0} \int_{\delta}^1 \left(\int_0^y \int_0^{y'} (u_{,yy}^L u_{,xx}^L + u_{,yyx}^L u_{,x}^L) \, dy'' \, dy' \right)^2 \, dx \, dy \\
 & \lesssim \frac{L_0^4}{L^2} \int_{\delta}^1 \left(\int_{-L_0}^{L_0} (u_{,yy}^L)^2 \, dy'' \right) \left(\int_{-L_0}^{L_0} (u_{,xx}^L)^2 \, dy'' \right) \, dx \\
 & \quad + \frac{L_0^4}{L^2} \int_{\delta}^1 \left(\int_{-L_0}^{L_0} (u_{,yyx}^L)^2 \, dy'' \right) \left(\int_{-L_0}^{L_0} (u_{,x}^L)^2 \, dy'' \right) \, dx. \tag{5.72}
 \end{aligned}$$

Next we show separately that:

$$\int_{-L_0}^{L_0} (u_{,yy}^L)^2 \, dy'' \lesssim \frac{1}{\varepsilon}, \quad \int_{-L_0}^{L_0} (u_{,x}^L)^2 \, dy'' \lesssim \frac{1}{x\varepsilon} \max\{x, \varepsilon\}, \tag{5.73}$$

$$\int_{\delta}^1 \int_{-L_0}^{L_0} (u_{,xx}^L)^2 \, dy'' \, dx \lesssim \frac{1}{\delta^2}, \quad \int_{\delta}^1 \int_{-L_0}^{L_0} (u_{,yyx}^L)^2 \, dy'' \, dx \lesssim \frac{1}{\delta\varepsilon}. \tag{5.74}$$

Since $u_{,yy}^L = \psi_{\delta}(x) f^L(x) \hat{u}_{,yy}^L$ by (5.19) and (5.24), we have

$$\int_{-L_0}^{L_0} (\hat{u}_{,yy}^L)^2 \, dy \lesssim \frac{1}{\varepsilon}.$$

By (5.55), (5.19), (5.24), (5.22) and (5.21), we have

$$\begin{aligned}
 \int_{-L_0}^{L_0} (u_{,x}^L)^2 dy &\lesssim \int_{-L_0}^{L_0} (f^L(x))^2 (\hat{u}_{,x}^L)^2 dy + \frac{1}{\delta^2} \int_{-L_0}^{L_0} \chi_{(\delta, 2\delta)} (f^L(x))^2 (\hat{u}^L)^2 dy \\
 &\quad + \int_{-L_0}^{L_0} (\dot{f}^L(x))^2 (\hat{u}^L)^2 dy \\
 &\lesssim \frac{1}{\varepsilon} + \frac{1}{\delta^2} \frac{\delta}{\varepsilon} \max\{x, \varepsilon\} \chi_{(\delta, 2\delta)} + \frac{1}{x\varepsilon} \max\{x, \varepsilon\} \\
 &\lesssim \frac{1}{\varepsilon} + \frac{1}{\delta} \chi_{(\delta, 2\delta)} + \frac{1}{x\varepsilon} \max\{x, \varepsilon\} \lesssim \frac{1}{x\varepsilon} \max\{x, \varepsilon\}.
 \end{aligned}$$

From (5.55) it follows

$$\begin{aligned}
 u_{,xx}^L(x, y) &= \psi_\delta(x) f^L(x) \hat{u}_{,xx}^L(x, y) + \psi_\delta(x) \ddot{f}^L(x) \hat{u}^L(x, y) \\
 &\quad + \ddot{\psi}_\delta(x) f^L(x) \hat{u}^L(x, y) + 2\dot{\psi}_\delta(x) f^L(x) \hat{u}_{,x}^L(x, y) \\
 &\quad + 2\dot{\psi}_\delta(x) \dot{f}^L(x) \hat{u}^L(x, y) + 2\psi_\delta(x) \dot{f}^L(x) \hat{u}_{,x}^L(x, y).
 \end{aligned}$$

Hence, by (5.19), (5.27), (5.21), (5.22) and (5.26) we have

$$\begin{aligned}
 \int_\delta^1 \int_{-L_0}^{L_0} (u_{,xx}^L)^2 dy dx &\lesssim \int_\delta^1 \int_{-L_0}^{L_0} (f^L(x))^2 (\hat{u}_{,xx}^L)^2 dy dx \\
 &\quad + \int_\delta^1 \int_{-L_0}^{L_0} (\ddot{f}^L(x))^2 (\hat{u}^L)^2 dy dx \\
 &\quad + \int_\delta^1 \int_{-L_0}^{L_0} (\dot{f}^L(x))^2 (\hat{u}_{,x}^L)^2 dy dx \\
 &\quad + \frac{1}{\delta^4} \int_\delta^{2\delta} \int_{-L_0}^{L_0} (f^L(x))^2 (\hat{u}^L)^2 dy dx \\
 &\quad + \frac{1}{\delta^2} \int_\delta^{2\delta} \int_{-L_0}^{L_0} (\dot{f}^L(x))^2 (\hat{u}^L)^2 dy dx \\
 &\quad + \frac{1}{\delta^2} \int_\delta^{2\delta} \int_{-L_0}^{L_0} (f^L(x))^2 (\hat{u}_{,x}^L)^2 dy dx \\
 &\lesssim \frac{1}{\varepsilon^2} + \int_\delta^1 \frac{1}{x^3 \varepsilon} \max\{x, \varepsilon\} dx + \frac{1}{\delta \varepsilon} \\
 &\quad + \frac{1}{\delta^4} \int_\delta^{2\delta} \frac{x}{\varepsilon} \max\{x, \varepsilon\} dx \\
 &\quad + \frac{1}{\delta^2} \int_\delta^{2\delta} \frac{1}{x \varepsilon} \max\{x, \varepsilon\} dx + \frac{1}{\delta^2} \\
 &\lesssim \frac{1}{\varepsilon^2} + \frac{1}{\delta^2} + \frac{1}{\delta \varepsilon} \lesssim \frac{1}{\delta^2}.
 \end{aligned} \tag{5.75}$$

Analogously by (5.55) it follows

$$\begin{aligned} u_{,yyx}^L(x, y) &= \psi_\delta(x) f^L(x) \hat{u}_{,yyx}^L(x, y) + \dot{\psi}_\delta(x) f^L(x) \hat{u}_{,yy}^L(x, y) \\ &\quad + \psi_\delta(x) \dot{f}^L(x) \hat{u}_{,yy}^L(x, y), \end{aligned}$$

from which together with (5.27), (5.19), (5.24) and (5.21)

$$\begin{aligned} \int_\delta^1 \int_{-L_0}^{L_0} (u_{,yyx}^L)^2 dy dx &\lesssim \int_\delta^1 \int_{-L_0}^{L_0} (f^L(x))^2 (\hat{u}_{,yyx}^L)^2 dy dx \\ &\quad + \frac{1}{\delta^2} \int_\delta^{2\delta} \int_{-L_0}^{L_0} (f^L(x))^2 (\hat{u}_{,yy}^L)^2 dy dx \\ &\quad + \int_\delta^1 \int_{-L_0}^{L_0} (\dot{f}^L(x))^2 (\hat{u}_{,yy}^L)^2 dy dx \\ &\lesssim \frac{1}{\varepsilon^2} + \frac{1}{\delta^2} \frac{\delta}{\varepsilon} \frac{\delta}{\varepsilon} + \frac{1}{\delta \varepsilon} \lesssim \frac{1}{\delta \varepsilon}. \end{aligned}$$

Now (5.73) and (5.74) yield

$$\frac{L_0^4}{L^2} \int_\delta^1 \left(\int_{-L_0}^{L_0} (u_{,yy}^L)^2 dy'' \right) \left(\int_{-L_0}^{L_0} (u_{,xx}^L)^2 dy'' \right) dx \lesssim \frac{L_0^4}{L^2} \frac{1}{\delta^2 \varepsilon}, \quad (5.76)$$

and

$$\begin{aligned} \frac{L_0^4}{L^2} \int_\delta^1 \left(\int_{-L_0}^{L_0} (u_{,yyx}^L)^2 dy'' \right) \left(\int_{-L_0}^{L_0} (u_{,x}^L)^2 dy'' \right) dx \\ \lesssim \frac{L_0^4}{L^2} \left(\frac{1}{\delta} + \frac{1}{\varepsilon} \right) \frac{1}{\delta \varepsilon} \lesssim \frac{L_0^4}{L^2} \frac{1}{\delta^2 \varepsilon}. \end{aligned} \quad (5.77)$$

Gathering together (5.72), (5.76) and (5.77), we find

$$\frac{1}{L^2} \int_{-L_0}^{L_0} \int_{-1}^1 \left(\int_0^y \int_0^{y'} (u_{,yy}^L u_{,xx}^L + u_{,yyx}^L u_{,x}^L) dy'' dy' \right)^2 dx dy \lesssim \frac{L_0^4}{L^2} \frac{1}{\delta^2 \varepsilon}. \quad (5.78)$$

It remains to estimate the third term on the right-hand side of (5.70). In a similar way, see in particular (5.76) and (5.77), we have

$$\begin{aligned}
 & \frac{1}{L^2} \int_{-L_0}^{L_0} \int_{-1}^1 (\dot{C}^L(x))^2 y^2 \, dx \, dy \\
 & \lesssim \frac{L_0^2}{L^2} \int_{-1}^1 \left(\int_{-L_0}^{L_0} \int_0^y (u_{,yy}^L u_{,xx}^L + u_{,yyx}^L u_{,x}^L) \, dy' \, dy \right)^2 \, dx \\
 & \lesssim \frac{L_0^4}{L^2} \int_{-1}^1 \int_{-L_0}^{L_0} \left(\int_{-L_0}^{L_0} (u_{,yy}^L)^2 \, dy'' \right) \left(\int_{-L_0}^{L_0} (u_{,xx}^L)^2 \, dy'' \right) \, dy \, dx \quad (5.79) \\
 & \quad + \frac{L_0^4}{L^2} \int_{-1}^1 \int_{-L_0}^{L_0} \left(\int_{-L_0}^{L_0} (u_{,yyx}^L)^2 \, dy'' \right) \left(\int_{-L_0}^{L_0} (u_{,x}^L)^2 \, dy'' \right) \, dy \, dx \\
 & \lesssim \frac{L_0^4}{L^2} \frac{1}{\delta^2 \varepsilon}.
 \end{aligned}$$

Gathering together (5.70), (5.71), (5.78) and (5.79), we infer

$$L^2 \int_{-L}^L \int_{-1}^1 (w_{1,x}^L - 1)^2 \, dx \, dy \lesssim \frac{L_0^4}{L^2} \frac{1}{\delta^2 \varepsilon},$$

which together with (5.68) and (5.63) implies

$$L^2 \int_{-L}^L \int_{-1}^1 \left(w_{1,x}^L + \frac{(u_{,x}^L)^2}{2L^2} - 1 \right)^2 \, dx \, dy \lesssim \frac{L_0^2}{L^2} \left(\frac{1}{\delta} + \frac{1}{\varepsilon^2} \right) + \frac{L_0^4}{L^2} \frac{1}{\delta^2 \varepsilon} \lesssim \frac{L_0^4}{L^2} \frac{1}{\delta^2 \varepsilon}.$$

Step 4: we show that

$$L^2 \int_{-L}^L \int_{-1}^1 \left(w_{2,y}^L + \frac{(u_{,y}^L)^2}{2} - x \right)^2 \, dx \, dy \leq L^2 \left(\frac{1}{3} + C\delta^3 \right). \quad (5.80)$$

Recalling the definition of w_2^L it holds $w_{2,y}^L + \frac{(u_{,y}^L)^2}{2} - x = \psi_\delta^2(x)x - x$, so that

$$\begin{aligned}
 \int_{-L}^L \int_{-1}^1 \left(w_{2,y}^L + \frac{(u_{,y}^L)^2}{2} - x \right)^2 \, dx \, dy &= \int_{-1}^1 (\psi_\delta^2(x)x - x)^2 \, dx \\
 &\leq \int_{-1}^{2\delta} x^2 \, dx = \frac{8}{3}\delta^3 + \frac{1}{3}.
 \end{aligned}$$

Step 5: we show that

$$\int_{-L}^L \int_{-1}^1 \left(L^2 w_{1,y}^L + w_{2,x}^L + u_{,x}^L u_{,y}^L \right)^2 \, dx \, dy = 0. \quad (5.81)$$

This is a trivial consequence of the following identity

$$w_{1,y}^L = -\frac{1}{L^2}(w_{2,x}^L + u_{,x}^L u_{,y}^L),$$

which follows from the definitions of w_1^L and w_2^L .

Step 6: we show that

$$\frac{1}{L^2} \int_{-L}^L \int_{-1}^1 \left(2(u_{,xy}^L)^2 + \frac{1}{L^2} (u_{,xx}^L)^2 \right) dx dy \lesssim \left(\frac{1}{L^2} + \frac{1}{L^4} \right) \frac{1}{\delta^2}. \quad (5.82)$$

We have

$$u_{,xy}^L(x, y) = \psi_\delta(x) f^L(x) \hat{u}_{,xy}^L(x, y) + \dot{\psi}_\delta(x) f^L(x) \hat{u}_{,y}^L(x, y) + \psi_\delta(x) \dot{f}^L(x) \hat{u}_{,y}^L(x, y).$$

Thus, by (5.18), (5.19), (5.27), (5.26) and (5.21)

$$\begin{aligned} \int_{-L}^L \int_{-1}^1 (u_{,xy}^L)^2 dx dy &\lesssim \int_{-L}^L \int_{\delta}^1 (f^L(x))^2 (\hat{u}_{,xy}^L)^2 dx dy \\ &\quad + \frac{1}{\delta^2} \int_{-L}^L \int_{\delta}^{2\delta} (f^L(x))^2 (\hat{u}_{,y}^L)^2 dx dy \\ &\quad + \int_{-L}^L \int_{\delta}^1 (\dot{f}^L(x))^2 (\hat{u}_{,y}^L)^2 dx dy \lesssim \frac{1}{\varepsilon^2} + 1 + \frac{1}{\varepsilon} + \frac{\varepsilon}{\delta} \lesssim \frac{1}{\delta\varepsilon}. \end{aligned}$$

This together with (5.75) implies (5.82).

Conclusions. By Step 1 we have that (ω^L, u^L) converges to μ in the sense of Definition 2.3. Moreover, by collecting the estimates showed in Steps 2–6, i.e., (5.54), (5.62), (5.80), (5.81) and (5.82) we find

$$\begin{aligned} \limsup_{L \rightarrow +\infty} L^2(\mathcal{E}_L(w^L, u^L) - \mathcal{E}_0) &\leq \mathcal{F}_\infty(\mu) + C \lim_{L \rightarrow +\infty} \omega(2M^2\delta) \log M \\ &\quad + C \lim_{L \rightarrow +\infty} \left(\frac{L_0^4}{L^2} \frac{1}{\delta^3 M} + L^2 \delta^3 + \frac{1}{L^2 \delta^2} \right). \end{aligned} \quad (5.83)$$

We now proceed with the choice of the parameters. We start by noticing that for every $\overline{M} \in \mathbb{N}$ there exists $L_{\overline{M}} \geq \overline{M}^{1/2}$ such that

$$\omega(2\overline{M}^2 L^{-2/3}) \leq \overline{M}^{-1} \quad \forall L \geq L_{\overline{M}}. \quad (5.84)$$

Since $L_{\overline{M}+1} \geq L_{\overline{M}}$ we set

$$M = M(L) := \overline{M} \quad \text{if } L \in [L_{\overline{M}}, L_{\overline{M}+1})$$

Next we define

$$\delta := \frac{\varepsilon}{M} = L^{-2/3} M^{-1/8} \iff \varepsilon = L^{-2/3} M^{7/8},$$

and we choose

$$L_0 := \frac{L}{n} \in [M^{1/8}, 2M^{1/8}) \iff n \in \left[\frac{L}{2M^{1/8}}, \frac{L}{M^{1/8}} \right).$$

These choices ensure the validity of (5.52) and (5.53). Indeed, we have

$$\varepsilon = \frac{1}{L^{2/3} M^{-7/8}} = \frac{1}{L^{2/3} \overline{M}^{-7/8}} \leq \frac{1}{\overline{M}^3 \overline{M}^{-7/8}} \quad \text{if } L \in [L_{\overline{M}}, L_{\overline{M}+1}),$$

where the last inequality follows from the fact that $L \geq L_{\overline{M}} \geq \overline{M}^{1/2}$. Hence, $\varepsilon \rightarrow 0$ and $\delta \rightarrow 0$ as $L \rightarrow +\infty$. In a similar way we have $M^2 \delta \rightarrow 0$ and $L_0, n \rightarrow +\infty$ as $L \rightarrow +\infty$.

Recalling that ω is monotone, we find

$$\begin{aligned} \omega(2M^2 \delta) &= \omega(2M^2 L^{-2/3} M^{-1/8}) \leq \omega(2M^2 L^{-2/3}) \\ &= \omega(2\overline{M}^2 L^{-2/3}) \quad \text{if } L \in [L_{\overline{M}}, L_{\overline{M}+1}), \end{aligned}$$

which together with (5.84) implies

$$\log M \omega(2M^2 \delta) \leq \log(\overline{M}) \overline{M}^{-1} \quad \text{if } L \in [L_{\overline{M}}, L_{\overline{M}+1}). \quad (5.85)$$

Moreover, if $L \in [L_{\overline{M}}, L_{\overline{M}+1})$ it holds

$$L^2 \delta^3 = L^2 L^{-2} M^{-3/8} = \overline{M}^{-3/8}, \quad (5.86)$$

$$\frac{1}{L^2 \delta^2} = \frac{L^{4/3} M^{1/4}}{L^2} = \frac{M^{1/4}}{L^{2/3}} = \frac{\overline{M}^{1/4}}{L^{2/3}} \leq \frac{\overline{M}^{1/4}}{(\overline{M}^{1/2})^{2/3}} = \overline{M}^{-1/12}, \quad (5.87)$$

and

$$\frac{L_0^4}{L^2} \frac{1}{\delta^3 M} \leq \frac{16M^{1/2}}{MM^{-3/8}} = 16M^{-1/8} = 16\overline{M}^{-1/8}, \quad (5.88)$$

Eventually collecting (5.83)–(5.88) we infer

$$\limsup_{L \rightarrow +\infty} \mathcal{F}_L(w^L, u^L) = \limsup_{L \rightarrow +\infty} L^2 (\mathcal{E}_L(w^L, u^L) - \mathcal{E}_0) \leq \mathcal{F}_\infty(\mu).$$

□

6 Existence and Regularity of Minimizers of $\mathcal{F}_\infty$

In this section we address the existence of minimizers of the limiting functional $\mathcal{F}_\infty$ and we discuss some properties such as equipartition of the energy. In order to do that, we need to introduce the definition of disintegration of measures in the k -variable, which is slightly different from the disintegration in the x -variable introduced in Sect. 3.

In the following for a given interval $I \subset \mathbb{R}$, we denote by $L^0(I)$ the space of functions $g: I \rightarrow \mathbb{R}$ that are Lebesgue measurable. Moreover, the map $\pi_2: I \times \mathbb{R} \rightarrow \mathbb{R}$ denotes the canonical projection, and for any $\mu \in \mathcal{M}_b(I \times \mathbb{R})$ we indicate by $(\pi_2)_\# \mu \in \mathcal{M}_b^+(\mathbb{R})$ its push-forward with respect to the map π_2 .

Definition 6.1 (Disintegration of measures in the k -variable) Let $I \subset \mathbb{R}$ be an interval and let $\mu \in \mathcal{M}_b(I \times \mathbb{R})$. We say that the family

$$(\lambda, (g_k)_{k \in \mathbb{R}}) \quad \text{with} \quad \lambda \in \mathcal{M}_b(\mathbb{R}) \quad \text{and} \quad g_k \in L^0(I) \quad \forall k \in \mathbb{R},$$

is a disintegration of μ (in the k -variable) if $k \mapsto g_k$ is λ -measurable, $\int_0^1 g_k \, dx = 1$ for λ -a.e. $k \in \mathbb{R}$ and

$$\int_{I \times \mathbb{R}} f(x, k) \, d\mu = \int_{\mathbb{R}} \int_I f(x, k) g_k(x) \, dx \, d\lambda(k), \quad (6.1)$$

for every $f \in L^1(I \times \mathbb{R}; |\mu|)$.

With this definition at hand, we can state the main result of this section.

Theorem 6.2 (Minimizers of $\mathcal{F}_\infty$) Let $\mathcal{M}_\infty$ and $\mathcal{F}_\infty$ be as in (2.8) and (2.9), respectively. Then there exists $\hat{\mu} \in \mathcal{M}_\infty$ such that

$$\mathcal{F}_\infty(\hat{\mu}) = \inf_{\mu \in \mathcal{M}_\infty} \mathcal{F}_\infty(\mu).$$

Moreover, every minimizer $\hat{\mu}$ satisfies the following properties: there exist a constant $C > 0$ and a $(\pi_2)_\# \hat{\mu}$ -measurable map $k \mapsto g_k$ with $g_k \in BV(0, 1)$ for $(\pi_2)_\# \hat{\mu}$ a.e. $k \in \mathbb{R}$, such that

$$((\pi_2)_\# \hat{\mu}, (g_k)_{k \in \mathbb{R}}) \quad \text{is a disintegration of } \hat{\mu},$$

$$\int_0^1 k^2 g_k(x) \, dx = \int_0^1 \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{\cdot, x}}{d\hat{\mu}} \right)^2 g_k(x) \, dx \quad \text{for } (\pi_2)_\# \hat{\mu} \text{ a.e. } k \in \mathbb{R}, \quad (6.2)$$

and

$$(\pi_2)_\# \hat{\mu} \left(\{|k| < C\} \right) = 0. \quad (6.3)$$

As a direct consequence, minimizers of $\mathcal{F}_\infty$ satisfy equipartition of the energy.

Corollary 6.3 (*Equipartition of the energy*) *Let $\hat{\mu} \in \mathcal{M}_\infty$ be a minimizer of $\mathcal{F}_\infty$. Then it holds*

$$\int_{(0,1) \times \mathbb{R}} k^2 d\hat{\mu} = \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{,x}}{d\hat{\mu}} \right)^2 d\hat{\mu}.$$

Proof By Theorem 6.2 it holds

$$\begin{aligned} \int_{(0,1) \times \mathbb{R}} k^2 d\hat{\mu} &= \int_{\mathbb{R}} \int_0^1 k^2 g_k(x) dx d(\pi_2)_\# \hat{\mu} \\ &= \int_{\mathbb{R}} \int_0^1 \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{,x}}{d\hat{\mu}} \right)^2 g_k(x) dx d(\pi_2)_\# \hat{\mu} \\ &= \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{,x}}{d\hat{\mu}} \right)^2 d\hat{\mu}. \end{aligned}$$

□

We divide the proof of Theorem 6.2 into several steps. Precisely, we need to show that the functional $\mathcal{F}_\infty$ is convex and lower semi-continuous and that the class of measures $\mathcal{M}_\infty$ admits a disintegration in the k -variable of the form $((\pi_2)_\# \hat{\mu}, (g_k)_{k \in \mathbb{R}})$. First of all we recall that by Remark 2.4 (ii) we have

$$\mathcal{F}_\infty(\mu) = \int_{(0,1) \times \mathbb{R}} k^2 d\mu + \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu}{d|\tilde{\mu}|} \right)^{-1} \left(\frac{d\mu_{,x}}{d|\tilde{\mu}|} \right)^2 d|\tilde{\mu}|,$$

with $\tilde{\mu} = (\mu, \mu_{,x})$ and $|\tilde{\mu}|$ its total variation. This alternative formulation turns out to be more convenient, in particular the term

$$\int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu}{d|\tilde{\mu}|} \right)^{-1} \left(\frac{d\mu_{,x}}{d|\tilde{\mu}|} \right)^2 d|\tilde{\mu}|$$

is reminiscent of the Benamou–Brenier functional used in optimal transport which enjoys nice properties such as lower semi-continuity and convexity. Here we consider a specific case and we refer to Santambrogio (2015, Section 5.3.1) for a general treatment of this topic.

For any $\rho, E \in \mathcal{M}_b((0,1) \times \mathbb{R})$ the Benamou–Brenier functional is defined as

$$\begin{aligned} \mathcal{B}_2(\rho, E) &:= \sup \left\{ \int_{(0,1) \times \mathbb{R}} a(x, k) d\rho \right. \\ &\quad \left. + \int_{(0,1) \times \mathbb{R}} b(x, k) dE : (a, b) \in C_b((0,1) \times \mathbb{R}; K_2) \right\}, \quad (6.4) \end{aligned}$$

where

$$K_2 := \left\{ (z_1, z_2) \in \mathbb{R}^2 : z_1 + \frac{1}{2}z_2^2 \leq 0 \right\}.$$

We next recall some properties of $\mathcal{B}_2$ which follow from Santambrogio (2015, Proposition 5.18). Then we state and prove two intermediate Lemmas (cf. Lemma 6.5 and Lemma 6.6) which will be used to show the validity of Theorem 6.2.

Proposition 6.4 [Properties of $\mathcal{B}_2$] *The functional $\mathcal{B}_2$ is convex and lower semi-continuous on the space $(\mathcal{M}_b((0, 1) \times \mathbb{R}))^2$. Moreover, the following property hold: if both ρ and E are absolutely continuous w.r.t. a same positive measure λ on $(0, 1) \times \mathbb{R}$, then*

$$\mathcal{B}_2(\rho, E) = \int_{(0,1) \times \mathbb{R}} \frac{1}{2} \left(\frac{d\rho}{d\lambda} \right)^{-1} \left(\frac{dE}{d\lambda} \right)^2 d\lambda.$$

Lemma 6.5 (Properties of $\mathcal{F}_\infty$) *The functional $\mathcal{F}_\infty$ is 1-homogeneous, convex and lower semi-continuous on the space $\mathcal{M}_b((0, 1) \times \mathbb{R})$. Moreover, let $(\mu_j) \subset \mathcal{M}_\infty$ be a minimizing sequence, i.e.,*

$$\mathcal{F}_\infty(\mu_j) \rightarrow \inf_{\mu \in \mathcal{M}_\infty} \mathcal{F}_\infty(\mu).$$

Then (μ_j) is pre-compact in $\mathcal{M}_\infty$, i.e., there exists $\hat{\mu} \in \mathcal{M}_\infty$ such that, up to subsequence, $\mu_j \xrightarrow{} \hat{\mu}$. Thus, in particular,*

$$\mathcal{F}_\infty(\hat{\mu}) = \inf_{\mu \in \mathcal{M}_\infty} \mathcal{F}_\infty(\mu).$$

Proof *1-homogeneity.* Let $\alpha > 0$ and let $\mu \in \mathcal{M}_\infty$. Then a direct computation shows that

$$\frac{d(\alpha\mu, x)}{d(\alpha\mu)} = \frac{d\mu, x}{d\mu},$$

from which we readily deduce $\mathcal{F}_\infty(\alpha\mu) = \alpha\mathcal{F}_\infty(\mu)$.

Convexity. Let $\mu_1, \mu_2 \in \mathcal{M}_\infty$, $t \in (0, 1)$. Assume that $\mathcal{F}_\infty(\mu_1), \mathcal{F}_\infty(\mu_2) < +\infty$, otherwise there is nothing to prove. Clearly $\mu_3 = t\mu_1 + (1-t)\mu_2 \in \mathcal{M}_\infty$ and

$$\mathcal{F}_\infty(\mu_3) = \int_{(0,1) \times \mathbb{R}} k^2 d\mu_3 + \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu_3}{d|\tilde{\mu}_3|} \right)^{-1} \left(\frac{d\mu_{3,x}}{d|\tilde{\mu}_3|} \right)^2 d|\tilde{\mu}_3|.$$

We have

$$\int_{(0,1) \times \mathbb{R}} k^2 d\mu_3 = t \int_{(0,1) \times \mathbb{R}} k^2 d\mu_1 + (1-t) \int_{(0,1) \times \mathbb{R}} k^2 d\mu_2. \quad (6.5)$$

Next for $i = 1, 2$ set $\rho_i := \mu_i$, $E_i := \frac{1}{\sqrt{2k}}\mu_{1,x}$, $\lambda_i = |\tilde{\mu}_i|$, and note that they belong to $\mathcal{M}_b((0, 1) \times \mathbb{R})$. Indeed ρ_i, λ_i are bounded by definition, whereas by Young inequality, it holds

$$\begin{aligned} E_i((0, 1) \times \mathbb{R}) &= \int_{(0,1) \times \mathbb{R}} \frac{1}{\sqrt{2k}} d\mu_{i,x} = \int_{(0,1) \times \mathbb{R}} \frac{1}{\sqrt{2k}} \left(\frac{d\mu_{i,x}}{d\mu_i} \right) d\mu_i \\ &\leq \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu_{i,x}}{d\mu_i} \right)^2 d\mu_i + \frac{1}{2} \int_{(0,1) \times \mathbb{R}} d\mu_i \\ &\leq \mathcal{F}_\infty(\mu_i) + \mu_i((0, 1) \times \mathbb{R}) < +\infty. \end{aligned}$$

Thus, we can invoke Proposition 6.4 and get

$$\begin{aligned} \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu_3}{d|\tilde{\mu}_3|} \right)^{-1} \left(\frac{d\mu_{3,x}}{d|\tilde{\mu}_3|} \right)^2 d|\tilde{\mu}_3| &= \mathcal{B}_2 \left(\mu_3, \frac{1}{\sqrt{2k}}\mu_{3,x} \right) \\ &\leq t \mathcal{B}_2 \left(\mu_1, \frac{1}{\sqrt{2k}}\mu_{1,x} \right) + (1-t) \mathcal{B}_2 \left(\mu_2, \frac{1}{\sqrt{2k}}\mu_{2,x} \right) \\ &\leq t \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu_1}{d|\tilde{\mu}_1|} \right)^{-1} \left(\frac{d\mu_{1,x}}{d|\tilde{\mu}_1|} \right)^2 d|\tilde{\mu}_1| \\ &\quad + (1-t) \int_{(0,1) \times \mathbb{R}} \frac{1}{4k^2} \left(\frac{d\mu_2}{d|\tilde{\mu}_2|} \right)^{-1} \left(\frac{d\mu_{2,x}}{d|\tilde{\mu}_2|} \right)^2 d|\tilde{\mu}_2|. \end{aligned} \quad (6.6)$$

Finally combining (6.5) with (6.6) we get

$$\mathcal{F}_\infty(\mu_3) \leq t \mathcal{F}_\infty(\mu_1) + (1-t) \mathcal{F}_\infty(\mu_2).$$

Lower semi-continuity. Let $(\mu_j) \subset \mathcal{M}_\infty$ be such that $\mu_j \xrightarrow{*} \mu$ for some $\mu \in \mathcal{M}_\infty$. Let $\varphi \in C_c^\infty((0, 1) \times \mathbb{R})$, then by duality we have

$$\begin{aligned} \lim_{j \rightarrow +\infty} \int_{(0,1) \times \mathbb{R}} \varphi d\mu_{j,x} &= - \lim_{j \rightarrow +\infty} \int_{(0,1) \times \mathbb{R}} \varphi_{,x} d\mu_j = - \int_{(0,1) \times \mathbb{R}} \varphi_{,x} d\mu \\ &= - \int_{(0,1) \times \mathbb{R}} \varphi d\mu_{,x}, \end{aligned}$$

so that $\mu_{j,x} \xrightarrow{*} \mu_{,x}$. Then by applying Reshetnyak Theorem (Ambrosio et al. 2000, Theorem 2.38) exactly as in (4.14) and (4.15) we deduce

$$\liminf_{j \rightarrow +\infty} \mathcal{F}_\infty(\mu_j) \geq \mathcal{F}_\infty(\mu).$$

Compactness. Let $(\mu_j) \subset \mathcal{M}_\infty$ be a minimizing sequence for $\mathcal{F}_\infty$. Then by Corollary 3.3 for every j there exists $x \mapsto \nu_{j,x}$ with $\nu_{j,x}$ probability measure on $\mathbb{R}$ such that

$$|\mu_j|((0, 1) \times \mathbb{R}) = \int_{(0,1) \times \mathbb{R}} d\mu_j = \int_0^1 \int_{\mathbb{R}} d\nu_{j,x} 2x dx = \int_0^1 2x dx = 1.$$

Thus, up to subsequence, $\mu_j \xrightarrow{*} \hat{\mu}$. Moreover, by Young's inequality and (Ambrosio et al. 2000, Proposition 1.23) we have

$$C \geq \mathcal{F}_\infty(\mu_j) \geq \int_{(0,1) \times \mathbb{R}} \left| \frac{d\mu_{j,x}}{d\mu_j} \right| d\mu_j = |\mu_{j,x}|((0,1) \times \mathbb{R}).$$

Thus, up to subsequence and arguing as in the proof of Proposition 4.1, we can deduce that $\mu_{j,x} \xrightarrow{*} \hat{\mu}_{,x}$ and $\hat{\mu} \in \mathcal{M}_\infty$.

Minimality of $\hat{\mu}$. By lower semi-continuity and compactness, we have

$$\inf_{\mu \in \mathcal{M}_\infty} \mathcal{F}_\infty(\mu) = \lim_{j \rightarrow +\infty} \mathcal{F}_\infty(\mu_j) \geq \mathcal{F}_\infty(\hat{\mu}) \geq \inf_{\mu \in \mathcal{M}_\infty} \mathcal{F}_\infty(\mu),$$

so that

$$\mathcal{F}_\infty(\hat{\mu}) = \inf_{\mu \in \mathcal{M}_\infty} \mathcal{F}_\infty(\mu).$$

□

Lemma 6.6 (Disintegration of $\mu \in \mathcal{M}_\infty$ in the k -variable) *Let $\mu \in \mathcal{M}_\infty$ with $\mathcal{F}_\infty(\mu) < +\infty$. Then there exists $k \mapsto g_k$ $(\pi_2)_\# \mu$ -measurable such that $((\pi_2)_\# \hat{\mu}, (g_k)_{k \in \mathbb{R}})$ is a disintegration of μ (in the k -variable). Moreover, for $(\pi_2)_\# \mu$ a.e. $k \in \mathbb{R}$*

$$g_k \in W^{1,1}(0,1) \quad \text{with} \quad \dot{g}_k = \frac{d\mu_{,x}}{d\mu}(\cdot, k) g_k \mathcal{L}^1.$$

Proof By the Disintegration Theorem (cf. Ambrosio et al. 2000, Theorem 2.28)) there exists $k \mapsto v_k \in \mathcal{M}_b^+(0,1)$ $(\pi_2)_\# \mu$ -measurable with $v_k(0,1) = 1$ such that

$$\int_{(0,1) \times \mathbb{R}} f(x,k) d\mu(x,k) = \int_{\mathbb{R}} \int_0^1 f(x,k) dv_k(x) d(\pi_2)_\# \mu(k), \quad (6.7)$$

for all $f \in L^1((0,1) \times \mathbb{R}; \mu)$. Let $\varphi(x,k) = \phi(x)\chi_A(k)$ with $\phi \in C_c^\infty(0,1)$ and $A \subset \mathbb{R}$ bounded and measurable. Then, being $\varphi_{,x}(x,k) = \dot{\phi}(x)\chi_A(k)$, from Remark 2.2 (i) we have

$$\int_{(0,1) \times \mathbb{R}} \varphi d\mu_{,x} = - \int_{(0,1) \times \mathbb{R}} \varphi_{,x} d\mu = - \int_{\mathbb{R}} \int_0^1 \varphi_{,x} dv_k(x) d(\pi_2)_\# \mu(k).$$

Moreover, since $\mu_{,x} \ll \mu$,

$$\int_{(0,1) \times \mathbb{R}} \varphi d\mu_{,x} = \int_{(0,1) \times \mathbb{R}} \varphi \frac{d\mu_{,x}}{d\mu} d\mu = \int_{\mathbb{R}} \int_0^1 \varphi \frac{d\mu_{,x}}{d\mu} dv_k(x) d(\pi_2)_\# \mu(k).$$

Therefore, we deduce

$$-\int_A \int_0^1 \dot{\phi}(x) \, dv_k(x) \, d(\pi_2)_\# \mu(k) = \int_A \int_0^1 \phi(x) \frac{d\mu_{\cdot,x}}{d\mu} \, dv_k(x) \, d(\pi_2)_\# \mu(k).$$

By the arbitrariness of A this implies

$$-\int_0^1 \dot{\phi}(x) \, dv_k(x) = \int_0^1 \phi(x) \frac{d\mu_{\cdot,x}}{d\mu}(\cdot, k) \, dv_k(x) \quad \text{for } (\pi_2)_\# \mu \text{ a.e. } k \in \mathbb{R},$$

from which, in turn, it follows that $(v_k)_{\cdot,x} \ll v_k$ and

$$(v_k)_{\cdot,x} = \frac{d\mu_{\cdot,x}}{d\mu}(\cdot, k) v_k \quad \text{for } (\pi_2)_\# \mu \text{ a.e. } k \in \mathbb{R}. \quad (6.8)$$

We next claim that (6.8) implies that $v_k \ll \mathcal{L}^1 \llcorner (0, 1)$ and there exists $g_k \in W^{1,1}(0, 1)$ such that $v_k = g_k \mathcal{L}^1 \llcorner (0, 1)$ for $(\pi_2)_\# \mu$ a.e. $k \in \mathbb{R}$. This is enough to conclude the proof since (6.7) becomes

$$\int_{(0,1) \times \mathbb{R}} f(x, k) \, d\mu(x, k) = \int_{\mathbb{R}} \int_0^1 f(x, k) g_k(x) \, dx \, d(\pi_2)_\# \mu(k).$$

It remains to show the claim. Let $\delta > 0$ and let $\rho_\delta(x) = \frac{1}{\delta} \rho(\frac{x}{\delta})$ be a smooth mollifier at scale δ . From (6.8) we have

$$(v_k * \rho_\delta)_{\cdot,x} = \left(\frac{d\mu_{\cdot,x}}{d\mu}(\cdot, k) v_k \right) * \rho_\delta \quad \text{for } (\pi_2)_\# \mu \text{ a.e. } k \in \mathbb{R},$$

where here is implicitly assumed v_k to be extended to 0 in $(0, 1)^c$. Thus

$$|v_k * \rho_\delta(x)| = \left| \int_{-\infty}^x \left(\frac{d\mu_{\cdot,x}}{d\mu}(\cdot, k) v_k \right) * \rho_\delta \, dt \right| \leq \int_0^1 \left| \frac{d\mu_{\cdot,x}}{d\mu}(\cdot, k) \right| \, dv_k < +\infty,$$

so that, in particular $v_k * \rho_\delta \in L^\infty(0, 1)$. This together with $v_k * \rho_\delta \xrightarrow{*} v_k$ implies $v_k = g_k \mathcal{L}^1 \llcorner (0, 1)$ for some $g_k \in L^\infty(0, 1)$. In addition, given $\psi \in C_c^\infty(0, 1)$,

$$\int_0^1 \dot{\psi} g_k \, dx = \int_0^1 \dot{\psi} \, dv_k = - \int_0^1 \psi \frac{d\mu_{\cdot,x}}{d\mu}(\cdot, k) \, dv_k,$$

from which we infer $\dot{g}_k = \frac{d\mu_{\cdot,x}}{d\mu}(\cdot, k)v_k = \frac{d\mu_{\cdot,x}}{d\mu}(\cdot, k)g_k\mathcal{L}^1$. Eventually by Young's inequality

$$\begin{aligned}\int_0^1 |\dot{g}_k(x)| \, dx &= \int_0^1 \frac{|\dot{g}_k(x)|}{\sqrt{2k}\sqrt{g_k(x)}} \sqrt{2k}\sqrt{g_k(x)} \, dx \\ &\leq \int_0^1 \left[k^2 g_k(x) + \frac{1}{4k^2} \frac{(\dot{g}_k(x))^2}{g_k(x)} \right] \, dx \\ &\leq \int_0^1 \left[k^2 + \frac{1}{4k^2} \left(\frac{d\mu_{\cdot,x}}{d\mu}(\cdot, k) \right)^2 \right] g_k(x) \, dx < +\infty\end{aligned}$$

for $(\pi_2)_\# \mu$ a.e. $k \in \mathbb{R}$, and thus in particular $g_k \in W^{1,1}(0, 1)$.

□

We are now ready to prove the main result of this section.

Proof of Theorem 6.2 By Lemma 6.5 we know there exists $\hat{\mu} \in \mathcal{M}_\infty$ minimizer of $\mathcal{F}_\infty$. Moreover, by Lemma 6.6 there exists $k \mapsto g_k(\pi_2)_\# \hat{\mu}$ measurable with $g_k \in W^{1,1}(0, 1)$ for $(\pi_2)_\# \hat{\mu}$ a.e. $k \in \mathbb{R}$, such that $((\pi_2)_\# \hat{\mu}, (g_k)_{k \in \mathbb{R}})$ is a disintegration of $\hat{\mu}$. Therefore, in particular we can rewrite

$$\mathcal{F}_\infty(\hat{\mu}) = \int_{\mathbb{R}} \int_0^1 \left[k^2 + \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{\cdot,x}}{d\hat{\mu}} \right)^2 \right] g_k(x) \, dx \, d(\pi_2)_\# \hat{\mu}.$$

Step 1: we show (6.2). Assume by contradiction that (6.2) does not hold true. Then the set

$$E := \left\{ k \in \mathbb{R} : \int_0^1 k^2 g_k(x) \, dx \neq \int_0^1 \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{\cdot,x}}{d\hat{\mu}} \right)^2 g_k(x) \, dx \right\}$$

is such that $(\pi_2)_\# \hat{\mu}(E) > 0$. Assume, without loss of generality, that the subset

$$E_1^+ := \left\{ k \in \mathbb{R}^+ : \int_0^1 k^2 g_k(x) \, dx < \int_0^1 \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{\cdot,x}}{d\hat{\mu}} \right)^2 g_k(x) \, dx \right\} \subset E$$

satisfies $(\pi_2)_\# \hat{\mu}(E_1^+) > 0$ (the other cases can be treated in a similar way). Since $\int_0^1 g_k(x) \, dx = 1$, we can rewrite

$$E_1^+ = \left\{ k \in \mathbb{R}^+ : 1 < \frac{1}{4k^4} \int_0^1 \left(\frac{d\hat{\mu}_{\cdot,x}}{d\hat{\mu}} \right)^2 g_k(x) \, dx \right\}.$$

Then there exists $\sigma > 0$ such that

$$E_\sigma := \left\{ k \in \mathbb{R}^+ : 1 + \sigma \leq \frac{1}{4k^2} \int_0^1 \left(\frac{d\hat{\mu}_{\cdot,x}}{d\hat{\mu}} \right)^2 g_k(x) \, dx \right\} \subset E_1^+$$

with $(\pi_2)_\# \hat{\mu}(E_\sigma) > 0$. Next fix $\delta > 0$ such that $(1 + \delta)^2 < 1 + \sigma$ and let $\tilde{\mu} \in \mathcal{M}_b((0, 1) \times \mathbb{R})$ be defined as follows

$$\tilde{\mu} := \hat{\mu} \llcorner (0, 1) \times (\mathbb{R} \setminus E_\sigma) + \mu_\delta, \quad (6.9)$$

where $\mu_\delta := (\tau_\delta)_\# \mu \llcorner ((0, 1) \times E_\sigma)$ is the push-forward of $\mu \llcorner ((0, 1) \times E_\sigma)$ with respect to the map $\tau_\delta: (0, 1) \times \mathbb{R} \rightarrow (0, 1) \times \mathbb{R}$, $\tau_\delta(x, k) := (x, k(1 + \delta))$. Note that $\tilde{\mu}$ is a positive measure. Setting $E_\sigma^\delta := (1 + \delta)E_\sigma$, then the support of μ_δ is contained in $(0, 1) \times E_\sigma^\delta$, and

$$\int_{(0,1) \times E_\sigma^\delta} f(x, k) \, d\mu_\delta = \int_{(0,1) \times E_\sigma} f(x, k(1 + \delta)) \, d\hat{\mu}$$

for every f summable with respect to μ_δ . Hence, by duality and using that $\hat{\mu}_{,x} \ll \hat{\mu}$ we have

$$\begin{aligned} \int_{(0,1) \times E_\sigma^\delta} \varphi(x, k) \, d(\mu_\delta)_{,x} &= - \int_{(0,1) \times E_\sigma^\delta} \varphi_{,x}(x, k) \, d\mu_\delta \\ &= - \int_{(0,1) \times E_\sigma} \varphi_{,x}(x, k(1 + \delta)) \, d\hat{\mu} \\ &= \int_{(0,1) \times E_\sigma} \varphi(x, k(1 + \delta)) \, d\hat{\mu}_{,x} \\ &= \int_{(0,1) \times E_\sigma} \varphi(x, k(1 + \delta)) \frac{d\hat{\mu}_{,x}}{d\hat{\mu}}(x, k) \, d\hat{\mu} \\ &= \int_{(0,1) \times E_\sigma^\delta} \varphi(x, k) \frac{d\hat{\mu}_{,x}}{d\hat{\mu}}\left(x, \frac{k}{1 + \delta}\right) \, d\mu_\delta, \end{aligned}$$

for all $\varphi \in C_c^\infty((0, 1) \times E_\sigma^\delta)$. As a consequence we readily deduce that $(\mu_\delta)_{,x} \ll \mu_\delta$ with

$$\frac{d(\mu_\delta)_{,x}}{d\mu_\delta}(x, k) = \frac{d\hat{\mu}_{,x}}{d\hat{\mu}}\left(x, \frac{k}{1 + \delta}\right), \quad (6.10)$$

so that, in particular, $\tilde{\mu}_{,x} \ll \tilde{\mu}$. Moreover, for every $\phi \in C_c^\infty(0, 1)$ we have

$$\begin{aligned} \int_0^1 2x\phi(x) \, dx &= \int_{(0,1) \times \mathbb{R}} \phi(x) \, d\hat{\mu} = \int_{(0,1) \times E_\sigma^c} \phi(x) \, d\hat{\mu} + \int_{(0,1) \times E_\sigma} \phi(x) \, d\hat{\mu} \\ &= \int_{(0,1) \times E_\sigma^c} \phi(x) \, d\hat{\mu} \\ &\quad + \int_{(0,1) \times E_\sigma^\delta} \phi(x) \, d\mu_\delta = \int_{(0,1) \times \mathbb{R}} \phi(x) \, d\tilde{\mu}, \end{aligned}$$

and thus $\tilde{\mu} \in \mathcal{M}_\infty$. We next show that

$$\mathcal{F}_\infty(\tilde{\mu}) < \mathcal{F}_\infty(\hat{\mu}),$$

which contradicts the fact that $\hat{\mu}$ is a minimizer. To this purpose, it is convenient to define the localized functional

$$\mathcal{F}_\infty(\mu, A) := \int_{(0,1) \times A} \left[k^2 + \frac{1}{4k^2} \left(\frac{d\mu_{,x}}{d\mu} \right)^2 \right] d\mu,$$

for any bounded measure μ with $\mu_{,x} \ll \mu$ and any $A \subset \mathbb{R}$ measurable. Observing that $\tilde{\mu} = \hat{\mu}$ on $(0, 1) \times (\mathbb{R} \setminus (E_\sigma \cup E_\sigma^\delta))$ and $\tilde{\mu} = \mu_\delta$ on $(0, 1) \times (E_\sigma^\delta \cap E_\sigma)$ we have

$$\mathcal{F}_\infty(\tilde{\mu}) = \mathcal{F}_\infty(\hat{\mu}, \mathbb{R} \setminus (E_\sigma \cup E_\sigma^\delta)) + \mathcal{F}_\infty(\tilde{\mu}, E_\sigma^\delta \setminus E_\sigma) + \mathcal{F}_\infty(\mu_\delta, E_\sigma^\delta \cap E_\sigma). \quad (6.11)$$

By Lemma 6.5 we know that $\mathcal{F}_\infty$ is convex and 1-homogeneous, which together with

$$\tilde{\mu} = \frac{2\hat{\mu} + 2\mu_\delta}{2} \quad \text{on } (0, 1) \times (E_\sigma^\delta \setminus E_\sigma),$$

yield

$$\begin{aligned} \mathcal{F}_\infty(\tilde{\mu}, E_\sigma^\delta \setminus E_\sigma) &= \mathcal{F}_\infty(\tilde{\mu} \llcorner ((0, 1) \times E_\sigma^\delta \setminus E_\sigma)) \\ &\leq \mathcal{F}_\infty(\hat{\mu} \llcorner ((0, 1) \times E_\sigma^\delta \setminus E_\sigma)) + \mathcal{F}_\infty(\mu_\delta \llcorner ((0, 1) \times E_\sigma^\delta \setminus E_\sigma)) \\ &= \mathcal{F}_\infty(\hat{\mu}, (E_\sigma^\delta \setminus E_\sigma)) + \mathcal{F}_\infty(\mu_\delta, (E_\sigma^\delta \setminus E_\sigma)). \end{aligned}$$

Combining this together with (6.11), we get

$$\begin{aligned} \mathcal{F}_\infty(\tilde{\mu}) &\leq \mathcal{F}_\infty(\hat{\mu}, \mathbb{R} \setminus E_\sigma) + \mathcal{F}_\infty(\mu_\delta, E_\sigma^\delta) \\ &= \mathcal{F}_\infty(\hat{\mu}) + \mathcal{F}_\infty(\mu_\delta, E_\sigma^\delta) - \mathcal{F}_\infty(\hat{\mu}, E_\sigma). \end{aligned}$$

Therefore, we would conclude the proof if we show that

$$\mathcal{F}_\infty(\mu_\delta, E_\sigma^\delta) - \mathcal{F}_\infty(\hat{\mu}, E_\sigma) < 0. \quad (6.12)$$

By the change of variable $k = \hat{k}(1 + \delta)$ and recalling (6.10), it holds

$$\begin{aligned} \mathcal{F}_\infty(\mu_\delta, E_\sigma^\delta) &= \int_{(0,1) \times E_\sigma^\delta} \left[k^2 + \frac{1}{4k^2} \left(\frac{d(\mu_\delta)_{,x}}{d\mu_\delta} \right)^2 \right] d\mu_\delta \\ &= \int_{(0,1) \times E_\sigma} \left[k^2(1 + \delta)^2 + \frac{1}{4k^2(1 + \delta)^2} \left(\frac{d\hat{\mu}_{,x}}{d\hat{\mu}} \right)^2 \right] d\hat{\mu}, \end{aligned}$$

from which it follows

$$\begin{aligned} & \mathcal{F}_\infty(\mu_\delta, E_\sigma^\delta) - \mathcal{F}_\infty(\hat{\mu}, E_\sigma) \\ &= \int_{(0,1) \times E_\sigma} \left[k^2((1+\delta)^2 - 1) + \frac{1 - (1+\delta)^2}{4k^2(1+\delta)^2} \left(\frac{d\hat{\mu}_{,x}}{d\hat{\mu}} \right)^2 \right] d\hat{\mu} \\ &= \frac{(1+\delta)^2 - 1}{(1+\delta)^2} \int_{(0,1) \times E_\sigma} \left[k^2(1+\delta)^2 - \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{,x}}{d\hat{\mu}} \right)^2 \right] d\hat{\mu}. \end{aligned}$$

Being $\delta > 0$ we have that $((1+\delta)^2 - 1)/(1+\delta)^2 > 0$. Moreover, by disintegration we can rewrite the integral as

$$\begin{aligned} & \int_{(0,1) \times E_\sigma} \left[k^2(1+\delta)^2 - \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{,x}}{d\hat{\mu}} \right)^2 \right] d\hat{\mu} \\ &= \int_{E_\sigma} \int_0^1 \left[k^2(1+\delta)^2 - \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{,x}}{d\hat{\mu}} \right)^2 \right] g_k(x) dx d(\pi_2)_\# \hat{\mu}. \end{aligned}$$

The above quantity is strictly negative if

$$\int_0^1 \left[k^2(1+\delta)^2 - \frac{1}{4k^2} \left(\frac{d\hat{\mu}_{,x}}{d\hat{\mu}} \right)^2 \right] g_k(x) dx < 0 \quad \text{for } (\pi_2)_\# \hat{\mu} \text{ a.e. } k \in E_\sigma.$$

From $\int_0^1 g_k(x) dx = 1$, this is equivalent to

$$(1+\delta)^2 < \frac{1}{4k^4} \int_0^1 \left(\frac{d\hat{\mu}_{,x}}{d\hat{\mu}} \right)^2 g_k(x) dx \quad \text{for } (\pi_2)_\# \hat{\mu} \text{ a.e. } k \in E_\sigma,$$

which holds thanks to the choice of δ and the definition of E_σ , and thus, we infer (6.12).

Step 2: we show (6.3). By Lemma 6.6 we have $\dot{g}_k = \frac{d\hat{\mu}_{,x}}{d\hat{\mu}}(\cdot, k) g_k \mathcal{L}^1$, so that from (6.2) we have

$$\begin{aligned} k^2 \int_0^1 (\sqrt{g_k(x)})^2 dx &= k^2 \int_0^1 g_k(x) dx = \frac{1}{4k^2} \int_0^1 \frac{(\dot{g}_k(x))^2}{g_k(x)} dx \\ &= \frac{1}{k^2} \int_0^1 \left(\frac{d}{dx} \sqrt{g_k(x)} \right)^2 dx. \end{aligned} \quad (6.13)$$

Since $\hat{\mu} \in \mathcal{M}_\infty$, then in particular $\hat{\mu}(\{0\} \times \mathbb{R}) = 0$ from which it follows $g_k(0) = 0$ for $(\pi_2)_\# \hat{\mu}$ a.e. $k \in \mathbb{R}$. Thus, we apply Poincaré's inequality to get

$$\int_0^1 (\sqrt{g_k(x)})^2 dx \leq C \int_0^1 \left(\frac{d}{dx} \sqrt{g_k(x)} \right)^2 dx,$$

for some constant $C > 0$. Now combining the above inequality with (6.13) we find that

$$\int_0^1 (\sqrt{g_k(x)})^2 dx \leq Ck^4 \int_0^1 (\sqrt{g_k(x)})^2 dx \iff k^4 \geq C^{-1}.$$

Hence, (6.2) holds true if $k^4 \geq C^{-1}$ which in turn implies $(\pi_2)_{\sharp} \hat{\mu}(|k| \leq C^{-1/4}) = 0$. $\square$

Acknowledgements This work was done while the second author was postdoc at TU Dortmund. Both authors were partially supported by the German Science Foundation DFG in context of the Emmy Noether Junior Research Group BE 5922/1-1. RM has been partially funded by the European Union - NextGenerationEU under the Italian Ministry of University and Research (MUR) National Centre for HPC, Big Data and Quantum Computing CN_00000013 - CUP: E13C22001000006. PB also thanks Carlos Román and Alaa Elshorbagy for discussions at the early stages of the project.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ambrosio, L., Fusco, N., Pallara, D.: Functions of bounded variation and free discontinuity problems. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York (2000)
- Bach, A., Esposito, T., Marziani, R., Zeppieri, C.I.: Interaction between oscillations and singular perturbations in a one-dimensional phase-field model. AWM Volume - Research in the Mathematics of Materials Science (2022)
- Bach, A., Esposito, T., Marziani, R., Zeppieri, C.I.: Gradient damage models for heterogeneous materials. SIAM J. Math. Anal. **55**, 4 (2023)
- Bach, A., Marziani, R., Zeppieri, C.I.: Γ -convergence and stochastic homogenisation of singularly-perturbed elliptic functionals. Calc. Var. **62**, 199 (2023)
- Bedrossian, J., Kohn, R.V.: Blister patterns and energy minimization in compressed thin films on compliant substrates. Commun. Pure Appl. Math. **68**(3), 472–510 (2015)
- Bella, P., Goldman, M.: Nucleation barriers at corners for a cubic-to-tetragonal phase transformation. Proc. R. Soc. Edinb. Sect. A **145**(4), 715–724 (2015)
- Bella, P., Goldman, M., Zwicknagl, B.: Study of island formation in epitaxially strained films on unbounded domains. Arch. Ration. Mech. Anal. **218**(1), 163–217 (2015)
- Bella, P., Kohn, R.V.: Wrinkles as the result of compressive stresses in an annular thin film. Commun. Pure Appl. Math. **67**(5), 693–747 (2014)
- Bella, P., Kohn, R.V.: The annulus problem, using a von Kármán membrane model. Preliminary version <https://math.nyu.edu/~kohn/pcmi/annulus-problem-notes-for-pcmi.pdf>
- Bella, P., Kohn, R.V.: Coarsening of folds in hanging drapes. Commun. Pure Appl. Math. **70**(5), 978–1021 (2017)
- Bella, P., Kohn, R.V.: Wrinkling of a thin circular sheet bonded to a spherical substrate. Philos. Trans. R. Soc. A **375**, 2093 (2017)
- Bella, P.: Wrinkles as a relaxation of compressive stresses. ProQuest LLC, Ann Arbor, MI, 2012, Thesis (Ph.D.)—New York University

- Bella, P.: The transition between planar and wrinkled regions in a uniaxially stretched thin elastic film. *Arch. Ration. Mech. Anal.* **216**(2), 623–672 (2015)
- Bella, P., Kohn, R.V.: Metric-induced wrinkling of a thin elastic sheet. *J. Nonlinear Sci.* **24**(6), 1147–1176 (2014)
- Bella, P., Román, C.: Wrinkling of an elastic sheet floating on a liquid sphere (2023)
- Ben Belgacem, H., Conti, S., DeSimone, A., Müller, S.: Rigorous bounds for the Föppl-von Kármán theory of isotropically compressed plates. *J. Nonlinear Sci.* **10**(6), 661–683 (2000)
- Ben Belgacem, H., Conti, S., DeSimone, A., Müller, S.: Energy scaling of compressed elastic films—three-dimensional elasticity and reduced theories. *Arch. Ration. Mech. Anal.* **164**(1), 1–37 (2002)
- Bourne, D., Conti, S., Müller, S.: Folding patterns in partially delaminated thin films. In: *Innovative Numerical Approaches for Multi-Field and Multi-Scale Problems. Lecture Notes in Applied and Computational Mechanics*, vol 81. Springer, Cham, pp. 25–39 (2016)
- Brandman, J., Kohn, R.V., Nguyen, H.-M.: Energy scaling laws for conically constrained thin elastic sheets. *J. Elast.* **113**(2), 251–264 (2013)
- Brietzke, B., Knüpfer, H.: Onset of pattern formation in thin ferromagnetic films with perpendicular anisotropy. *Calc. Var. Partial Differ. Equ.* **62**(4), 133 (2023)
- Capella, A., Otto, F.: A rigidity result for a perturbation of the geometrically linear three-well problem. *Commun. Pure Appl. Math.* **62**(12), 1632–1669 (2009)
- Capella, A., Otto, F.: A quantitative rigidity result for the cubic-to-tetragonal phase transition in the geometrically linear theory with interfacial energy. *Proc. R. Soc. Edinb. Sect. A* **142**(2), 273–327 (2012)
- Choksi, R.: Scaling laws in microphase separation of diblock copolymers. *J. Nonlinear Sci.* **11**(3), 223–236 (2001)
- Conti, S.: Branched microstructures: scaling and asymptotic self-similarity. *Commun. Pure Appl. Math.* **53**(11), 1448–1474 (2000)
- Conti, S., Garroni, A., Marziani, R.: Line-tension limits for line singularities and application to the mixed-growth case. *Calc. Var. PDE* **62**, 228 (2023)
- Conti, S., Maggi, F.: Confining thin elastic sheets and folding paper. *Arch. Ration. Mech. Anal.* **187**(1), 1–48 (2008)
- Davidovitch, B., Schroll, R.D., Cerda, E.: Nonperturbative model for wrinkling in highly bendable sheets. *Phys. Rev. E* **85**, 066115 (2012)
- Davidovitch, B., Schroll, R.D., Vella, D., Adda-Bedia, M., Cerda, E.: Prototypical model for tensional wrinkling in thin sheets. *Proc. Natl. Acad. Sci.* **108**(45), 18227–18232 (2011)
- Garroni, A., Marziani, R., Scala, R.: Derivation of a line-tension model for dislocations from a nonlinear three-dimensional energy: the case of quadratic growth. *SIAM J. Math. Anal.* **53**, 4252–4302 (2021)
- Géminard, J.-C., Bernal, R., Melo, F.: Wrinkle formations in axi-symmetrically stretched membranes. *Eur. Phys. J. E* **15**(2), 117–126 (2004)
- Huang, J., Davidovitch, B., Santangelo, C.D., Russell, T.P., Menon, N.: Smooth cascade of wrinkles at the edge of a floating elastic film. *Phys. Rev. Lett.* **105**, 038302 (2010)
- Jin, W., Sternberg, P.: Energy estimates for the von Kármán model of thin-film blistering. *J. Math. Phys.* **42**(1), 192–199 (2001)
- Knüpfer, H., Kohn, R.V., Otto, F.: Nucleation barriers for the cubic-to-tetragonal phase transformation. *Commun. Pure Appl. Math.* **66**(6), 867–904 (2013)
- Kohn, R.V., Müller, S.: Surface energy and microstructure in coherent phase transitions. *Commun. Pure Appl. Math.* **47**(4), 405–435 (1994)
- Kohn, R.V.: Energy-driven pattern formation, *International Congress of Mathematicians*, vol. I, pp. 359–383. Eur. Math. Soc., Zürich (2007)
- Kohn, R.V., Nguyen, H.-M.: Analysis of a compressed thin film bonded to a compliant substrate: the energy scaling law. *J. Nonlinear Sci.* **23**(3), 343–362 (2013)
- Kohn, R.V., O’Brien, E.: The wrinkling of a twisted ribbon. *J. Nonlinear Sci.* **28**(4), 1221–1249 (2018)
- Kohn, R.V., Wirth, B.: Optimal fine-scale structures in compliance minimization for a uniaxial load. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **470**(2170), 20140432 (2014)
- Kohn, R.V., Wirth, B.: Optimal fine-scale structures in compliance minimization for a shear load. *Commun. Pure Appl. Math.* **69**(8), 1572–1610 (2016)
- Maor, C., Shachar, A.: On the role of curvature in the elastic energy of non-Euclidean thin bodies. *J. Elast.* **134**(2), 149–173 (2019)
- Marziani, R.: Γ -convergence and stochastic homogenisation of phase-transition functionals. *ESAIM Control Optim. Calc. Var.* **29**, 44 (2023)

- Marziani, R., Solombrino, F.: Non-local approximation of free-discontinuity problems in linear elasticity and application to stochastic homogenization. *Proc. R. Soc. Edinb. Sect. A Math.* **154**(4), 1060–1094 (2024). <https://doi.org/10.1017/prm.2023.51>
- Olbermann, H.: The shape of low energy configurations of a thin elastic sheet with a single disclination. *Anal. PDE* **11**(5), 1285–1302 (2018)
- Olbermann, H.: On a boundary value problem for conically deformed thin elastic sheets. *Anal. PDE* **12**(1), 245–258 (2019)
- Otto, F.: Cross-over in scaling laws: a simple example from micromagnetics. In: *Proceedings of the International Congress of Mathematicians, Vol. III (Beijing, 2002)*, Higher Ed. Press, Beijing, pp. 829–838 (2002)
- Padilla-Garza, D.: An asymptotic variational problem modeling a thin elastic sheet on a liquid, lifted at one end. *Math. Methods Appl. Sci.* **43**(8), 4956–4973 (2020)
- Petrache, M., Serfaty, S.: Next order asymptotics and renormalized energy for Riesz interactions. *J. Inst. Math. Jussieu* **16**(3), 501–569 (2017)
- Pipkin, A.C.: Relaxed energy densities for large deformations of membranes. *IMA J. Appl. Math.* **52**(3), 297–308 (1994)
- Ried, T., Román, C.: Domain branching in micromagnetism: scaling law for the global and local energies. *Rev. Mat. Iberoam.* (2024). <https://doi.org/10.4171/RMI/1508>
- Román, C., Sandier, E., Serfaty, S.: Bounded vorticity for the 3D Ginzburg-Landau model and an isoflux problem. *Proc. Lond. Math. Soc. (3)* **126**(3), 1015–1062 (2023)
- Rüland, A., Tribuzio, A.: On scaling laws for multi-well nucleation problems without gauge invariances. *J. Nonlinear Sci.* **33**(2), 25 (2023a)
- Rüland, A., Tribuzio, A.: On the energy scaling behaviour of singular perturbation models with prescribed Dirichlet data involving higher order laminates. *ESAIM Control Optim. Calc. Var.* **29**, 68 (2023b)
- Santambrogio, F.: *Optimal transport for applied mathematicians. Progress in Nonlinear Differential Equations and their Applications*, vol. 87, Birkhäuser/Springer, Cham (2015), *Calculus of variations, PDEs, and modeling*
- Serfaty, S.: *Coulomb gases and Ginzburg-Landau vortices, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich (2015)*
- Tobasco, I.: Axial compression of a thin elastic cylinder: bounds on the minimum energy scaling law. *Commun. Pure Appl. Math.* **71**(2), 304–355 (2018)
- Tobasco, I.: Curvature-driven wrinkling of thin elastic shells. *Arch. Ration. Mech. Anal.* **239**(3), 1211–1325 (2021)
- Vandeparre, H., Piñeirua, M., Brau, F., Roman, B., Bico, J., Gay, C., Bao, W., Lau, C.N., Reis, P.M., Damman, P.: Wrinkling hierarchy in constrained thin sheets from suspended graphene to curtains. *Phys. Rev. Lett.* **106**, 224301 (2011)
- Venkataramani, S.: Lower bounds for the energy in a crumpled elastic sheet—a minimal ridge. *Nonlinearity* **17**(1), 301–312 (2004)