Measurement of Deeply Virtual Compton Scattering at HERA

Vom Fachbereich Physik
der Universität Dortmund
und der Faculté des Sciences
Université Libre de Bruxelles
zur Erlangung des akademischen Grades
eines Doktors der Naturwissenschaften
genehmigte

Dissertation

von
Diplom–Physiker Rainer Stamen
aus Ense

Dortmund
November 2001

Abstract

In this thesis the results of the first measurement of Deeply Virtual Compton Scattering in electron–proton collisions at HERA are presented. The analysed data were taken with the H1 detector during the 1997 data taking period corresponding to an integrated luminosity of $8\,\mathrm{pb}^{-1}$. The differential cross sections $\frac{d\sigma}{dQ^2}$ and $\frac{d\sigma}{dW}$ of the reaction $ep \rightarrow e\gamma p$ have been measured in the kinematic domain $2 < Q^2 < 20\,\mathrm{GeV}^2$, $30 < W < 120\,\mathrm{GeV}$ and $|t| < 1\,\mathrm{GeV}^2$. After subtracting the contribution of the Bethe–Heitler process the photon proton cross section for the DVCS process $\sigma^{\gamma^* p \longrightarrow \gamma p}$ has been determined. The results are compared to QCD based calculations which are able to describe the measurement.

Kurzfassung

Die Ergebnisse der ersten Messung tiefvirtueller Comptonstreuung in Elektron–Proton Kollisionen bei HERA werden vorgestellt. Die analysierten Daten wurden mit dem H1 Detektor während der Datennahmeperiode 1997 aufgezeichnet und entsprechen einer integrierten Luminosität von $8\,\mathrm{pb}^{-1}$. Die differentiellen Wirkungsquerschnitte $\frac{d\sigma}{dQ^2}$ und $\frac{d\sigma}{dW}$ für die Reaktion $ep \rightarrow e\gamma p$ wurden im kinematischen Bereich $2 < Q^2 < 20\,\mathrm{GeV}^2$, $30 < W < 120\,\mathrm{GeV}$ und $|t| < 1\,\mathrm{GeV}^2$ gemessen. Nach Abzug des Beitrags des Bethe–Heitler Prozesses, wurde der Photon Proton Wirkungsquerschnitt $\sigma^{\gamma^* p \longrightarrow \gamma p}$ bestimmt. Die Ergebnisse werden mit QCD basierten Berechnungen verglichen, die in der Lage sind die Messung zu beschreiben.

Resumé

La présente thèse de doctorat porte sur l'analyse des données accumulées par l'expérience H1 auprès du collisionneur électron-proton HERA. Ce travail présente, pour la première fois, la mesure de la section efficace de la diffusion Compton profondément virtuel. Cette mesure utilise les évènements collectés durant la prise de données de 1997, accumulant une luminosité de 8 pb^{-1}. La section efficace de la réaction $ep \rightarrow e\gamma p$ est présentée differentiellement en $\frac{d\sigma}{dQ^2}$ et $\frac{d\sigma}{dW}$, dans la domaine $2 < Q^2 < 20\,\mathrm{GeV}^2$, $30 < W < 120\,\mathrm{GeV}$ et $|t| < 1\,\mathrm{GeV}^2$. Aprés la soustration du contribution Bethe–Heitler la section efficace photon–proton $\sigma^{\gamma^* p \longrightarrow \gamma p}$ est extrahit. Les résultats obtenus sont en bon accords avec certaines prédictions basées sur les calculs de chromodynamique quantique.

Contents

Chapter 1

Introduction

In high energy lepton–hadron scattering experiments an elementary particle, the lepton, is probing the hadron, a composite particle and thus delivers an excellent tool to investigate the hadrons structure. The results of early experiments led to the partonic picture of hadrons, in which they (e.g. the proton) consist of quarks. This was refined by more precise data and was one of the foundations of Quantum Chromo Dynamics (QCD) the quantum field theory of the strong interaction which describes hadron interactions on the basis of quarks and gluons (partons). Nowadays one of the aims of such lepton–hadron scattering experiments is the precise measurement of the parton density functions and the determination of the strong coupling constant α_s. This provides the possibility for very sensitive tests of the validity of different approximation methods which are applied in perturbative QCD calculations.

At the *ep*–collider HERA[1] a class of reactions has been 'rediscovered' the so called diffractive interactions which in the past have been observed in soft hadron–hadron scattering experiments and which are successfully described in the framework of Regge theory. The aim of current measurements of diffractive interactions is to understand these in the framework of perturbative QCD. Deeply Virtual Compton Scattering (DVCS), the reaction under investigation in this thesis, is a diffractive process for which pQCD calculations are expected to be reliable, and which furthermore opens the possibility to extract information about the structure of the proton which exceeds the meanwhile well known parton density functions.

The measurement of DVCS was proposed to access the generalised parton distributions which are generalisations of the ordinary parton densities and which exhibit also information about elastic form factors and the spin structure of the proton.

1.1 Deep Inelastic Scattering and the Structure of the Proton

In this section a general introduction to Deep Inelastic Scattering (DIS) is presented. The kinematic variables which define the process are introduced and the cross section is defined in terms of the two structure functions F_2 and F_L. These are discussed in the framework of the quark–parton model which is then refined by discussing the basic properties of perturbative QCD. At the end of this section our current knowledge about the partonic structure of the proton is presented.

[1] **H**adron **E**lektron **R**ing **A**nlage

Deep Inelastic Scattering (DIS)

In deep inelastic electron–proton scattering [1, 2, 3] the incoming electron interacts with the incoming proton by the exchange of a single gauge boson as shown in Figure 1.1. In neutral current events (NC) a virtual photon (γ^*) or a Z^0–boson is exchanged and an electron and the hadronic final state X are observed; in charged current interactions a W–boson is exchanged and the final state consists of a neutrino and the hadronic system X.



Figure 1.1: *This diagram for deep inelastic electron–proton scattering shows the exchange of a gauge boson between the incoming lepton and the hadron. In the quark–parton model the electron scatters elastically off a constituent quark of the proton. In neutral current events a neutral gauge boson is exchanged and the final state consists of the electron and the hadronic system X whereas in charged current events a charged W–boson is exchanged and a neutrino and the hadronic system X compose the final state of the reaction. The four vectors of the particles are denoted k and k' for the incoming and outgoing lepton respectively and p for the incoming proton.*

Kinematic Variables

The cross section is defined in terms of Lorentz invariant variables which ensures a convenient comparison of the measurements for different experimental setups. The variable Q^2 is the negative of the squared momentum of the exchanged boson,

$$Q^2 = -q^2 = -(k - k')^2, \tag{1.1}$$

where k is the four momentum of the incoming electron and k' denotes the four momentum of the outgoing lepton. The two Bjorken scaling variables x and y are defined as:

$$x = \frac{-q^2}{2pq}, \qquad y = \frac{pq}{pk}, \tag{1.2}$$

where p denotes the four momentum of the incoming proton. Thus x and y are dimensionless variables defined in the range $0 < x, y \leq 1$. The electron–proton centre of mass energy squared s is given by the equation

$$s = (k + p)^2. \tag{1.3}$$

At fixed centre of mass energy $\sqrt{s}$, only two of these four variables are independent due to energy momentum conservation[2]. When neglecting the electron and proton masses they are connected by the relation

$$Q^2 = xys. \tag{1.4}$$

In addition W is defined as the centre of mass energy of the exchanged boson and the incoming proton

$$W^2 = (q+p)^2 \simeq ys - Q^2. \tag{1.5}$$

Definition of the Cross Section and the Structure Function F_2

The cross section for neutral current *ep* interactions is defined such that it takes into account the different couplings of the gauge bosons to the electron and the propagator terms. In the region of low Q^2 ($Q^2 \ll M_{Z^0}^2$) only photon exchange contributes, resulting in a drastic simplification of the cross section expression. The Z^0 exchange and the interference between Z^0 and photon exchange can be neglected safely since the gauge boson masses enter the cross section by the propagator terms which leads to the following ratios for the contributions of the Z^0 exchange and the interference w.r.t. the contribution from single photon exchange to the cross section:

$$\frac{\sigma\left(Z^0\right)}{\sigma\left(\gamma^*\right)} \sim \left(\frac{Q^2}{Q^2 + M_{Z^0}^2}\right)^2, \tag{1.6}$$

$$\frac{\sigma\left(\gamma^* Z^0\right)}{\sigma\left(\gamma^*\right)} \sim \left(\frac{Q^2}{Q^2 + M_{Z^0}^2}\right). \tag{1.7}$$

At low Q^2 the differential electron–proton cross section can be written as:

$$\frac{d^2\sigma\left(x,Q^2\right)}{dx dQ^2} = \frac{4\pi\alpha_{em}^2}{xQ^4}\left[\left(1 - y + \frac{y^2}{2}\right) F_2\left(x,Q^2\right) - \frac{y^2}{2} F_L\left(x,Q^2\right)\right] \tag{1.8}$$

where α_{em} denotes the electromagnetic coupling constant and the structure functions F_2 and F_L depend on the internal structure of the proton. They have to be determined experimentally since they currently cannot be calculated from first principles although non perturbative QCD methods based on lattice QCD exist [4]. The structure function F_2 is proportional to the sum of the cross sections for the exchange of longitudinally and transversely polarised photons whereas F_L only depends on the cross section for the exchange of longitudinally polarised photons. In the kinematic region of not too large y the contribution of the structure function F_L can be neglected and the cross section mainly depends on F_2. A compilation of the most precise F_2 measurements is shown in Figure 1.2 [5, 6, 7, 8, 9, 10, 11, 12]. The structure function F_2 is shown as a function of the photon virtuality Q^2 for different values of the scaling variable x. For values of $x \sim 0.2$ F_2 is independent of Q^2 which was first observed at SLAC [13, 14] and is known as *scaling*. For larger x values the structure functions decreases with increasing Q^2 whereas the opposite behaviour is observed for smaller values of x. The observation of scaling led to the advent of the quark–parton model [15, 16, 17, 18] which was refined by the observation of scaling violations [19] for large and small values of x. These effects have been successfully described in the framework of Quantum Chromo Dynamics (QCD) [20, 21, 22, 23].

[2]The reaction is assumed to be independent on the azimuthal angle and therefore this degree of freedom has not been considered.



Figure 1.2: *Compilation of the most precise F_2 measurements up to date in dependence of Q^2 for different values of x. The data in the region of low Q^2 and medium to large x are taken by fixed target experiments. These kinematic region has been extended by about 2 orders of magnitude in x and Q^2 by the HERA experiments H1 and ZEUS. The data are compared to next to leading order QCD fits which are able to reproduce the measurements in the whole kinematic range* [5, 6, 7, 8, 9, 10, 11, 12].

The Quark Parton Model

The observation of scaling at SLAC in the kinematic region $1 < Q^2 < 10\,\mathrm{GeV}^2$ for $x \simeq 0.2$ was interpreted by Bjorken and Paschos [15, 16] and by Feynman [17, 18] as being due to the partonic structure of the proton. In a Thompson–model of the proton in which the proton is an extended particle consisting of a continuous charge distribution a steeply falling structure function F_2 would be expected for all values of x as may be understood as follows. The wavelength of the virtual photon is proportional to the inverse of its virtuality $\lambda \sim 1/\sqrt{-q^2}$ which means that the photon probes smaller distances for larger Q^2 values and therefore it would be sensitive to smaller and smaller fractions of the total electric charge of the proton. Since the coupling of the photon is proportional to the charge this would lead to an F_2 falling with increasing values of Q^2 for all values of x.

In the quark–parton model the proton consists of three point-like partons which can be identified with the quarks introduced by Gell–Mann and Zweig [24, 25] to explain the spectroscopic hadron data. The proton is built from two up–quarks with fractional charge 2/3 and one down–quark with fractional charge -1/3. In this model the electron scatters elastically off one of the three quarks by single photon exchange. Since the time scale of the interaction is very small in comparison to interactions between the constituent quarks the proton can be treated for cross section calculations as an incoherent sum of these three quarks. This model leads to the observed scaling behaviour and the structure function F_2 may then be written as

$$F_2(x) = \sum_f e_f^2 x q_f(x) \tag{1.9}$$

where e_f is the charge of the struck quark and $q_f(x)$ are the quark density functions. In this model, x can be interpreted, in the infinite momentum frame, as the momentum fraction of the protons momentum carried by the struck quark.

By exploring a larger region in the x–Q^2 plane the violation of the scaling behaviour was observed as mentionend before. In addition it was found when summing up the momentum carried by the quarks that only a fraction of the total proton momentum is carried by these partons. These effects can be understood in the framework of Quantum Chromo Dynamics (QCD).

Quantum Chromo Dynamics (QCD)

Quantum Chromo Dynamics is the quantum field theory for strong interactions. With the large success of Quantum Electrodynamics (QED) in describing electromagnetic interactions at high energies it was recognised that renormisable quantum field theories are the most appropriate way to describe interactions of high energy particles. Until the advent of QCD a phenomenological theory the so called Regge theory existed which was able to describe *soft* hadron–hadron interactions (see chapter 1.2). With the observation of the substructure of hadrons a new theory for strong interactions became necessary.

In QCD the proton is built from quarks (as in the quark–parton model) which are spin 1/2 fermions. These are bound together by gluons which are the spin 1 gauge bosons of QCD, a non–abelian gauge theory invariant under SU(3) colour symmetry. In the framework of QCD colour corresponds to a new degree of freedom which takes the rôle of the electric charge in QED. It is carried by the quarks and in contrast to QED also by the gauge bosons, the gluons as a consequence of the non–abelian character of the theory. This leads to a self coupling of the gluons and hence to fundamental differences between QCD and QED.

Renormalisation and the Running Coupling

In perturbative QCD calculations divergent integrals appear which are treated by a renormalisation scheme and hence lead to an effective coupling constant $\alpha_s\left(\mu_r^2\right)$ depending on the energy scale μ_r^2. The requirement that the calculated cross sections are independent of the renormalisation scale μ_r^2 leads to the Renormalisation Group Equation (RGE) which in leading log. approximation (LLA) leads to the scale dependence of the coupling constant:

$$\alpha_s\left(\mu_r^2\right) = \frac{12\pi}{(33-2n_f)\ln\left(\mu_r^2/\Lambda_{QCD}^2\right)}. \tag{1.10}$$

Λ_{QCD} is a free parameter which has to be determined by experiments. It represents the lower limit of μ_r for which perturbative QCD calculations are expected to be predictive. n_f is the number of quark flavours with mass $m_q^2 < \mu_r^2$. In contrast to QED the effective strong coupling constant α_s is decreasing for larger values of μ_r^2 as consequense of the gluon self coupling.

The two extremes of small and large μ_r^2 which by the uncertainty principle correspond to large and small distances respectively, are called infrared slavery and asymptotic freedom. In the region of infrared slavery α_s is large and perturbative QCD calculations are not applicable. This long range effect is responsible for the confinement of partons in bound states, the hadrons. In the region of asymptotic freedom α_s is small and the perturbative expansion series (in powers of α_s) of QCD calculations is expected to converge fast which means that quarks can be treated as free particles.

In a pQCD calculation to a fixed order in α_s the cross section depends on the renormalisation scale μ_r^2 through the coupling constant $\alpha_s\left(\mu_r^2\right)$. When calculating up to all orders the terms involving μ_r^2 cancel and the cross section becomes independent of the renormalisation scale.

The Factorisation in DIS

The theorem of factorisation has been proven for hard scattering [26] and states that short range effects in the scattering amplitude, calculable in pQCD, can be separated from long range effects which are accounted to the parton density functions (pdf's). The proton structure function F_2 can be written as

$$F_2\left(x,Q^2\right) = \sum_{i=q,\bar{q},g}\int_x^1 d\xi f_i\left(\xi,\mu_r^2,\mu_f^2,\alpha_s\right)\cdot C_i^V\left(\frac{x}{\xi},\frac{Q^2}{\mu_r^2},\mu_f^2,\alpha_s\right). \tag{1.11}$$

In this decomposition μ_f^2 is the factorisation scale, f_i are the parton density functions and C_i^V are the coefficient functions. These describe the interaction of the exchanged boson V with a quark i (see Figure 1.3) which is calculable in perturbative QCD. The factorisation scale μ_f^2 defines the energy scale above which additional parton emissions from the quark are included in the perturbative QCD calculation. Long range effects which are not calculable in pQCD (e.g. parton emission with $k_T^2 < \mu_f^2$) are absorbed in the parton density functions f_i which therefore also become dependent on the factorisation scale μ_f^2. The parton density functions f_i can be interpreted in leading order calculations as probability to find a parton of type i for a certain value of x in the hadron. These are universal functions which only depend on the hadron type whereas the coefficient functions C_i^V are process dependent.



Figure 1.3: *a) Leading order diagram for the interaction of the virtual photon with a quark of the proton. b) In NLO calculations diagrams with e.g. additional parton emission have to be taken into account. The factorisation theorem states that parton emission with a large scale is taken into account in the scattering amplitude, calculable in pQCD whereas parton emission below the scale μ_f is absorbed in the parton density function which has to be determined experimentally.*

Parton Density Functions

Since the parton density functions cannot be calculated from first principles they have to be extracted from the experimental data although perturbative QCD calculations predict the evolution (e.g. in Q^2) of parton densities once they are known at at a certain starting point $\left(f_i\left(x, Q_0^2\right)\right)$. The proton structure function F_2 can be developed in a perturbation series in powers of α_s which contains terms $\alpha_s \ln\left(Q^2/Q_o^2\right)$, $\alpha_s \ln\left(1/x\right)$ and mixed terms $\alpha_s\left(Q^2/Q_o^2\right)\ln\left(1/x\right)$. Three different kind of evolution equations exist differing in the choice of terms added up from this perturbative expansion. The evolution of the parton density functions corresponds to a ladder diagram shown in Figure 1.4 and the summation of certain terms can be interpreted as ordering in either k_T or x of the rungs from the ladder (see e.g. [27]).

DGLAP Evolution and Operator Product Expansion

With the Dokshitzer, Gribov, Lipatov, Altarelli and Parisi (DGLAP) evolution equation [28, 29, 30, 31] the evolution of the parton density functions in dependence of Q^2 is predicted. In leading order the two coupled equations can be written as:

$$\frac{dq_i\left(x,Q^2\right)}{d\ln Q^2} = \frac{\alpha_s\left(Q^2\right)}{2\pi}\int_x^1 \frac{dz}{z}\left[\sum_j q_j\left(z,Q^2\right)P_{ij}\left(\frac{x}{z}\right) + g\left(z,Q^2\right)P_{ig}\left(\frac{x}{z}\right)\right], \tag{1.12}$$

$$\frac{dg\left(x,Q^2\right)}{d\ln Q^2} = \frac{\alpha_s\left(Q^2\right)}{2\pi}\int_x^1 \frac{dz}{z}\left[\sum_j q_j\left(z,Q^2\right)P_{gj}\left(\frac{x}{z}\right) + g\left(z,Q^2\right)P_{gg}\left(\frac{x}{z}\right)\right]. \tag{1.13}$$

Here $q_i\left(z,Q^2\right)$ and $g\left(z,Q^2\right)$ denote the quark and the gluon density functions. P_{ij} are the splitting functions (see Figure 1.5) which describe the parton emission calculable in perturbative QCD. Starting from a certain input density function $f_i\left(x,Q_0^2\right)$ the parton density is calculable for higher Q^2 values by solving the DGLAP evolution equations. This scheme corresponds to a strong k_T ordering $\left(k_{T,1} \ll k_{T,2} \ll ... \ll Q^2\right)$ of the subsequent parton emission as can be seen in Figure 1.4 whereas the ordering in x is only weak $(x_1 > x_2 > ... > x)$.



Figure 1.4: *In perturbative QCD the evolution of the structure function F_2 can be described by such a ladder diagram. In the DGLAP approach the subsequent parton emissions are strongly ordered in k_T $\left(k_{T,1} \ll k_{T,2} \ll ... \ll Q^2\right)$ and only weak ordering in x $(x_1 > x_2 > ... > x)$ is required. In the BFKL approach which is expected to be valid at very low x a strong ordering in x $(x_1 \gg x_2 \gg ... \gg x_i)$ is required whereas in k_T no ordering is present.*



Figure 1.5: *First order diagrams for three different splitting functions $P_{ij}(z)$.*

In a more formal approach the DGLAP evolution equations can be derived by the method of Operator Product Expansion (OPE) [32]. The matrix element of the reaction $ep \longrightarrow eX$ is proportional to the hadronic transition amplitude

$$M \sim < X|O|p > \tag{1.14}$$

where O is the operator which describes the transition from the initial proton $|p>$ to the hadronic final state $< X|$ (see Figure 1.1). The total cross section is proportional to the square of the matrix element

$$\sigma \sim |M|^2 \sim \sum_X < p|O'|X >< X|O|p > \tag{1.15}$$

which can be simplified when summing up all contributions of the hadronic final state X which leads to the relation

$$|M|^2 \sim < p|O'O|p > . \tag{1.16}$$

This method can be visualised in the quark–parton model by the diagram in Figure 1.6. The Operator Product Expansion consists of expanding the product $O'O$ in terms of quark and gluon fields where the leading diagram (Figure 1.6) can be interpreted as a if a quark is projected out



Figure 1.6: *left: leading order diagram for the matrix element $X|O|p>$; right: leading order diagram for the matrix element $<p|O'O|p>$.*

the proton which then interacts with the virtual photon and subsequently enters the proton again. In the context of the factorisation theorem the upper part of the diagram represents the hard scattering amplitude calculable in pQCD and the lower part is a representation for the parton densities. Note that the quark line between the vertices of the coupling gauge boson is on shell due to energy momentum conservation.

BFKL and CCFM Evolution

Several other approaches exist to model the structure function F_2 in regions where the DGLAP approximation does not hold. In the approach by Balitsky, Fadin, Kuraev and Lipatov (BFKL) [33, 34, 35] the terms in $\ln(1/x)$ of the perturbative expansion of F_2 are summed up which leads to a strong ordering in x in the ladder diagram (Figure 1.4) $(x_1 \gg x_2 \gg ... \gg x_i)$ The BFKL equation can be written in terms of the unintegrated gluon density $f_g\left(x, k_T^2\right)$ from which the ordinary gluon distribution can be calculated by integration over k_T. It thus predicts the evolution of F_2 as function of x.

The approach by Catani, Ciafaloni, Fiorani and Marchesini (CCFM) [36, 37, 38] gives the BFKL solution at low x and the DGLAP one at large Q^2. It is based on a strong angular ordering of subsequent parton emission.

Todays structure function measurements can be entirely described in the DGLAP scheme for $Q^2 \gg 1\,\mathrm{GeV}^2$ but there is indication that the DGLAP approach fails to describe certain exclusive measurements of the hadronic final state in ep collisions [39, 40] which might be the first sign for BFKL effects.

Parameterisations of Parton Density Functions

In order to model the structure function F_2 several parameterisations for parton densities exist where most of them are based on a polynomial ansatz (e.g. $xg = Ax^{\delta}(1-x)^{\eta}$) at a certain starting scale Q_0^2. The evolution to higher values of Q^2 is then performed by solving the DGLAP evolution equations and the free parameters of this ansatz are determined by a fit to the data.

The approach of Glück, Reya and Vogt (GRV) [41] is based on a dynamical model in which at a very low scale (chosen to be $Q_0^2 \simeq 0.34\,\mathrm{GeV}^2$), only valence like partons are assumed to exist. The parton density functions at higher values of Q^2 are then calculated by the DGLAP evolution equations.

Figure 1.7 shows the GRV parton density functions at a) the starting scale $Q^2 = \mu^2$ and b) after evolution to $Q^2 = 5\,\mathrm{GeV}^2$ [42]. It can be seen that the valence like gluon density very fastly evolves to low values of x which can be understood by gluon radiation from the valence partons which is resolved for larger values of Q^2. In Figure 1.7c) a fit to the data for the different parton flavours is shown [43] at the scale $Q^2 = 10\,\mathrm{GeV}^2$ which compares qualitatively with the GRV prediction.



Figure 1.7: *a) parton density functions in the GRV approach at the starting scale $Q^2 = \mu^2$; b) pdf's after evolution at $Q^2 = 5\,\mathrm{GeV}^2$; c) QCD fit of the pdf's at $Q^2 = 10\,\mathrm{GeV}^2$* [42, 43].

1.2 Diffractive Scattering

The investigation of hadron–hadron interactions at high energies has been a rich source of information about the strong interaction. Especially the study of events with low momentum transfer has been of great interest [44, 45, 46]. In the pre QCD era these kind of events have been successfully described in the framework of Regge-theory [47] which is based on the S-Matrix approach. In this model the hadron–hadron interactions are described by the exchange of meson and baryon trajectories. At high energies the pomeron trajectory, corresponding to a pseudo particle with the quantum numbers of the vacuum, exchange was proposed to explain the rise of cross sections at high energies. Measurements of diffractive events with jet production gave the first hint of the partonic structure of the pomeron.

Diffractive events have also been observed in deep inelastic electron–proton scattering at HERA where the pomeron is exchanged between the photon fluctuated into a hadronic state and the proton. The large range of the photon virtuality Q^2 available at HERA opened the possibility to measure precisely the partonic structure of the diffractive exchange.

In a QCD based model for the pomeron exchange, diffractive events were proposed to be described by a two gluon exchange model which can be best tested studying vector meson production. This model then leads consequently to the concept of generalised parton distributions which relates also to the partonic structure of the proton.

Diffractive Scattering in Soft Hadron–Hadron Interactions

Experimental Observations

In Figure 1.8 the cross section for elastic proton–proton scattering is plotted for different centre of mass energies as a function of $|t|$ where t is the squared momentum transfer between the two hadrons [46]. For low centre of mass energies a falling cross–section is observed where the slope gets steeper towards higher centre of mass energies. In addition a minimum and a maximum appear. This shape is similar to the diffraction pattern as observed when a plane wave of light passes a disc. The shape of the cross section down to the first minimum can be approximated



Figure 1.8: *Compilation of measurements of the differential cross section for elastic pp scattering as a function of $|t|$ for different values of the centre of mass energy* [46].

by an exponential function of the form[3]:

$$\frac{\mathrm{d}\sigma}{\mathrm{d}t} = \left(\frac{\mathrm{d}\sigma}{\mathrm{d}t}\right)\Bigg|_{t=0} e^{-b|t|} \tag{1.17}$$

with $b \sim 10\,\mathrm{GeV}^{-2}$.

The data can be interpreted in the spirit of the similarity between optical and hadronic diffraction [45]. In this picture one discusses the scattering of a fast hadron represented by a plane wave in the field of a hadron at rest. In optical diffraction the following formula holds for $kR \gg 1$:

$$\frac{I}{I_0} \simeq 1 - \frac{R^2}{4}k^2\theta^2 \tag{1.18}$$

which describes the fraction of observed intensity w.r.t. the initial intensity as a function of the diffraction angle θ, where R is the radius of the disc and k denotes the wave number of the incident light wave. If one relates the scattering angle θ to the momentum transfer t in hadron–hadron interactions, the wave number k to the momentum of the hadrons, and the radius of the disc R to the mass of the pion one can deduce a t-slope b of $b = 12.5\,\mathrm{GeV}^{-2}$ resulting in the right order of magnitude as measured.

Aspects of Regge Theory

The only theory to describe soft hadron–hadron interactions is Regge theory [47]. Apart from the fact that it was invented well before QCD, it still provides the only possibility to predict cross sections for these kind of reactions since perturbative QCD calculations are not applicable due to the absence of a hard scale in the process.

Regge theory was introduced in the framework of non–relativistic quantum mechanics and generalised the idea of Yukawa explaining particle interactions by the exchange of virtual particles. Later it was generalised to also describe relativistic particle interactions. It is based on S-Matrix theory which connects the wave function of the initial state $|i>$ with that of the final state $|f>$ by the relation:

$$|f>= S|i> . \tag{1.19}$$

The foundations of Regge–theory are the following axioms:

- The superposition principle states that a particle state ψ_γ can be decomposed linearly in the form $|\psi_\gamma >= \alpha|\psi_\alpha > +\beta|\psi_\beta >$ correctly taking into account the selection rules.
- The forces are required to be of short range. This is fulfilled for strong interactions since the interaction range is of the order of the inverse of the pion mass: $R \sim 1/m_\pi \sim 10^{-15}\,\mathrm{m}$.
- The Lorentz invariance principle states that the S-Matrix is a Lorentz scalar which itself only depends on Lorentz scalars.
- The unitarity requirement which corresponds to the conservation of probability states that $S^\dagger S = 1 = SS^\dagger$.
- Maximal analyticity of first order is required (see below).

[3]Note that in a more careful analysis the sum of two exponatial funtions has to be used [46]

- Generalisation of angular momentum in the complex plane is introduced (see below).

If one considers the reaction $A + B \longrightarrow C + D$. The Mandelstam variables s_{ij} are real numbers defined by relationes of the form $s_{ij} = (\pm P_i \pm P_j)^2$ with $i, j \in (A, B, C, D)$. Maximal analyticity of first order means that these variables can be extended into the complex plane which allows the use of complex analysis. The S-Matrix can now be written as a function of these complex numbers. In the limit of real numbers one gets back the S-Matrix for particle reactions in the physical region. A consequence of this axiom is the crossing symmetry which states that if the reaction $A + B \longrightarrow C + D$ is described by the variable $s = (P_A + P_B)^2$ and $t = (P_A - P_C)^2$ by the S–Matrix $S(s,t)$ one can calculate the amplitude[4] for the reaction $A + \bar{C} \longrightarrow D + \bar{B}$ by exchange of s and t which leads to the amplitude $S(t,s)$ as depicted in Figure 1.9.



Figure 1.9: *The two diagrams show the reactions $AB \longrightarrow CD$ and $A\bar{C} \longrightarrow \bar{B}D$ respectively. These processes share the same scattering amplitude. If the s-channel reaction a) can be described by the exchange of a particle in the t-channel then Regge theory predicts that the t-channel reaction b) is dominated by production of a resonance which subsequently decays into the final state. This prediction originates from the crossing symmetry being the consequence of one of the axioms of Regge theory.*

The solution of the non relativistic Schrödinger equation are eigenstates of the angular momentum with integer or half integer eigenvalues. In Regge theory the angular momentum is continued in the complex plane. The physical meaningful results are achieved by projecting the results on the real axis of the complex angular momentum plane.

If one considers the reaction $\pi^+\pi^- \longrightarrow p\bar{p}$ one can describe it by the fusion of the two pions to a ρ, ρ' or even higher excited states which than decay into the $p\bar{p}$ pair. This corresponds to the diagram in Figure 1.9 b with $A = \pi^+$, $\bar{C} = \pi^-$, $\bar{B} = \bar{p}$ and $D = p$. All intermediate states $(\rho, \rho', ...)$ must have the same quantum numbers except for the angular momentum J which only depends on the relative angular momentum of the two incident pions. The allowed states are hence a discrete series of states with integer spin. Regge theory states now that the scattering amplitude of the crossed channel $(\pi^+ p \longrightarrow \pi^+ p)$ can be calculated by exchanging the Mandelstam variables s and t in the scattering amplitude

$$S_{\pi^+p \longrightarrow \pi^+p}(s,t) = S_{\pi^+\pi^- \longrightarrow p\bar{p}}(t,s)\,. \tag{1.20}$$

It was observed that all intermediate states of the reaction $\pi^+\pi^- \longrightarrow p\bar{p}$ lie on a straight line the so called ρ–trajectory when plotting the spin of the physical states as function of the squared

[4]Note that the s and t channel are defined according to [47].

mass which is shown in Figure 1.10. The ρ–trajectory connects the s–channel ($t < 0$) and t–channel ($t > 0$) region which corresponds in this example to the reaction $\pi^+ p \longrightarrow \pi^+ p$ with $t = m^2$. When extending the trajectory into the s–channel region the value $\alpha(t) = Re(J)$ is used to predict cross sections for the crossed channel (see below) which agrees well with the observations. The trajectory depends on the intercept $\alpha(0) = \alpha_0$ and the slope α'

$$\alpha(t) = \alpha_0 + \alpha' \cdot t. \tag{1.21}$$

Figure 1.10 shows that the measurements of the ρ trajectory in the s–channel region agree well with the prediction when continuing the Regge trajectory from the t–channel region into the s–channel region. A large amount of data from hadron–hadron scattering can be explained by Regge theory using the known meson and baryon spectra as origin for Regge trajectories.



Figure 1.10: *The ρ trajectory: In the t–channel region the bound states (ρ, ρ_3) are shown. A straight line through these states is continued into the s–channel region where it is compared to data from hadron–hadron interactions.*

The total cross section for hadron–hadron interactions is connected to the imaginary part of the forward scattering amplitude $\mathcal{A}$ by the optical theorem which states

$$\sigma_{tot}(AB) \sim \frac{1}{s}\mathrm{Im}\mathcal{A}\,(AB \longrightarrow AB)\,(s, t = 0). \tag{1.22}$$

Hence this relation can be used to calculate total cross sections when knowing the scattering amplitude. Regge theory provides predictions e.g. for the energy dependence of total cross section:

$$\sigma_{tot} \sim s^{2\alpha(0)-2}, \tag{1.23}$$

where s is the centre of mass energy of the hadron–hadron interaction. The measurement of the energy dependence of the cross section therefore provides access to the intercept α_0 of the trajectory. When measuring the t–dependence of cross sections also the determination of the slope α' becomes possible (see Figure 1.10).

It was observed (see Figure 1.11) that the total cross section decreases with energy for low energies which is explained by Regge–trajectories originating from the known baryon and meson spectra whereas the rise of the cross section for high energies was not expected. To account for



Figure 1.11: *Compilation of measurements for total hadron–hadron cross sections as a function of the centre of mass energy* $\sqrt{s}$ *for different hadron types. a) proton–proton and proton–antiproton cross sections; b)* π^+p *and* π^-p *cross sections; c) photon–proton cross sections, where the photon can be treated as hadron in the vector meson dominance model (see text)* [48].

this rise the Pomeron trajectory was introduced which is characterised by the quantum numbers of the vacuum. From the rise of σ_{tot} for different reactions the Pomeron trajectory can be derived [48]:

$$\alpha(t) = 1.08 + 0.25 \cdot t. \tag{1.24}$$

It is measured in the s–channel region and a real particle in the t–channel regime is expected but not yet found. This particle is most probably glueball, i.e. a composite state of two valence gluons with the quantum numbers of the vacuum.

With introduction of the Pomeron trajectory also the observations discussed at the beginning of this section can be explained. The shrinkage of the diffractive peak can be predicted since Regge theory states:

$$b = b_0 + 2\alpha' ln \left(\frac{s}{s_0} \right) \tag{1.25}$$

and the double peak structure of the differential cross section as function of t at high energies (see Figure 1.8) can be explained by interference between single and multiple Pomeron exchange.

Following the idea by Ingelmann and Schlein [49] the structure of the vacuum exchange was investigated using diffractive events in presence of a hard scale suitable to resolve the partonic structure of the Pomeron. In proton–antiproton interactions this was done studying diffractive events with high transverse momentum jets. The measured cross section for this kind of events has been interpreted to be dominated by a large gluon content of the Pomeron [50, 51]. Following the same idea at HERA diffractive events have been investigated in the presence of a hard scale.

Diffraction in Electron–Proton Interactions

In the first years of HERA operation a widely unexpected large amount of diffractive events was observed [52, 53, 54, 55].

Inclusive Diffraction in Electron–Proton Interactions

Such events are identified by the presence of a large gap in rapidity η[5] in which no particle production is observed. The corresponding diagram for diffraction in electro–production is shown in Figure 1.12. The hadronic system Y propagates in the direction of the incoming proton while the system X is observed in the central part of the detector. Due to the high centre of mass energy in the photon–proton system it can be assumed that for the main part of the available phase space Pomeron exchange is dominant. In order to account for the new degrees of freedom for diffractive processes it is necessary to introduce the additional kinematic variables

$$M_X^2 = p_X^2, \qquad M_Y^2 = P_Y^2, \qquad t = (P - p_Y)^2,$$

$$x_{\mathbb{P}} = \frac{q \cdot (P - p_Y)}{q \cdot P}, \qquad \beta = \frac{x}{x_{\mathbb{P}}} = \frac{-q^2}{2q \cdot (P - P_Y)},$$

where M_X and M_Y are the invariant masses of the hadronic systems X and Y and t is the momentum transfer squared at the proton vertex. $x_{\mathbb{P}}$ can be interpreted as the longitudinal momentum carried by the exchanged Pomeron whereas β is the momentum fraction of the struck quark w.r.t. the pomeron momentum.

The cross section for diffractive DIS as depicted in Figure 1.12 can be written as a five fold differential cross section depending on: $x_{\mathbb{P}}$, β, Q^2, M_Y and t. Since M_Y and t cannot be measured at present one must integrate over these variables obtaining a three fold differential

[5] η is defined as $\eta = -\frac{1}{2} \ln \left(\tan \frac{\theta}{2} \right)$

cross section which is then related to the diffractive structure function $F_2^{D(3)}$ introduced similarly to the inclusive proton structure function F_2 [57]:

$$\frac{d^3\sigma_{ep\longrightarrow eXY}}{dx d\beta dQ^2} = \frac{4\pi\alpha_{em}^2}{\beta^2 Q^4}\left(1 - y + \frac{y^2}{2}\right) F_2^{D(3)}. \qquad (1.26)$$

Following the idea of Ingelmann and Schlein [49] the virtual photon which is emitted by the incoming electron can probe the structure of the Pomeron with different spatial resolution according to its virtuality Q^2. As factorisation was proven to hold for this process [56] the differential cross section can be analysed in terms of a hard scattering amplitude and parton density functions similarly to inclusive DIS. The structure function $F_2^D(3)$ was further assumed to factorise into a Pomeron flux and a Pomeron structure function which then was analysed in terms of parton distributions for the Pomeron. The QCD analysis indicated a large gluon component [57] which is in line with the results from jet production in hadron–hadron interactions [51].



Figure 1.12: *The diagram for diffractive electroproduction at HERA. The photon emitted by the incoming electron interacts with the Pomeron and resolves its structure. In the hadronic final state two systems X and Y are separated by a gap in rapidity with no hadron production.*

The diffractive exchange can be studied in more detail by selecting diffractive events with specific properties of the hadronic final state X (see Figure 1.12). Events with two high energetic particle jets have been studied [58] and also events with charmed mesons in the final state have been analysed [59] where a hard scale is provided either by the transverse momentum of the jets or the mass of the charm quark to resolve the partonic structure of the Pomeron. The results of these analysis are consistent with the QCD analysis of the diffractive structure function.

Vector Meson Production

In order to test the diffractive exchange more exclusive events have been analysed where the hadronic system X consists only of a vector meson. Vector meson production ($ep \longrightarrow e\mathrm{VM}p$) has been extensively studied at HERA for different types of vector mesons and also in the different kinematic regimes [60, 61, 62, 63, 64, 65, 66, 67, 68]. The data can be interpreted in terms of Regge theory in connection with the vector meson dominance model (VDM) [69, 70] as well as in the framework of perturbative QCD [71]. It therefore opens the possibility to study the applicability of both approaches and allows a separation of effects calculable in perturbative QCD from soft physics observed in hadron–hadron interactions.

In the VDM the photon emitted by the incoming electron fluctuates into a vector meson which then scatters diffractively on the proton as depicted in Figure 1.13. In the VDM the production of vector mesons with the same quantum numbers as the photon is described (i.e. ρ, ω, ϕ, ...) since the Pomeron is characterised by the quantum numbers of the vacuum.



Figure 1.13: *Diagram for vector meson production in the vector meson dominance model in conjunction with Regge theory. The incoming electron emits a photon (real or virtual) which fluctuates into a hadronic system, the vector meson, which subsequently interacts with the proton by pomeron exchange. Only the production of vector mesons with the same quantum numbers as the photon are described within this model since the pomeron carries the quantum numbers of the vacuum.*

Figure 1.14 shows the measured cross section for vector meson production in photoproduction $\left(Q^2 \simeq 0\right)$ as a function of W, the photon–proton centre of mass energy. For the production of light vector mesons (ρ, ϕ and ω) the rise of the cross section at high W can be described as before in the framework of Regge theory by the exchange of the known Pomeron trajectory. The photoproduction of light vector mesons belongs therefore to the class of soft diffractive interactions. In the presence of a hard scale μ^2 as is the case for J/ψ production with $\mu^2 = m_c^2$, where the charm mass provides the hard scale, the rise of the cross section is steeper than predicted by Regge theory.

For light vector meson production a hard scale can be introduced by selecting events in the kinematic domain $Q^2 >$ few GeV^2. Here a similar behaviour can be observed indicating, that the rise of the cross section for high values of W becomes steeper when Q^2 rises. This can be seen in Figure 1.15 where the slope δ of the cross section as function of W $\left(\sigma \sim W^\delta\right)$ is shown for different values of Q^2. For low Q^2 the rise of the cross section is comparable to the rise predicted by Regge theory. For large values of Q^2 the rise is significantly larger. These kind of interactions are called hard diffractive interactions and expected to be calculable within the framework of perturbative QCD since the presence of a hard scale ensures convergence of the perturbation series.

In summary this means that in the presence of a hard scale Regge theory is not able anymore



Figure 1.14: *Measured cross sections for the production of various types of vector mesons in photoproduction* $(Q^2 \simeq 0)$ *as a function of* W, *the photon proton centre of mass energy, is shown* [60, 61, 62, 63, 64, 65, 66, 67, 68]. *For light VM production the rise at large values of* W *is comparable to that predicted by pomeron exchange. For heavy vector meson production (i.e.* J/ψ*) the observed rise is steeper than for light vector mesons.*



Figure 1.15: *The slope of the cross section* δ *as function of* W $(\sigma \sim W^{\delta})$ *for diffractive production of* ρ *mesons is shown as function of the photon virtuality* Q^2 [72]. *For low values of* Q^2 *the fitted slope* δ *is comparable to the one predicted by pomeron exchange. For large values of* Q^2 *the slope is significantly larger.*

to describe diffractive processes. Due to the presence of a hard scale these kind of interactions are expected to be calculable in the framework of perturbative QCD. The investigation of these processes can be used to study the transition from soft to hard processes by varying the hard scale either by variation of the mass of the final vector meson, by the investigation of events in different regions of the photon virtuality Q^2 or by studying events with a large momentum transfer t at the proton vertex.

The Two Gluon Exchange Model

The two gluon exchange model [73, 74] was invented as model for the Pomeron and has been applied to the diffractive production of vector mesons. Figure 1.16 shows one of the diagrams to be calculated for this process. The virtual photon emitted by the incoming electron fluctuates in a $q\bar{q}$ system which then interacts with the proton by the exchange of two gluons. The quark–antiquark system subsequently recombines to the vector meson. The exchange of two gluons is the simplest way to establish the quantum numbers of the vacuum. The two quarks which build the final state vector meson have to be very close in phase space to avoid a break up before the VM has been formed. This condition is fulfilled in case of the two gluon exchange since the quark–antiquark pair stems from the photon and a small transverse momentum between the partons is possible. When exchanging two quarks which then couple directly to the incoming photon and the outgoing vector meson a large transverse momentum is enforced, due to energy momentum conservation, between the two partons leading to a suppression of this contribution for the reaction at low x.



Figure 1.16: *Diagram for the production of vector mesons in the framework of QCD. The photon which is emitted by the incoming electron splits in a quark–antiquark pair which then interacts with the proton by the exchange of two gluons. After the interactions the quark–antiquark pair recombines to a vector meson. The diagram is fully calculable in perturbative QCD except the transition from the quark–antiquark pair to the final vector meson which has to be modelled.*

The factorisation theorem for this process has been proven [75] and states that the upper part of the diagram (see Figure 1.16) can be calculated in perturbative QCD and the non perturbative effects may be absorbed into parton density functions similar to ordinary DIS. In the common calculations for vector meson production the two gluons are approximated by the square of the gluon density as derived from DIS. These kind of calculations have been successfully applied to predict the rise of the cross section for diffractive J/ψ production as function of W.

The approximation of the two gluon exchange by the square of the ordinary gluon density holds for low Q^2 and not too heavy vector mesons. In the domain of large Q^2 or for the production of heavy vector mesons (e.g. Υ) it is expected that an additional effect has to be taken into account when calculating cross sections. In case of photoproduction $\left(Q^2 \simeq 0\right)$ of heavy vector mesons the almost real photon interacting with the proton is transformed to a vector meson with the mass squared M^2_{VM} (see Figure 1.16). Energy momentum conservation implies a longitudinal momentum transfer from the lower to the upper part of the diagram and hence $x_1 \neq x_2$ where x_1 and x_2 are the longitudinal momentum fractions of the two exchanged gluons. The difference in longitudinal momentum of the two participating partons is:

$$x_1 - x_2 = \frac{Q^2 + M_V^2/4}{W^2} \tag{1.27}$$

where M_V is the mass of the produced vector meson. In order to account for this effect the ordinary parton densities have been extended to the so called generalised parton distributions (GPD's) which become important for the calculations of cross sections for large values of Q^2, heavy vector mesons and for low values of W. In particular for the production of Υ mesons GPD's have to be used to explain the measured cross sections (see below).

1.3 Generalised Parton Distributions

Generalised Parton distributions (GPD's) [6] are functions which generalise and interpolate between different types of functions characterising the structure of the proton (nucleons). In particular the parton density functions as measured in DIS do not provide enough information needed for cross section predictions of certain reactions (e.g. vector meson production). Elastic form factors, unpolarised and polarised parton distributions can be derived from GPD's when integrating over or taking boundary values of these functions.

The concept of GPD's arose from three different theoretical studies[7]. Geyer et al. studied the relation between the Altarelli–Parisi evolution for parton distributions and the Brodsky–Lepage evolution for meson wave functions [77, 78, 79, 80]. The introduced interpolating functions are essentially what is nowadays called GPD's. Jain and Ralston studied hard processes involving hadron helicity flip in terms of an off–diagonal transition amplitude [81]. These can be used to derive elastic form factors. Finally Ji introduced off–forward parton distributions to describe the spin structure of the nucleon [82, 83]. He also proposed the measurement of Deeply Virtual Compton Scattering to extract these new distributions from the data [83].

Definition of GPD's

In this section the formal definition of GPD's is reviewed following the nomenclature of Ji [82, 83]. Additional definitions exist (e.g. [84]) which are basically equivalent and transformation formulae have been provided to switch between different nomenclatures.

[6]The term *Generalised Parton Distribution* is nowadays commonly used and comprises all names introduced in the past which originate from the different usage of these functions. Historically the terms: *off–diagonal*, *non–diagonal*, *off–forward* and *non–forward parton distribution* are used as well as *skewed parton distributions*.

[7]For a historical review see [76].

In section 1 it has been discussed that the standard parton distributions are obtained in the framework of Operator Product Expansion (OPE) from the squared proton wave function for all partonic configurations containing a parton with specified polarisation and longitudinal momentum x. After application of the factorisation theorem the hadronic matrix element $< p|O'O|p >$ can be identified with the parton density function which is interpreted as the probability for finding such a parton in the proton. Figure 1.17a) shows the graphical representation of this matrix element.

In contrast to this GPD's are defined as hadronic matrix elements of two unequal hadronic wave functions representing the incoming and outgoing nucleon (e.g. for VM production) $< p'|O|p >$ where $|p>$ denotes the wave function of the incoming proton and $|p'>$ represents the outgoing proton [85]. In Figure 1.17b) and c) the corresponding diagrams are shown.

In comparison to standard DIS two kinematic regimes exist, the so called DGLAP region (Figure 1.17b) and the ERBL region (Figure 1.17c). In the DGLAP region ($|x| > \xi$) the matrix element is interpreted as emission of a parton with longitudinal momentum $(x + \xi) \cdot p$ where p is the momentum of the incident proton and subsequent absorption of a parton with longitudinal momentum $(x - \xi) \cdot p$. For the ERBL region ($|x| < \xi$) which has no counterpart to standard DIS the matrix element is interpreted as emission of a di–parton system with longitudinal momenta $(x + \xi) \cdot p$ and $(-x + \xi) \cdot p$. In case of quarks the two parton system consists of a quark–antiquark pair.

The standard parton density functions are defined on the cross section level whereas the GPD's are functions defined on the amplitude level, i.e. when calculating cross sections (e.g. for vector meson production) the GPD's enter the calculation of the scattering amplitude which then is squared to achieve the cross section expression.



Figure 1.17: *Diagrams representing the hadronic matrix elements: a) for standard DIS and b) and c) for the generalised parton distribution. In contrast to parton densities in standard DIS the GPD's are defined by hadronic matrix elements of unequal hadronic wave functions. One distinguishes between the DGLAP b) and ERBL c) region.*

The generalised parton distributions can formally be defined by Fourier transforms of the hadronic matrix elements:

$$\begin{aligned} \int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle P'|\overline{\psi}(-\lambda n)\gamma^\mu \psi(\lambda n)|P\rangle &= H(x,\xi,t)\ \overline{u}(P')\gamma^\mu u(P) \\ &+ E(x,\xi,t)\ \overline{u}(P')\frac{i\sigma^{\mu\nu}\Delta_\nu}{2M}u(P), \end{aligned} \tag{1.28}$$

$$\begin{aligned} \int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle P'|\overline{\psi}(-\lambda n)\gamma^\mu \gamma_5 \psi(\lambda n)|P\rangle &= \widetilde{H}(x,\xi,t)\ \overline{u}(P')\gamma^\mu \gamma_5 u(P) \\ &+ \widetilde{E}(x,\xi,t)\ \overline{u}(P')\frac{\Delta^\mu \gamma_5}{2M}u(P) \end{aligned} \tag{1.29}$$

where $|P\rangle$ and $\langle P'|$ represent the quantum numbers of the incoming and outgoing proton respectively (including differences for the spin state). The operators $\overline{\psi}(-\lambda n)\gamma^\mu\psi(\lambda n)$ and $\overline{\psi}(-\lambda n)\gamma^\mu\gamma_5\psi(\lambda n)$ select the parton with certain properties from the hadronic wave functions. Δ^μ is defined as $\Delta^\mu = p'^\mu - p^\mu$ in the infinite momentum frame, $t = \Delta^2$, $u(P)$ and $\overline{u}(P')$ denote the Dirac spinors of the hadron. An examination of the helicity structure of the scattering amplitude shows that there are exactly four independent generalised parton distributions. The functions H, E, $\widetilde{H}$ and $\widetilde{E}$ are defined for the different quark flavours as well as for the gluon.

The reference system is chosen such that the initial and final nucleon have longitudinal momenta $(1+\xi)p^\mu$ and $(1-\xi)p^\mu$, and the outgoing and incoming partons carry the longitudinal momentum $(x+\xi)p^\mu$ and $(x-\xi)p^\mu$, respectively. Due to energy momentum conservation x and ξ are restricted to

$$0 < \xi < \frac{\sqrt{-t}}{\sqrt{M^2 - t/4}}, \tag{1.30}$$

$$-1 < x < 1. \tag{1.31}$$

The distributions H_q and E_q are summed over the quark helicities, whereas $\widetilde{H_q}$ and $\widetilde{E_q}$ involve the difference between right and left handed quarks and therefore contain information about the spin structure of the proton. The chiral-even distributions H and $\widetilde{H}$ survive in the forward limit, (i.e. $H(x,\xi,t) \neq 0$ and $E(x,\xi,t) \neq 0$ for $\xi = 0$ and $t = 0$) in which the nucleon helicity is conserved, while the chiral-odd distributions E and $\widetilde{E}$ allow for the possibility of a nucleon helicity flip. The change of the nucleon helicity in case of helicity conservation for the quarks is possible when orbital momentum is transferred by the two partons.

The evolution equations for the generalised parton distributions are modified DGLAP evolution equations in the DGLAP regime ($|x| > \xi$) whereas in the ERBL region ($|x| < \xi$) the Brodsky–Lepage evolution equations are used which originate from the study of evolution equations for meson wave functions[92, 93].

Sum rules and boundary conditions for GPD's

Several sum rules and boundary conditions for GPD's exist, thus drastically restricting the possibilities of models for GPD's.

The Nucleon Spin Structure

The spin of the nucleon can be decomposed in components which are carried by the quarks and by the gluons [82]:

$$\overrightarrow{J}_{\mathrm{N}} = \overrightarrow{J}_q + \overrightarrow{J}_g. \tag{1.32}$$

It can now be shown that:

$$J_{q,g} = \frac{1}{2}\Big(A_{q,g}(0) + B_{q,g}(0)\Big) \tag{1.33}$$

with

$$\int_{-1}^{1} dx\; x\Big(H(x,\xi,t) + E(x,\xi,t)\Big) = A(t) + B(t) \tag{1.34}$$

where the integral over the functions H and E has to be evaluated for the quark and for the gluon distributions. The ξ–dependence drops out when calculating the integral.

The Parton Density Functions

The generalised parton distributions contain information of both the ordinary forward parton distributions and electromagnetic nucleon form factors. In the limit $\Delta^\mu \longrightarrow 0$ one restores the ordinary quark and the quark helicity distributions $q(x)$ and $\Delta q(x)$ for $x > 0$:

$$H(x,0,0) = q(x), \quad \widetilde{H}(x,0,0) = \Delta q(x). \tag{1.35}$$

The Elastic Form Factors

The first moment of the generalised parton distributions is related to the nucleon form factors:

$$\int_{-1}^{1} dx H(x,\xi,t) = F_1(t) \ , \tag{1.36}$$

$$\int_{-1}^{1} dx E(x,\xi,t) = F_2(t) \ , \tag{1.37}$$

$$\int_{-1}^{1} dx \widetilde{H}(x,\xi,t) = G_A(t) \ , \tag{1.38}$$

$$\int_{-1}^{1} dx \widetilde{E}(x,\xi,t) = G_P(t) \ , \tag{1.39}$$

where F_1 and F_2 are the Dirac and Pauli form factors which can be related by linear combinations to the electric and magnetic form factors G_E and G_M. G_A and G_P represent the axial and pseudo-scalar form factors of the nucleon.

Models for GPD's

In this section the model for GPD's from Radyushkin based on Double Distributions will be outlined [80, 84, 86, 87]. Other approaches based on the MIT bag model [76], hadronic light cone wave functions [85] or the chiral quark soliton model [88, 89] are not discussed.

In addition to the boundary conditions discussed for GPD's they have to fulfil a polynomiality condition which is a non trivial consequence from Lorentz invariance. It states that the Mellin moments of the order N of the GPD should by polynomials of the maximal order $N+1$:

$$\int_{-1}^{1} \mathrm{d}x \ x^N \ H^q(x,\xi) = h_0^{q(N)} + h_2^{q(N)}\xi^2 + ... + h_{N+1}^{q(N)}\xi^{N+1}, \tag{1.40}$$

$$\int_{-1}^{1} \mathrm{d}x \ x^N \ E^q(x,\xi) = e_0^{q(N)} + e_2^{q(N)}\xi^2 + ... + e_{N+1}^{q(N)}\xi^{N+1}. \tag{1.41}$$

This condition is satisfied when deriving GPD's from Double Distributions (DD) which are two dimensional functions $F(x',y')$ defined in a certain region of the x'–y' plane. The GPD is then calculated from the Double Distribution by the reduction formula:

$$H^q(x,\xi) = \int_{-1}^{1} \mathrm{d}x' \int_{-1+|x'|}^{1-|x'|} \mathrm{d}y' \ \delta\left(x' + \xi y' - x\right) F\left(x',y'\right). \tag{1.42}$$

In this ansatz the t–dependence has already been factorised out

$$H^q(x,\xi,t) = H^q(x,\xi) \cdot G(t) \tag{1.43}$$

which is a reasonable for HERA kinematics $\left(x_{\mathrm{Bj}} < 10^{-2}\right)$ [83].

Using this ansatz the polynomiality condition is automatically fulfilled independent of the actual form of the double distribution $F\left(x', y'\right)$. In the model from Radyushkin the Double Distribution is further factorised in two functions:

$$F\left(x', y'\right) = \pi^{q}\left(x', y'\right) f^{q}\left(x'\right) \tag{1.44}$$

where $\pi^{q}\left(x', y'\right)$ is the profile function and $f^{q}\left(x'\right)$ is the ordinary parton distribution. This choice restaurates the ordinary parton density for $\xi = 0$ as required by equation 1.35. In order to satisfy this condition the profile function $\pi\left(x', y'\right)$, which is the only unknown quantity in this model, has to fulfil the following integral equation:

$$\int_{-1+|x'|}^{1-|x'|} \mathrm{d}y' \pi\left(x', y'\right) = 1. \tag{1.45}$$

The shape of the profile function then determines the effect of the GPD's on the cross section calculation.

Since the Double Distribution fulfil the polynomial condition except that the highest term in the polynomial is always 0 an additional term has been introduced by Polyakov and Weiss [90] which accounts for the highest term in the polynomial.

This ansatz is chosen for a certain scale Q_0^2. Once the GPD's are determined at $Q^2 = Q_0^2$ they can be evolved in the DGLAP region ($|x| > \xi$) similar to the DGLAP evolution equations with a kernel that depends on the longitudinal momentum transfer ξ and in the ERBL region ($|x| < \xi$) evolve according to equations similar to the ERBL evolution equations [92, 93] for meson distribution amplitudes.

GPD's in Practice

The first comparison of GPD based calculations with measurements has been done in an analysis by Martin, Ryskin and Teubner when they calculated the cross section for photoproduction of Υ mesons [91]. In Figure 1.18 the cross section measurements from H1 and ZEUS are compared to two different calculations, one based on the leading order expression and the other one including an enhancement factor originating from GPD's. The leading log QCD calculation leads to the simple formula:

$$\left.\frac{d\sigma}{dt}\left(\gamma p \rightarrow V p\right)\right|_{t=0} = \frac{\alpha_s^2 \Gamma_{ee}^V}{3\alpha M_V^5}\, 16\pi^3 \left[xg\left(x, \frac{M_V^2}{4}\right)\right]^2 \tag{1.46}$$

where Γ_{ee}^V is the partial width of the $V \rightarrow ee$ decay, α_s is the QCD coupling constant, $\alpha = 1/137$ is the QED coupling constant and $g(x, \mu^2)$ is the gluon density measured at $x = M_V^2/4W^2$ and the scale $\mu = m_Q \simeq M_V/2$. The LO prediction is well below the measurements although the statistical errors are still quite large. This calculation has been corrected for four different effects.

Relativistic corrections and NLO QCD corrections lead to a change of the predicted cross section by 7 % and 20 % respectively. Formula 1.46 accounts only for the imaginary part of the scattering amplitude. When restoring the real part of the QCD amplitude the cross section has to be enhanced by about 50 %. The largest correction originates from effects when using generalised parton distributions. It has been found to account for an enhancement of about a factor of 2. When comparing the measured cross section with the corrected calculation a better description of the data can be observed.



Figure 1.18: *The measurements for photoproduction of Υ mesons are compared to LO calculations (dashed line) and to a calculation including NLO effects as well as corrections due to the real part of the scattering amplitude and a correction due to the GPD's which account for the non forward kinematic situation* [91].

Chapter 2

Deeply Virtual Compton Scattering

2.1 Deeply Virtual Compton Scattering

Deeply Virtual Compton Scattering (DVCS) is defined as the elastic scattering of a virtual photon off a proton with a real photon in the final state. In electron–proton interactions the photon with a virtuality $Q^2 >$ few GeV^2 is emitted by the incoming electron leading to the reaction $ep \longrightarrow e\gamma p$ as depicted in Figure 2.1 a). This process is investigated to study the structure of the proton; in particular it provides access to the generalised parton distributions as introduced in Chapter 1.

It has been proven that in presence of a hard scale ($Q^2 \to \infty$) the DVCS scattering amplitude factorises in a hard scattering coefficient, calculable in perturbative QCD and a soft part which is absorbed in generalised parton distributions [94]. Figure 2.2 shows examples of the leading (a)



Figure 2.1: *The contributing diagrams for the reaction $ep \longrightarrow e\gamma p$: a) diagram for the Deeply Virtual Compton Scattering process; b) and c) diagrams for the Bethe–Heitler process.*

and next to leading order (b) diagrams of the DVCS process in a QCD approach. In the leading order process the virtual photon scatters off a quark originating from the proton. After emission of the real photon the quark is reabsorbed by the proton. In the NLO diagram the virtual photon interacts via a quark loop with two gluons from the proton. The real photon is also emitted from the quark loop. In contrast to DVCS the contribution of quark exchange to the vector meson production process is suppressed w.r.t. the two–gluon exchange since the two quarks which form the vector meson have to be very close in phase space to form a stable meson. However this situation is suppressed for HERA kinematics (large W). For low energies (e.g. HERMES kinematics) the quark exchange also has to be taken into account when considering vector meson production.

DVCS calculations are expected to be more reliable than predictions for vector meson production due to the point–like coupling of the final state photon to the quark line. In vector meson production the VM wave function has to be modelled which describes the transition from the quark–antiquark system to the vector meson.



Figure 2.2: *Examples for diagrams of the DVCS process in leading and next to leading order QCD: a) The leading order diagram is the so called handbag diagram where the virtual photon interacts with a quark from a proton with subsequent emission of the real photon. b) In the next to leading order diagram the photon interacts with two gluons from the proton by a quark loop from which also the real photon is emitted.*

In addition to the DVCS process the Bethe–Heitler (BH) process where the real photon is emitted from the electron line contributes to the reaction $ep \longrightarrow e\gamma p$ (see Figure 2.1b) and c)). It leads to a direct contributions to the cross section and a contribution due to the interference of the two processes. The interference term provides the key to extract GPD's from $ep \longrightarrow e\gamma p$ data (see below).

It was found [95] that the DVCS and BH amplitude exhibit a different spin dependence which leads to a non vanishing ϕ dependence of the interference term, where ϕ is defined as the angle between the electron scattering plane and the photon production plane calculated in the virtual photon–proton centre of mass system (see Figure 2.3). The amplitude for the DVCS process shows a ϕ dependence according to:

$$\mathcal{A}_{DVCS} \sim e^{-i\lambda\phi}, \tag{2.1}$$

where λ is the helicity of the intermediate virtual photon γ^*. In contrast to this the BH process exhibits a ϕ dependence following:

$$\mathcal{A}_{BH} \sim e^{-2i\lambda'\phi}, \tag{2.2}$$

where λ' is the helicity of the final state photon. For the pure DVCS and BH contributions to the cross section the ϕ dependences cancel when going from the amplitude to the cross section expression, whereas for the interference term it does not. When evaluating the interference term for unpolarised ep scattering it turns out that its contribution to the cross section is proportional to $\cos\phi$. This can be exploited by an asymmetry measurement in which one gets access to the real part of the QCD amplitude (see below). Moreover, when studying ep scattering with polarised beams, different ϕ dependencies appear for different beam polarisations which all can be used to access different terms of the scattering amplitude. Once the different contributions to the scattering amplitude are measured the generalised parton distributions can be determined.



Figure 2.3: *The angle ϕ is defined as the angle between the scattering plane of the electron and the production plane of the real photon in the centre of mass system of the virtual photon and the incoming proton. The production plane of the real photon is defined by its vector and the virtual photon proton axis.*

2.2 Determination of GPD's from the DVCS Measurement

Although this analysis does not aim for a determination of the generalised parton distribution from the measurement, the method will be shortly outlined. As already discussed a measurement of the azimuthal angle asymmetry provides access to the real part of the QCD DVCS amplitude.

The azimuthal angle asymmetry is defined by the formula

$$A = \frac{\int_{-\pi/2}^{\pi/2} \mathrm{d}\phi \, \frac{\mathrm{d}\sigma^{ep\longrightarrow e\gamma p}}{\mathrm{d}\phi} - \int_{\pi/2}^{3\pi/2} \mathrm{d}\phi \, \frac{\mathrm{d}\sigma^{ep\longrightarrow e\gamma p}}{\mathrm{d}\phi}}{\int_0^{2\pi} \mathrm{d}\phi \, \frac{\mathrm{d}\sigma^{ep\longrightarrow e\gamma p}}{\mathrm{d}\phi}}, \tag{2.3}$$

where ϕ is the angle between the electron scattering plane and the photon production plane in the photon–proton centre of mass system. It is proportional to the real part of the DVCS amplitude which can be written in the factorised expression as

$$\mathrm{Re}\, \mathcal{A}\left(x, Q^2\right) = \int_{-1+x}^{1} \frac{\mathrm{d}x_1}{x_1} \mathrm{Re}\, C_i\left(x/x_1, Q^2\right) f_i\left(x_1, x, Q^2\right) \tag{2.4}$$

where Re C_i denotes the real part of the hard scattering coefficient and f_i are the generalised parton distributions for quarks and gluons. x_1 denotes the momentum of one of the exchanged partons and x denotes the Bjorken scaling variable.

In order to solve the deconvolution problem a method was proposed [96] based on the ansatz of a polynomial expansion for the GPD at a starting scale Q_0^2. The free parameters are obtained from fits to the data. When evolving the GPD to higher values of Q^2 the validity of this ansatz can be checked by comparing the results to data at these Q^2 values. Although this method is very convenient to apply, it has the disadvantage that the extraction of the GPD is model dependent.

2.3 Theoretical Predictions

The Model from Frankfurt, Freund and Strikman

Similar to ordinary DIS it is possible to calculate the Q^2 evolution for the scattering amplitude of Deeply Virtual Compton Scattering above a normalisation point $Q_0^2 \sim$ few GeV^2 [97]. Below this point the scattering amplitude is dominated by non–perturbative effects which prevents the usage of pQCD.

It was found ([98] and references therein) that the aligned jet model provides a reasonable description of the structure function F_2. When using the Gribov dispersion relation the ratio R of the imaginary parts for the DIS amplitude (in terms of the photon–proton amplitude) and the DVCS amplitude can be calculated as:

$$R = \frac{\mathrm{Im}\mathcal{A}\left(\gamma^* + p \longrightarrow \gamma^* + p\right)_{t=0}}{\mathrm{Im}\mathcal{A}\left(\gamma^* + p \longrightarrow \gamma + p\right)_{t=0}} \approx 0.5 \tag{2.5}$$

for $Q_0^2 \approx 2 - 3\,\mathrm{GeV}^2$. The DVCS process can now be calculated by a sum of soft and hard contributions where the soft contribution has been taken from this aligned jet model estimate and the hard contribution can be calculated in the framework of QCD evolution equations. This leads to the following expression for the evolution of the DVCS scattering amplitude[1]:

$$\begin{aligned} \mathrm{Im}\mathcal{A}\left(x, Q^2, Q_0^2\right) &= \mathrm{Im}\mathcal{A}\left(x, Q_0^2\right) \\ &+ 4\pi^2\alpha_{em} \int_{Q_0^2}^{Q^2} \frac{\mathrm{d}Q'^2}{Q'^2} \int_x^1 \frac{\mathrm{d}x_1}{x_1} \left[P_{qg}\left(x/x_1, \Delta/x_1\right) g\left(x_1, x_2, Q'^2\right)\right. \\ &\qquad \left. + P_{qq}\left(x/x_1, \Delta/x_1\right) q\left(x_1, x_2, Q'^2\right)\right] \end{aligned} \tag{2.6}$$

where P_{qg} and P_{qq} are the evolution kernels for generalised parton distributions. Using this relation the value for R has been estimated for the kinematic range accessible in this analysis as $R \approx 0.55$, almost independent on Q^2 and x [97]. The real part of the DVCS amplitude has been calculated using a dispersion relation:

$$\eta = \frac{\mathrm{Re}\mathcal{A}}{\mathrm{Im}\mathcal{A}} = \frac{\pi}{2} \frac{\mathrm{d}\ln\left(x\mathrm{Im}\mathcal{A}\right)}{\mathrm{d}\ln\frac{1}{x}}. \tag{2.7}$$

These relationes can now be used to calculate the complete cross section for the DVCS process. The imaginary part of the DIS amplitude $\mathrm{Im}\mathcal{A}\left(\gamma^* + p \longrightarrow \gamma^* + p\right)_{t=0}$ has been connected via the optical theorem to the DIS cross section which can be expressed in terms of the proton structure function F_2, such that F_2 enters the cross section expression (see below). The value of R accounts for the effect of the generalised parton densities.

The BH process is calculable within the framework of QED using the electro–magnetic form factors G_E and G_M. The complete differential cross section for the reaction $ep \longrightarrow e\gamma p$ is decomposed into the sum of three terms; the DVCS contribution, the BH term and the interference term.

$$\frac{d\sigma^{ep\rightarrow e\gamma p}}{dxdydtd\phi} = \frac{d\sigma^{\mathrm{DVCS}}}{dxdydtd\phi} + \frac{d\sigma^{\mathrm{BH}}}{dxdydtd\phi} + \frac{d\sigma^{\mathrm{Int}}}{dxdydtd\phi}, \tag{2.8}$$

[1]Note the different convention w.r.t. chapter 1.3.

$$\frac{d\sigma^{\mathrm{DVCS}}}{dxdydtd\phi} = \frac{\pi\alpha^3 s\left(1+(1-y)^2\right)}{4R^2Q^6}e^{-b|t|}F_2^2\left(x,Q^2\right)\left(1+\eta^2\right), \tag{2.9}$$

$$\frac{d\sigma^{\mathrm{BH}}}{dxdydtd\phi} = \frac{\alpha^3 sy^2\left(1+(1-y)^2\right)}{\pi|t|Q^4\left(1-y\right)}\frac{G_E^2(t)+\frac{|t|}{4m_p^2}G_M^2(t)}{1+\frac{|t|}{4m_p^2}}, \tag{2.10}$$

$$\frac{d\sigma^{\mathrm{Int}}}{dxdydtd\phi} = \pm\frac{\eta\alpha^3 sy\left(1+(1-y)^2\right)}{2RQ^5\sqrt{|t|(1-y)}}e^{-b|t|/2}F_2\left(x,Q^2\right)\frac{G_E(t)+\frac{|t|}{4m_p^2}G_M(t)}{1+\frac{|t|}{4m_p^2}}\cos\phi. \tag{2.11}$$

Note that the prediction for the DVCS process is valid at $t = t_{\min}$ where t is the momentum transfer at the proton vertex. The model assumes the t-dependence to follow an exponential function: $\frac{d\sigma}{dt} \sim e^{-b|t|}$ which is an ansatz that has been successfully applied to diffractive processes in the region of low $|t|$ (e.g. ρ vector meson production [60]). The interference term shows the $\cos\phi$ dependence which has been discussed already at the beginning of this chapter. The sign of the interference term depends on the polarity of the lepton beam.

Colour Dipole Models

General Properties of Colour Dipole Models

The Colour dipole model of diffraction [99] provides a simple unified picture of diffractive processes. It is able to reproduce the inclusive diffractive structure function $F_2^{D(3)}$ as well as exclusive reactions like vector meson production. Recently this model has been used to predict cross sections for the DVCS process. It is based on the assumption of a factorisation of the reaction in three subprocesses which are well separated in time. In the proton rest frame the incoming virtual photon fluctuates into a quark antiquark pair well before the interaction with the proton. The formation time for the dipole is $\tau \sim 1/M_p x$ which for HERA kinematics ($x \leq 0.01$) is much longer than the typical time scale for strong interactions $\tau \sim 1/m_\pi$. The subsequent formation of the hadronic final state (i.e. the photon in case of DVCS) from the scattered dipole is again much longer than the interaction time with the target. This justifies the basic assumption of the dipole model that the cross section for the interaction of the dipole is independent of how it is formed. The following factorisation formula for the scattering amplitude $\mathcal{A}$, also visualised in Figure 2.4, can be derived

$$\mathcal{A}\left(\gamma^* p \longrightarrow \gamma p\right) = \int \psi_{\gamma^*}^{in}\sigma_{\mathrm{d}}\psi_{\gamma}^{out}, \tag{2.12}$$

where the incoming and outgoing photon are denoted by the wave functions $\psi_{\gamma^*}^{in}$ and ψ_{γ}^{out}, respectively. It is integrated over all dipole sizes R and all momentum fractions z of the quark from the quark–antiquark dipole (see below). The cross section σ_{d} describes the interaction of the dipole with the proton. The colour dipole is assumed to be unchanged during the interaction. The different types of dipole models differ very strongly in their assumptions and the method used to derive the dipole cross section. In general the models make assumptions on the decomposition of dipole cross section; free parameters are derived from fits to the data. There exists no model which is able to derive the dipole cross section from first principles. Although the photon wave function is in principle calculable within perturbation theory (at least in leading order) some theories modify it to account for confinement effects for large dipoles.



Figure 2.4: *In colour dipole models the incoming virtual photon fluctuates in a quark–antiquark system with radius R which then interacts with the incoming proton. The dipole leaves unchanged during the interaction and subsequently recombines to a real photon.*

The dipole models have been constructed such that they incorporate soft and hard contributions to the cross section. The transverse momentum k_T of the quark–antiquark system is given by

$$k_T^2 \simeq z(1-z)Q^2 + m_q^2 \tag{2.13}$$

where z determines the longitudinal momentum fraction carried by the quark and m_q is the quark mass. This can be related by the uncertainty principle to the radial distance R between the quark and the antiquark. Thus at low Q^2 the reaction is dominated by the interaction of large dipoles with the proton while small dipoles dominate at large Q^2.

The dipole models presented here are valid for $t = t_{\text{min}}$. The cross section as a function of t is calculated by assuming an exponential t dependence

$$\frac{d\sigma}{dt} \sim e^{-b|t|}. \tag{2.14}$$

The Model from Donnachie and Dosch

In a model by Donnachie and Dosch [100] the dipole approach is coupled with the concept of soft and hard pomeron exchange. The scattering amplitude is written as

$$\mathcal{A}(s, t=0) = 2\pi \int_0^1 \mathrm{d}z \int \mathrm{d}R\, R\, \rho_\gamma^\lambda(Q_1^2, Q_2^2, R, z)\, \sigma_{\mathrm{d}}(R) \tag{2.15}$$

where z is the longitudinal momentum fraction of the incident photon carried by the quark and R is the diameter of the quark-antiquark dipole. The dipole cross section $\sigma_{\mathrm{d}}(R)$ has been evaluated in a non–perturbative model and has two free parameters which have been determined by fits to $p\bar{p}$ and pp cross section data for which the nucleons have been decomposed in a quark-diquark system and thus can be treated as dipoles. The quark antiquark overlap density of photons $\rho_\gamma^\lambda(Q_1^2, Q_2^2, R, z)$, the product of the incoming and outgoing photon wave functions, for which Q_1^2 and Q_2^2 denotes the virtualities of the incoming and outgoing photon and λ is their helicity, has been calculated in lowest order perturbation theory.

The energy dependence is included by multiplying the cross section with the factor $(W/W_0)^{2\epsilon_{\text{soft}}}$ where $1 + \epsilon_{soft}$ is the intercept of the pomeron trajectory with $\epsilon_{\text{soft}} \sim 0.08$ and $W_0 = 20\,\text{GeV}$. To account for the stronger energy dependence observed in reactions with a hard scale, the two pomeron approach [101] was included in the model such that small dipoles predominantly interact by the exchange of the hard pomeron component whereas large dipoles interact via the soft pomeron component. Finally Donnachie and Dosch obtain the expression for the cross section at $t = t_{\rm min}$.

$$\begin{aligned}
&\frac{d\sigma}{dt}(Q^2, W, t = t_{min}) = \\
&\frac{1}{16\pi}\Big(r_{\rm soft}(Q^2)(W/W_0)^{0.16} + r_{\rm hard}(Q^2)(W/W_0)^{0.84}\Big)^2 \times 311\ \mu{\rm b/GeV}^2
\end{aligned} \tag{2.16}$$

where $W_0 = 20$ GeV and

$$\begin{aligned}
r_{\rm soft}(Q^2) &= \Big(5.33 - 1.33 \cdot e^{(-4Q^2/Q_0^2)} + 5.37\, Q^2/Q_0^2\Big)^{-1}, \\
r_{\rm hard}(Q^2) &= \Big(49.42 - 7.65 \cdot e^{(-4Q^2/Q_0^2)} + 4.94\, Q^2/Q_0^2\Big)^{-1}
\end{aligned} \tag{2.17}$$

with $Q_0^2 = 1$ GeV^2.

The Model of Forshaw, Kerley and Shaw

A model by Forshaw, Kerley and Shaw (FKS) [102, 103] assumes that the dipole cross section depends solely on the properties of the dipole proton system itself, independent of the photon virtuality Q^2, which means that it depends only on the photon–proton centre of mass energy W and on the size of the dipole R. The idea is to extract information on the dipole cross section by assuming a reasonable flexible form to fit the diffractive structure function and the total photon–proton cross section. The cross section is decomposed into two terms to account for soft and hard contributions

$$\sigma_{\rm d}\left(W^2, R\right) = \sigma_{\rm d\ soft}\left(W^2, R\right) + \sigma_{\rm d\ hard}\left(W^2, R\right). \tag{2.18}$$

The functions $\sigma_{\rm d\ soft}\left(W^2, R\right)$ and $\sigma_{\rm d\ hard}\left(W^2, R\right)$ have been chosen such that for small dipoles the hard contribution dominates while at low Q^2 $\sigma_{\rm d\ soft}\left(W^2, R\right)$ contributes most.

The wave functions of the incoming and outgoing photon have been calculated in leading order perturbation theory and multiplied by an enhancement factor to increase contributions from large dipoles. This procedure is justified by the fact that for small $Q^2 < 4m_q^2$ the wavefunction becomes sensitive to the quark mass and that confinement effects are expected to become sizable. The quark mass has been chosen such that $m_q^2 = 0.08\,{\rm GeV}^2$ which corresponds to the constituent quark mass.

The Model of McDermott, Frankfurt, Guzey and Strikman

In another model of McDermott, Frankfurt, Guzey and Strikman (MFGS) [102, 104] the dipole cross section $\sigma_{\rm d}\left(Q^2, W, R\right)$ depends on both, the virtuality of the photon Q^2 and the photon proton centre of mass energy W. It is divided into three parts depending on the dipole size R. For small dipoles $R < R_C$ the cross section is directly related to the LO gluon distribution of the proton

$$\sigma_{\rm d}\left(x, R < R_C\right) = \frac{\pi^2 R^2}{3}\alpha_s\left(\bar{Q}^2\right) xg\left(x', \bar{Q}^2\right), \tag{2.19}$$

where x' and $\bar{Q}^2$ are functions of x and Q^2 depending on the dipole size R. The underlying model for this cross section is the two–gluon–exchange model discussed earlier[2]. The upper bound R_C depends on Q^2 and is $R_C = 0.246\,\mathrm{fm}$ for $Q^2 = 2.6\,\mathrm{GeV}^2$. For large dipoles above the pion radius $R > R_\pi = 0.65$ fm the cross section has been modelled by the function

$$\sigma_\mathrm{d}\left(R > R_\pi\right) = \sigma\left(\pi p\right) \frac{3R^2}{2R^2 + R_\pi} \left(\frac{x_0}{x}\right)^{0.08} \tag{2.20}$$

independent on Q^2 where $x_0 = 0.01$ has been chosen such that at $R = R_\pi$ σ_d matches the total pion proton cross section $\sigma\left(\pi p\right) = 24$ mb. In the intermediate region $R_C < R < R_\pi$ the dipole cross section is interpolated linearly.

This model focuses on the region of small dipoles and is hence only expected to be valid for $Q^2 > 1$ GeV^2 where the contribution from large dipoles is expected to be negligible. For the actual predictions of the cross section a generalised gluon distribution has been used.

Predictions Based on Generalised Parton Distributions

All models discussed above depend on different assumptions how hard and soft effects contribute to the DVCS cross section. Freund and McDermott presented [105, 106, 107] the first complete QCD calculation based on generalised parton densities. The used GPD is based on the model of Radyushkin (see chapter 1.3) which uses Double Distributions to derive them. The profile function $\pi^i\left(x', y'\right)$ has been chosen as:

$$\pi^i\left(x', y'\right) = \frac{3}{4} \frac{\left(1 - |x'|\right)^2 - y'^2}{\left(1 - |x'|\right)^3} \tag{2.21}$$

which leads in combination with the ordinary parton distributions (GRV [42] and MRST' [108] have been used) to predictions for the DVCS amplitude. It was calculated in leading order and next to leading order QCD by calculating the coefficient function and solving the evolution equations.

Depending on the actually used parton distribution large differences (factor 2–3) on the amplitude level can be observed which illustrates the high sensitivity of the cross section on effects due to the skewedness effects. These prediction do not yet exist on cross section level which would be a crucial test for GPD based models.

2.4 MC Programs

The final state of the reaction $ep \longrightarrow e\gamma p$ consists of an electron, a photon and a proton, where the electron and photon are detected with the H1 detector[3]. In order to extract the cross section from the measured event rate, MC simulations are performed using event generators as input. They generate the four vectors for the final state particles of the interactions which are then subjected to a detailed detector simulation. Six different MC programs have been used to study the signal as well as different background contributions.

[2] Note that the dipole cross section σ_d enters the amplitude (Equation 2.12). When calculating the cross section it depends quadratically on the dipole cross section and the gluon density as naively expected for the two gluon exchange model.

[3] For a detailed discussion of the analysis strategy see chapter 4.1.

TINTIN

The event generator TINTIN, developed in the framework of this thesis (for a detailed description see appendix A), is a program which generates the four vectors of the final state particles for the reaction $ep \longrightarrow e\gamma p$ where the kinematic variables are generated according to the predicted cross section by the calculations of Frankfurt, Freund and Strikman (see chapter 2.3). The program is able to generate events for the BH process, the DVCS process and the sum of the BH and DVCS process taking the interference term into account[4]. The most dominant QED radiative correction, the emission of a real photon from the incoming electron is implemented in the program.

COMPTON 2.0

The MC program COMPTON 2.0 [110, 111, 112] is used to generate events for the Bethe–Heitler process. It generates the elastic contribution as well as inelastic events. The program uses an exact calculation based on helicity amplitudes where the coupling to the proton is given by the electromagnetic form factors (G_E and G_M) for the elastic contributions. The inelastic contribution is divided into two regimes the resonance region ($M_X < 1.8\,\mathrm{GeV}$) and the inelastic continuum ($M_X > 1.8\,\mathrm{GeV}$) where M_X is the invariant mass of the hadronic final state. For the resonance region the three proton resonances $\Delta(1232)$, $N^*(1520)$ and $N^*(1688)$ have been taken into account and the corresponding cross sections are derived from measurements. In the region of the inelastic continuum the cross section has been taken from early F_2 measurements. As will be discussed in chapter 4 the cross section simulation of the inelastic Bethe–Heitler events has been modified according to a revised set of experimental data. The emission of a real photon from the initial electron which is expected to be the dominant higher order QED effect is implemented.

DIFFVM

The MC program DIFFVM [113] was designed to simulate the process of diffractive vector meson production $ep \longrightarrow e$ VM X where VM denotes the produced vector meson and X the hadronic final state, separated by a rapidity gap from the vector meson. The predictions of the program are based on the vector meson dominance model (VDM) and Regge theory as discussed in section 1.2. The virtual photon emitted by the incoming electron fluctuates into a vector meson which subsequently scatters off the proton. For the hadronic system X the resonant region as well as the inelastic continuum are taken into account. The program generates the four vectors of the final state particles including the decay products of the produced vector meson.

The DIFFVM program generates different vector meson channels, i.e. all vector mesons which have the same quantum numbers as the virtual photon (ω, ρ, ϕ ...). It can be used to generate elastic vector meson production as well as resonance production and the inelastic continuum.

[4]The free parameters have been set in the event generation to the following values: the t–slope parameter $b = 7\,\mathrm{GeV}^{-2}$, the phase of the QCD amplitude, $\eta_{\mathrm{QCD}} = 1 - \frac{\pi}{2}\left(0.176 + 0.33 \ln Q^2\right)$ [109] and the sensitivity to the skewedness of the parton densities $R = 0.55$ [97]. The proton structure function F_2 as extracted from the H1 data has been used [5].

GRAPE

The event generator GRAPE [114] is used to simulate the reaction $e^+p \longrightarrow e^+e^-e^+p$. The predicted cross section is based on a calculation of the full matrix element taking all contributing diagrams into account. In the kinematic range relevant for this analysis the cross section is dominated by the two photon process where the photon emitted by the incoming positron interacts with a photon emitted from the proton by the production of an electron positron pair. Using the GRAPE MC program it is again possible to generate the full final state for the elastic and the inelastic reaction.

DJANGO and HERWIG

The MC program DJANGO ([115] and references therein) is a standard DIS event generator which is used for the generation of inclusive deep inelastic scattering events. It is a combination of the programs LEPTO, HERACLES, ARIADNE and JETSET where the program LEPTO calculates the DIS Matrix element. It is interfaced to HERACLES for calculation of the QED radiative corrections. The program ARIADNE simulates the subsequent emission of partons with a large transverse momentum. The results of this program are then interfaced to JETSET which performs the simulation of the fragmention based on the Lund String model. The program produces the four vectors of all final state particles.

The HERWIG program [116] is also a standard DIS event generator similar to the DJANGO program. The basic difference between the two programs relevant for this analysis is the use of a different fragmentation scheme. In contrast to the Lund fragmentation scheme which is used in DJANGO the HERWIG program is based on the cluster fragmentation.

Chapter 3

HERA and H1

In this chapter the HERA[1] storage ring and the H1 experiment are introduced. After a brief description of the HERA machine, an overview of the H1 detector in its 1997 setup is given, concentrating on the components which were used for this analysis.

3.1 The Electron–Proton Collider HERA



Figure 3.1: *The electron proton collider HERA with its pre–accelerators.*

The HERA collider is located at the DESY[2] laboratory in Hamburg/Germany. It is the first electron–proton[3] collider ever built. The beam energies are 27.5 GeV for the electrons and 820 GeV for the protons respectively. This leads to an energy in the centre of mass system of

[1] **H**adron **E**lektron **R**ing **A**nlage
[2] **D**eutsches **E**lektronen **SY**nchrotron
[3] In the 1997 data taking period HERA accelerated positrons instead of electrons.

$\sqrt{s} \simeq 300\,\mathrm{GeV}$, which is more than 10 times higher than in formerly performed fixed target experiments. The two beam lines are located in the same tunnel with a circumference of 6.3 km. The electron ring is equipped with dipole magnets with a field strength of 0.17 T. The proton ring consists of superconducting dipole magnets with a field strength of 4.7 T. The proton energy is limited by the field strength of the bending magnets while the maximum electron energy is determined by the power of the cavities which accelerate the electrons and compensate for the energy loss due to synchrotron radiation. The energy loss of protons due to synchrotron radiation can be neglected due to their high mass.

Figure 3.1 shows an overview of HERA and its pre–accelerators. The electrons are provided by a 500 MeV linear accelerator which are then ramped up in DESY II (to 7 GeV) and PETRA (to 12 GeV). After injection into HERA they are brought to their final beam energy. The protons are produced by passing H^- ions with an energy of 50 MeV through a stripping foil where the ions loose their two electrons. The protons are collected in bunches and accelerated in DESY III (to 7.5 GeV) and in PETRA (to 40 GeV) and afterwards injected into HERA and accelerated to their final beam energy. Both the electrons and the protons are stored in bunches of 10^{10} to 10^{11} particles which circulate in opposite direction in the HERA storage ring. The time between two bunch crossings is 96 ns which leads to a total number of 220 bunches for both the electron and the proton ring. They are filled such that typically 175 of these bunches collide and the additional bunches are filled either for the electrons, the protons or none of them. The lifetime of the proton beam is about 100 hours, whereas the lifetime of the electron beam is about 10 hours which leads to a typical duration of 12 hours for a luminosity fill. By the year 2000 an electron current of 40 mA and a proton current of 90 mA was routiniously achieved leading to peak luminosity of $1.5 * 10^{31}\,\mathrm{cm}^{-2}\mathrm{s}^{-1}$ which corresponds to the design value.

The electron and the proton beam are brought to collisions at two interaction points. Around these the experiments H1 and ZEUS are constructed and take data since 1992. There are two additional fixed target experiments built at the HERA collider. The HERMES detector is a spectrometer which takes data since 1995 and studies interactions of polarised electrons from the electron beam with a polarised gas target (H_2, D, He). The aim of this experiment is to study the spin structure of the proton and related processes. The HERA–B experiment was commissioned in 2000 and is a forward spectrometer which uses an internal wire target to produce interactions between protons from the Halo of the proton beam and nucleons of the target wire. The aim of this experiment is to study CP–violation in the system of neutral B–mesons. This experiment serves also as a charm factory, which opens the possibility to study QCD with heavy quarks at high energies.

Figure 3.2 shows the time structure of a typical proton bunch in the HERA storage ring [117]. A threefold structure is observed where the main bunch is accompanied by two so called satellite bunches 5 ns before and after the main peak which are typically a factor 10 to 100 smaller. The interval of 5 ns between the main and the satellite bunch originates from the properties of the high frequency system of the HERA accelerating units. The frequency of the accelerating system is 200 MHz leading to a stable point in the seperatrix each 5 ns. The distribution of the proton bunches is very broad in the PETRA accelerating system. When transmitting them to the HERA ring about 7 % of the protons do not enter the main bunch but one of the satellite bunches. In the present analysis one cannot distinguish whether the final state particles originate from an interaction of an electron with a proton from the main or the satellite bunch. Therefore the effect of satellite bunch interactions has to be taken into account when analysing the data.

Since September 2000 the data taking has been stopped and the HERA accelerator and the



Figure 3.2: *The bunch structure of protons in the HERA storage ring is subdivided into three parts: The main bunch is accompanied by two satellite bunches* 5 ns *before and after the main peak* [117]. *The sidebands originating from protons outside stable points in the separatrix are a factor 1000 smaller w.r.t. the main bunch.*

experiments undergo major upgrades. Superconducting quadrupole magnets are built and installed inside the H1 and ZEUS detectors, which enable a better focussing of the beams in the interaction region and therefore lead to a luminosity 4–5 times higher than before.

3.2 The H1 Experiment

The H1 detector [118] is a multi–purpose apparatus to study the final state particles from electron–proton scattering. It is located in the hall north of HERA. In order to study the wide range of the HERA physics program the detector has to satisfy several basic requirements. In neutral current events the kinematic variables are reconstructed from the properties of the scattered electron which requires a good calorimetric measurement for all scattering angles. The electron is scattered in the backward direction[4] for small values of the photon virtuality $Q^2 < 100\,\mathrm{GeV}^2$, whereas for larger values of Q^2 the electron tends to be scattered in the central or forward direction. Furthermore it is essential that the detector is hermetic in order to measure the momentum of all final state particles to determine the missing momentum which is needed to identify charged current interactions where the neutrino leaves the detector unseen.

Further instrumentation is used to measure muons and particles which leave the main detector

[4]The backward direction is defined w.r.t. the incoming proton beam.