

## Guest editorial:

### POSSIBILITIES AND LIMITATIONS OF INTRAVITAL IMAGING

Reham Hassan

Forensic Medicine and Toxicology Department, Faculty of Veterinary Medicine,  
South Valley University, Qena, Egypt; [Reham\\_Hassan@vet.svu.edu.eg](mailto:Reham_Hassan@vet.svu.edu.eg)

<http://dx.doi.org/10.17179/excli2016-863>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

Recently, Raymond Reif and colleagues from the Leibniz Research Center in Dortmund have published a technical article about two-photon based intravital imaging (Reif et al., 2016). The strength of this article is the outstanding quality of the presented videos, which show organs of living mice, namely liver, kidney and intestine, with a resolution of approximately 200 nm. Key features of the established imaging technique are two-photon lasers with a broad wavelength spectrum, long-distance objectives with high numerical aperture, a sensitive detector and an optimized anesthesia (Reif et al., 2016). Under these conditions the technique can be applied to multiple organs to study transport and elimination of xenobiotics and endogenous compounds. A fundamental prerequisite of intravital imaging is the use of appropriate fluorophore-coupled compounds and fluorescent reporter mice. The authors used this method to image and quantify the flux of fluorescent bile salts and drugs from the sinusoids of the liver through the Dissé space into hepatocytes and finally into bile canaliculi. In the kidney it was possible to analyze glomerular filtration into the Bowman's capsule followed by transport into the proximal and then to the distal tubules. After bolus injections of fluorescent compounds even the passage time from proximal to distal tubules via the Henle loop could be determined. Finally, the same approach was used to image intestinal crypts

and study the transfer of drugs from capillaries to lymph vessels.

The presented intravital technique has major implications for three fields of research. First, pharmacokinetic modelling can be improved by intravital imaging. PBPK modelling is of high relevance for simulation of compound concentrations in blood or organs (Golubovskaya et al., 2015; Stamy et al., 2015; Widera, 2015; Ghallab, 2015) as well as for interspecies extrapolation (Thiel et al., 2015). However, current PBPK models consider individual organs as single compartments (Schug et al., 2013; Mielke et al., 2011). Using intravital microscopy it became clear that sub-compartments of tissues may accumulate specific compounds. Integrating these dynamics into PBPK models may further improve simulations. Second, establishment of in vitro systems represents a cutting-edge topic (Ghallab and Bolt, 2014; Godoy et al., 2013; Krug et al., 2013), particularly in the fields of cell systems for hepatotoxicity (Reif et al., 2015; Böttger et al., 2015; Grinberg et al., 2014), nephrotoxicity (Wassermann et al., 2013; Valente et al., 2012; Buhrke et al., 2015) and intestinal transport (Niu et al., 2015). It is important to be aware of similarities but also differences of these cultivated cells to their in vivo counterparts. Two-photon imaging can be used to study compound transport in vitro and in vivo by precisely the same method. Third, spatio-tem-

poral modelling has become a well-established technique in systems biology (Drasdo et al., 2014a, b; Ghallab et al., 2016; Hammad et al., 2014). A basic principle of spatio-temporal modelling is that tissues are reconstructed in a way that the position of each cell in a three-dimensional space is known; next further processes of interest, e.g. metabolic models, can be programmed into each individual cell to simulate the consequences at the tissue level (Hoehme et al., 2010; Schliess et al., 2014). Two-photon based imaging represents an excellent technique to validate predictions from spatio-temporal tissue models.

In conclusion, the two-photon based imaging technique presented by Reif and colleagues will support spatio-temporal modelling studies of physiological and pathophysiological processes in all organs of interest.

## REFERENCES

- Böttger J, Arnold K, Thiel C, Rennert C, Aleithe S, Hofmann U, et al. RNAi in murine hepatocytes: the agony of choice--a study of the influence of lipid-based transfection reagents on hepatocyte metabolism. *Arch Toxicol.* 2015;89:1579-88.
- Buhrke T, Frenzel F, Kuhlmann J, Lampen A. 2-Chloro-1,3-propanediol (2-MCPD) and its fatty acid esters: cytotoxicity, metabolism, and transport by human intestinal Caco-2 cells. *Arch Toxicol.* 2015;89:2243-51.
- Drasdo D, Bode J, Dahmen U, Dirsch O, Dooley S, Gebhardt R, et al. The virtual liver: state of the art and future perspectives. *Arch Toxicol.* 2014a;88:2071-5.
- Drasdo D, Hoehme S, Hengstler JG. How predictive quantitative modelling of tissue organisation can inform liver disease pathogenesis. *J Hepatol.* 2014b;61:951-6.
- Ghallab A. Systems toxicology. *EXCLI J.* 2015;14:1267-9.
- Ghallab A, Bolt HM. In vitro systems: current limitations and future perspectives. *Arch Toxicol.* 2014;88:2085-7.
- Ghallab A, Cellière G, Henkel SG, Driesch D, Hoehme S, Hofmann U, et al. Model-guided identification of a therapeutic strategy to reduce hyperammonemia in liver diseases. *J Hepatol.* 2016;64:860-71.
- Godoy P, Hewitt NJ, Albrecht U, Andersen ME, Ansari N, Bhattacharya S, et al. Recent advances in 2D and 3D in vitro systems using primary hepatocytes, alternative hepatocyte sources and non-parenchymal liver cells and their use in investigating mechanisms of hepatotoxicity, cell signaling and ADME. *Arch Toxicol.* 2013;87:1315-530.
- Golubovskaya V, Curtin L, Groman A, Sexton S, Cance WG. In vivo toxicity, metabolism and pharmacokinetic properties of FAK inhibitor 14 or Y15 (1, 2, 4, 5-benzenetetramine tetrahydrochloride). *Arch Toxicol.* 2015;89:1095-101.
- Grinberg M, Stöber RM, Edlund K, Rempel E, Godoy P, Reif R, et al. Toxicogenomics directory of chemically exposed human hepatocytes. *Arch Toxicol.* 2014;88:2261-87.
- Hammad S, Hoehme S, Friebel A, von Recklinghausen I, Othman A, Begher-Tibbe B, et al. Protocols for staining of bile canaliculi and sinusoidal networks of human, mouse and pig livers, three-dimensional reconstruction and quantification of tissue microarchitecture by image processing and analysis. *Arch Toxicol.* 2014;88:1161-83.
- Hoehme S, Brulport M, Bauer A, Bedawy E, Schormann W, Hermes M, et al. Prediction and validation of cell alignment along microvessels as order principle to restore tissue architecture in liver regeneration. *Proc Natl Acad Sci USA.* 2010;107:10371-6.
- Krug AK, Kolde R, Gaspar JA, Rempel E, Balmer NV, Meganathan K, et al. Human embryonic stem cell-derived test systems for developmental neurotoxicity: a transcriptomics approach. *Arch Toxicol.* 2013; 87:123-43.
- Mielke H, Anger LT, Schug M, Hengstler JG, Stahlmann R, Gundert-Remy U. A physiologically based toxicokinetic modelling approach to predict relevant concentrations for in vitro testing. *Arch Toxicol.* 2011;85:555-63.
- Niu X, de Graaf IA, Langelaar-Makkinje M, Horvato-vich P, Groothuis GM. Diclofenac toxicity in human intestine ex vivo is not related to the formation of intestinal metabolites. *Arch Toxicol.* 2015;89:107-19.
- Reif R, Karlsson J, Günther G, Beattie L, Wrangborg D, Hammad S, et al. Bile canaliculi dynamics in hepatocyte sandwich cultures. *Arch Toxicol.* 2015; 89:1861-70.

Reif R, Ghallab A, Beattie L, Günther G, Kuepfer L, Kaye PM, et al. In vivo imaging of systemic transport and elimination of xenobiotics and endogenous molecules in mice. *Arch Toxicol.* 2016 Dec 20. doi: 10.1007/s00204-016-1906-5. [Epub ahead of print]

Schliess F, Hoehme S, Henkel SG, Ghallab A, Driesch D, Böttger J, et al. Integrated metabolic spatial-temporal model for the prediction of ammonia detoxification during liver damage and regeneration. *Hepatology.* 2014;60:2040-51.

Schug M, Stöber R, Heise T, Mielke H, Gundert-Remy U, Godoy P, et al. Pharmacokinetics explain in vivo/in vitro discrepancies of carcinogen-induced gene expression alterations in rat liver and cultivated hepatocytes. *Arch Toxicol.* 2013;87:337-45.

Stamyr K, Mörk AK, Johanson G. Physiologically based pharmacokinetic modeling of hydrogen cyanide levels in human breath. *Arch Toxicol.* 2015;89:1287-96.

Thiel C, Schneckener S, Krauss M, Ghallab A, Hofmann U, Kanacher T, et al. A systematic evaluation of the use of physiologically based pharmacokinetic modeling for cross-species extrapolation. *J Pharm Sci.* 2015;104:191-206.

Valente MJ, Henrique R, Vilas-Boas V, Silva R, Bastos Mde L, Carvalho F, et al. Cocaine-induced kidney toxicity: an in vitro study using primary cultured human proximal tubular epithelial cells. *Arch Toxicol.* 2012;86:249-61.

Wassermann L, Halwachs S, Baumann D, Schaefer I, Seibel P, Honscha W. Assessment of ABCG2-mediated transport of xenobiotics across the blood-milk barrier of dairy animals using a new MDCKII in vitro model. *Arch Toxicol.* 2013;87:1671-82.

Widera A. Highlight report: Interspecies extrapolation by physiologically based pharmacokinetic modeling. *EXCLI J.* 2015;14:1014-6.